1966 cultures of meningococci isolated from patients in different areas of England and Northern Ireland have been sent to the Manchester Public Health Laboratory for serotyping and for testing for sensitivity to sulphonamides. From 1966 to the end of April 1972, we have typed 342 cultures isolated from blood or CSF: 27 strains were group A (8%), 185 group B (54%), 113 group C (33%), one strain each group X and Y, and 15 were untypable (4%). The majority of untypable strains (13/15) were received before 1970 and had been stored for long periods before typing. During the same period 370 cultures were tested for sensitivity to sodium sulphadiazine. The majority of cultures 319 (86%) were inhibited by 0.16 mg% sodium sulphadiazine, and a further 18 (5%) by 0.32 mg%. Thirty-three strains (9%) had a minimal inhibitory concentration of 0.64 mg% or greater, and these we have classed as resistant; 12 of these (3%) had a minimal inhibitory concentration of 0.64–1 mg%, and 21 (6%) a minimal inhibitory concentration of 5 mg% or more. The proportion of resistant strains isolated recently appears to be increasing—22% of those isolated in the first four months of 1972 compared with 7% of those in 1971. Resistant strains were isolated from different areas of England and Northern Ireland. The proportion of resistant strains amongst different serotypes varies—6% of group B, 8% of group C, and 44% of group A strains. These findings and the report of sulphonamide-resistant strains in Scotland (Fallon, 1971) clearly indicate that sulphonamides alone cannot be relied upon for the treatment of meningococcal infection in this country, and the efficacy of sulphonamides for chemoprophylaxis may be affected by the presence of sulphonamide-resistant strains.

Reference

Fallon, R. J. (1971). *Brit. med. J.*, 2, 403.

Meningococcal Infection: Serological Studies D. M. JONES (*Public Health Laboratory, Withington Hospital, Manchester*) The recrudescence of interest in meningococcal disease, particularly in the United States, has resulted in much progress being made towards the understanding of immunity in this disease (Artenstein *et al.* 1971). A variety of serological techniques have been used to study the antibody response in the carrier state, in disease, and after administration of the recently developed meningococcal polysaccharide vaccines. The indirect haemagglutination technique has been found to be very sensitive, and to detect type specific antibody. This method has been used to investigate the normal antibody responses in carriers and cases in a small outbreak of meningitis in a school in the north west

of England. With this basic information it has been possible to assess situations where multiple cases occur in one family and where there may be a deficiency in immunity. The occurrence of a community outbreak of group B meningococcal meningitis in Bolton and district during 1970 and 1971 enabled the incidence of antibody in the general population of epidemic and non-epidemic areas to be compared. The incidence of antibody was found to increase with age but no significant differences were found in antibody distribution between Bolton and elsewhere in NW England. The distribution of type-specific antibody was found to differ from that reported from the United States; this may be related to the differing distribution of serotypes causing disease in the US compared with England and Wales (Abbott and Graves, 1972).

References

- Abbott, J. D., and Graves, J. F. R. (1972). Serotype and sulphonamide sensitivity of meningococci isolated from 1966 to 1971. *J. clin. Path.*, 25, 528–530.
- Artenstein, M. S., Brandt, B. L., Tramont, E. C., Branche, W. C., Fleet, H. D., and Cohen, R. L. (1971). Serologic studies of meningococcal infection and polysaccharide vaccination. *J. infect. Dis.*, 124, 277–288.

Aspects of the Epidemiology of Haemophilus Meningitis D. C. TURK (*Bacteriology Department, Radcliffe Infirmary, Oxford*) The incidence of this disease has increased greatly in the past 30–40 years, notably in N America but also in Britain and many other countries. It is now, at least in those countries, the commonest form of meningitis in childhood. Mortality, almost 100% without treatment, can be reduced to 5% or so with suitable antibiotics, but residual disability is common. Virtually all cases are caused by capsulated type B strains of *Haemophilus influenzae*—a type carried in the nasopharynx of some 1–3% of the general population. The disease occurs as sporadic cases, with some seasonal concentration but no true epidemics and seldom with demonstrable case-to-case spread. A personal study of the families of 17 affected children confirmed the high frequency of carriers of type B strains in the environment of the children, and provided serological evidence that, in most cases at least, the strain had been in the family for some weeks before the child (usually the youngest) became ill. Little information about precipitating factors is available; evidence about the influence of social and economic factors is conflicting. The age incidence was explained in 1933 by demonstration that bactericidal antibodies for *H. influenzae* type B are commonly to be found in blood from newborn babies or older children or adults, but rarely in blood from children in the usual age range for the disease. Whether this pattern still applies is a matter currently under dispute—