Selective medium for the isolation of *Bordetella pertussis* and *parapertussis*

ENID M. SUTCLIFFE AND J. D. ABBOTT *From the Public Health Laboratory, Withington Hospital, Manchester*

Benzyl penicillin is widely used as a selective agent in media for the isolation of *Bordetella pertussis* and *parapertussis*. We report the use of cephalixin (Muggleton, O'Callaghan, Foord, Kirkby, and Ryan, 1968) as a selective agent with a wider spectrum than penicillin.

Preliminary experiments were carried out with a freeze-dried strain of *Bord. pertussis* originally isolated by Dr E. C. Armstrong on an antibiotic-free medium, and a number of freeze-dried strains of *Bord. parapertussis*. Surface viable counts were made on Oxoid charcoal agar (CM119) with 10% horse blood and various concentrations of cephalixin (93% potency). The strain of *Bord. pertussis* was not inhibited by 10 to 80 µg/ml of cephalixin, but there was some inhibition at 100 µg/ml. The strains of *Bord. parapertussis* were more sensitive to cephalixin: there was little or no inhibition by 40 µg/ml, slight inhibition by 60 µg/ml, and greater inhibition by levels of 80 to 100 µg/ml. At 60 µg/ml of cephalixin some strains of *Bord. parapertussis* showed variation of colony size after five days' incubation but this had disappeared after seven days. From these preliminary experiments we chose a level of 40 µg/ml of cephalixin for use as a selective agent in the charcoal-blood-agar base. During these experiments we also found that some strains of *Bord. parapertussis* showed marked variation of colony size on certain batches of charcoal-blood-agar containing no antibiotic; other strains were unaffected. This inhibitory effect of some batches of charcoal-blood-agar was removed by the addition of a final concentration of 1% Difco proteose peptone no. 3.

Comparison of Isolation of *Bord. pertussis* on Two Media

MEDIUM A (ROUTINE)

Charcoal agar plates with a final concentration of 10% horse blood, 1% Difco proteose peptone no. 3, and 0.25 units benzyl penicillin per ml.

MEDIUM B (EXPERIMENTAL)

As above with the substitution of 40 µg/ml cephalixin for the benzyl penicillin.

Plates were incubated for seven days at 36°C before final reading. Fresh batches were poured each week and stored at 4°C before use. Twenty-three routine pernasal swabs were inoculated directly on medium A, then placed in stabs of charcoal agar without blood or antibiotic and later the same day inoculated onto medium B. For the next 23 swabs the procedure was reversed and a further 29 swabs were inoculated onto both media from the charcoal agar stabs. The results of isolation of *Bord. pertussis* are shown in Table I and the growth of commensals in Table II.

Medium	Number of Isolations of <i>B. pertussis</i>
A only	2
B only	3
A & B	18
Total number of swabs	75

Table I Isolation of *Bord. pertussis* on routine (A) and experimental (B) media

Medium	No. of Swabs According to Growth of Commensals				Total No. of Swabs
	< 20 Colonies	20-50 Colonies	> 50 Colonies	Overgrown by Moulds	
A	38	5	31	1	75
B	64	2	9	0	75

Table II Growth of commensals on routine (A) and experimental (B) media

Although the isolation rate on each medium was similar, the experimental medium gave a significant reduction in growth of commensals, which facilitated the examination of the plates. The inhibition included penicillin-resistant staphylococci, some coliforms, and many, but not all, strains of *Haemophilus influenzae*. In a previous series of tests the results of direct inoculation of the routine medium was compared with a lower concentration of cephalixin (30 µg/ml) in charcoal-blood-agar inoculated from charcoal agar stabs. Of 146 swabs tested, *Bord. pertussis* was grown from 25 swabs on both media, from two swabs on the routine penicillin medium only, and from six swabs on the cephalixin medium only.

If serotyping is to be carried out it may be necessary to subculture strains of *Bord. pertussis* isolated on the cephalixin medium onto penicillin charcoal-blood-agar or antibiotic-free charcoal-blood-agar,

as there may be poor development of type-specific antigen on the cephalixin medium (N. W. Preston, personal communication).

Conclusions

A medium containing Oxoid charcoal agar (CM119) with 10% horse blood, 1% Difco proteose peptone no. 3, and 40 µg/ml of cephalixin has proved satisfactory for the isolation of *Bord. pertussis*. The isolation rate was as good on this medium as on the routine medium containing penicillin, but plates were easier to read due to the reduction in numbers of commensals. The cephalixin plates should be incubated for seven days at 36°C before being

reported as negative, since some strains of *Bord. parapertussis* may show variation in colony size with shorter periods of incubation.

We thank Mr R. W. Boyd of Glaxo Laboratories for the supply of cephalixin; Dr Hans Lautrop, Dr N. W. Preston, and Dr F. W. Sheffield for freeze-dried strains of *Bord. parapertussis*, and Dr E. C. Armstrong for the strain of *Bord. pertussis*.

Reference

Muggleton, P. W., O'Callaghan, Cynthia H., Foord, R. D., Kirkby, Susan M., and Ryan, D. M. (1968). In *Laboratory Appraisal of Cephalixin. Antimicrobial Agents and Chemotherapy*, edited by Gladys L. Hobby, pp. 353-360.

The July 1972 Issue

THE JULY 1972 ISSUE CONTAINS THE FOLLOWING PAPERS

Obituary A. G. Signy, Editor, *Journal of Clinical Pathology*, 1944-1972

Congenital dyserythropoietic anaemia with erythroblastic multinuclearity MALCOLM BRIGHT, JEREMY COBB, BYRON EVANS, AND T. E. PARRY

The influence of spleen size on the distribution of red cells and plasma P. J. TOGHILL AND SHEILA GREEN

Effect of feeding fat on fibrinolysis, stypven time, and platelet aggregation in elderly Africans J. C. FERGUSON, N. MACKAY, AND G. P. MCNICOL

Exfoliative cytology of diffuse mesothelioma G. HEFIN ROBERTS AND G. M. CAMPBELL

Immunoelectrosmophoresis in the diagnosis of meningococcal infections BARBARA M. TOBIN AND D. M. JONES

In-use testing of disinfectants in hospitals JEAN PRINCE AND G. A. J. AYLIFFE

Serum α-macroglobulins in renal disease and preeclampsia C. H. W. HORNE, J. D. BRIGGS, P. W. HOWIE, AND A. C. KENNEDY

Data processing in haematology M. G. NELSON AND E. ELDER

Red cell size as measured by the Coulter model S TAKASHI OKUNO

Some observations and improvements on SMA 6/60 determinations of creatinine MADELEINE CLOSE-MOLL AND J. G. LINES

Present day practice
Immunoglobulin and intrinsic factor antibody in the sera of patients with pernicious anaemia HING-YAN SHUM, ARTHUR M. STREETER, AND BARRY J. O'NEILL

Technical methods
Simultaneous staining of phospholipids, basic proteins, and glycogen on the same slide B. PRESENTLEY AND K. PERK

Book reviews

Supplement: Problems Related to Fibrinolysis—Theoretical and Clinical

Copies are still available and may be obtained from the PUBLISHING MANAGER, BRITISH MEDICAL ASSOCIATION, TAVISTOCK SQUARE, LONDON, WC1H 9JR, price £1.05.