- Cooper, H. L., and Rubin, A. D. (1965). RNA metabolism in lymphocytes stimulated by phytohemagglutinin: initial responses to phytohemagglutinin. *Blood*, 25, 1014-1027.
- Gardikas, C., and Israels, M. C. G. (1948). The Feulgen reaction applied to clinical haematology. *J. clin. Path.*, 1, 226-228.
- Gillis, E. M., and Baikie, A. E. (1964). Method for the demonstration of nucleoli in lymphocytes and other blood and bone marrow cells. *J. clin. Path.*, 17, 573-574.
- Greig, H. B. W. (1959). A substitute for the Feulgen staining technique. *J. clin. Path.*, 12, 93-94.
- Huhn, D., and Stich, W. (1969). *Fine Structure of Blood and Bone Marrow*, pp. 100-107. Lehmann, Munich. Hafner, New York.

- Potměšil, M., Smetana, K., and Wienerová, E. (1965). The nucleolar coefficient of the lymphocytes in the peripheral blood of patients with Hodgkin's disease. *Neoplasma*, 12, 205-214.
- Smetana, K., Janele, J., and Malinský, L. (1966). Nucleolar coefficient of lymphocytes in the peripheral blood of female patients with breast carcinoma. *Neoplasma*, 13, 643-648.
- Smetana, K., Lejnar, J., and Potměšil, M. (1969). A further contribution to the demonstration of RNA and nucleoli of blood cells in smear preparations. *Folia haemat. (Lpz.)*, 91, 381-394.
- Studzinski, G. P., and Gierthy, J. F. (1972). Cytologic appearance of the nucleolus of normal and neoplastic cells in relation to the synthesis of RNA. *Acta cytol. (Baltimore)*, 16, 245-248.

Letter to the Editor

Candida Isolates

Regarding Dr R. C. Bartlett's letter commenting on our paper (Stieritz, Law and Holder, *J. clin. Path.*, 1973, 26, 405-408), we did not mean to imply all *Candida* isolates from any patient should be speciated. Our paper concerned burned patients infected with *Candida* and in the context of *Candida* isolated from the wounds, urine, and blood of compromised hosts we still feel that speciation is important.

While I concur with Dr Bartlett's comment that 'the decision to treat should be based on clinical evidence of infection, not the species isolated', we have found this not to be the case when dealing with *Candida* infections. Because of the highly toxic nature of Amphotericin B, the decision to treat *Candida* infections is frequently delayed until the clinician is forced to treat or lose the patient. As we pointed out, many clinicians consider *C. albicans* the only *Candida* which is of clinical concern. When they receive a culture report stating that *Candida* other than *albicans* has been isolated from an infected patient, they may be lulled into a false sense of security and then finally be forced to treat with too little too late.

Our data showed that while all of the Amphotericin B MICs for *C. albicans* were 0.16 µg/ml or below, most of the MICs for *C. tropicalis* were at least one dilution higher, and in some cases two and three dilutions higher. This may imply that *C. tropicalis* infection could be a treatment problem, whereas *C. albicans* infection may not be. Support for this idea may be found in the fact that the first report, to my knowledge, of clinical isolates of *Candida* being resistant to Amphotericin B has recently been

published (Woods, R. A., *et al*, 1974). These resistant isolates were *C. tropicalis* not *C. albicans*.

While we all would like to keep the costs of health care to a minimum, the attitude that we should only do that clinical microbiology which is 'essential to diagnosis and treatment', I believe to be incorrect. In the evolution of infectious diseases and in defining infection problems, the microbiology that was 'essential to diagnosis and treatment' yesterday is not the same as the microbiology that is 'essential to diagnosis and treatment' today; nor will today's microbiology suffice tomorrow. We, the clinical microbiologists, must be alert to identifying new infection problems; the latest 'emerging' pathogens, and it must be our responsibility to warn the medical fraternity of these new dangers. If we don't, who then?

IAN ALAN HOLDER

Department of Microbiology,
Shriners Burns Institute,
Cincinnati, Ohio 45219

Reference

- Woods, R. A., *et al* (1974). Resistance to polyene antibiotics and correlated sterol changes in two isolates of *Candida tropicalis* from a patient with an amphotericin B-resistant funguria. *J. infect. Dis.*, 129, 53.

Book reviews

Transplacental Carcinogenesis Proceedings of a meeting held at the Medizinische Hochschule, Hanover, 1971 (IARC Scientific Publications No. 4). Edited by L. Tomatis and U. Mohr. (Pp. xiii + 181; illustrated. £5.00; \$12.00; Sw. fr. 40.-) Geneva: World Health Organization. International Agency for Research on Cancer. 1973. Available from HMSO, London.

Interest in the transplacental action of certain carcinogens was widely stimulated by the report from Boston in 1971 of a cluster of eight cases of adenocarcinoma of the vagina in young women; the mothers of all but one of them had received stilboestrol during their pregnancy. Although the transplacental carcinogenic activity of stilboestrol has not yet been reproduced in experimental animals, the general subject of transplacental carcinogenesis has been extensively investigated. A useful summary of current knowledge is presented in this book. Particular attention has been paid to the nitrosamines, and one group of them—the nitrosoalkylureas—has been clearly shown to act as transplacental carcinogens with the brain as the main target organ. The dose of the compound given and the stage of foetal development both appear to be crucial in determining the outcome—in particular, death or induction of malformations rather than tumours. Predictably, no tidy relationship appears to exist between teratogenic and carcinogenic activities, and many details of the actual mechanisms of transplacental carcinogenesis are still quite unknown.

The subject is, however, one of considerable potential importance. Tumours are now a major cause of death among children in advanced societies, and the