A note on the use of immunofluorescent methods for the detection of *Pseudomonas aeruginosa* in bronchitic sputum

R. L. SANDS AND I. D. GREEN

From the Department of Bacteriology and Department of Medicine, University of Birmingham, Birmingham B15 2TJ

SYNOPSIS By immunofluorescent microscopy of sputum from 67 cases of bronchitis *Pseudomonas aeruginosa* was detectable in 14 as compared with nine by cultural methods.

Recent studies have shown that immunological methods may be of value for the detection of infective agents in respiratory disease (Nicholls *et al*, 1975). Immunofluorescence has been employed for the diagnosis of *Haemophilus* in sputum (Sell *et al*, 1963).

Severe respiratory tract infections may be caused by *Ps. aeruginosa* during hospitalization and this organism may prove invasive. Thus its identification, in the early infective stages, is desirable (Perry and Sellors, 1963; Fraser and Paré, 1970).

Material and Methods

HOMOLOGOUS ANTISERUM

Ps. aeruginosa was obtained from a case of pneumonia in the Queen Elizabeth Hospital, Birmingham and found to be a non-typable polyagglutinating strain by the Cross-infection Laboratory, Colindale. Antiserum was raised in rabbits by the method of Yager *et al* (1960) using the inoculation schedule of Thomason and Cherry (1963). The agglutination titre of the antiserum against the homologous strain of *Ps. aeruginosa* was found by the Widal technique to be 1/250, and it was subsequently shown by immunofluorescence to react with this strain and 14 newly isolated strains but failed to react with one. No cross-reactions were observed by either method with *Haemophilus influenzae*, *Staphylococcus epidermidis*, *Proteus mirabilis*, *Klebsiella pneumoniae*, *Klebsiella aerogenes*, group A streptococcus, α -haemolytic streptococcus or *Neisseria catarrhalis*.

ANTISPECIES IMMUNOFLUORESCENT CONJUGATE

Commercial reagents having proved unsatisfactory, Received for publication 15 July 1975

immunofluorescent conjugate was prepared; sheep anti-rabbit immunoglobulin G was obtained from the Department of Experimental Pathology, University of Birmingham, and conjugated to fluorescein isothiocyanate (FITC) isomer 1 using the method of Nairn (1969). Unconjugated fluorochrome and impurities present in the FITC were removed by passing *c* 80 mg of protein through a 2.5 × 42 cm chromatography column containing Sigma G-25 Sephadex in a gel sample ratio of 6:1 (v/v).

The purified conjugate was checked for the presence of unconjugated dye using polyacrylamide gel electrophoresis (PAGE) (Ornstein, 1964; Davis, 1964) and running the gels at 493 nm in a microdensitometer (Cartwright and Jaynes, 1972). The preparation was absorbed with ox liver homogenate supplied by the Department of Rheumatism Research, University of Birmingham by the technique of Nairn (1969).

The prepared conjugate was examined by PAGE, double gel diffusion (Ouchterlony, 1948), immunoelectrophoresis (Scheidegger, 1955), chessboard titration, incorporating normal rabbit serum against conjugate dilutions as a test for non-specific staining (Hale and Bergquist, 1971), and the fluores-

PAGE	No free dye detected
Double gel diffusion	Single arc of identity
Immunoelectrophoresis	Single arc of the IgG class
Chessboard titration	PEP 1/64
	PT 1/200 to 1/500
Molar fluorescein to protein ratio	2:3

PEP = plateau end-point

PT = plateau titre (Hale and Bergquist, 1971)

Table I *Assessment of a fluorescent antibody conjugate*

cein to protein ratio (F:P) was calculated (Jobbágy and Király, 1966) (table I).

The conjugate was also tested for specificity against the previously listed bacteria, using the homologous antibacterial antiserum, prepared in this laboratory, for the indirect immunofluorescence test. Staining reactions were controlled for inhibition of fluorescent antibodies by pretreatment with unconjugated sheep anti-rabbit immunoglobulin G, and substitution of normal rabbit serum for the homologous antiserum.

INDIRECT STAINING TECHNIQUE

Heat-fixed bacterial smears were covered with homologous antiserum and incubated at room temperature for 1 hour in Petri dishes containing moist filter paper. Slides were washed in phosphate buffered saline (PBS) pH 7.2, 0.145M for 30 minutes and after the removal of excess moisture were incubated with homologous antispecies conjugate for 30 minutes. After the slides had been washed in PBS for 30 minutes the preparations were mounted in glycerol/PBS (9:1) and the coverslips were sealed with wax. Preparations were examined by dark-ground, blue fluorescence with the $\times 55$ oil immersion lens of a Gillett and Siebert 'Research Conference' microscope.

BACTERIOLOGICAL METHODS

Sputum samples were homogenized after pancreatin digestion and, for the purposes of comparison, were examined by Gram's stain and inoculated on horse blood agar and McConkey agar, incubated at 37°C, and inspected after 24 and 48 hours. Colonies were picked, Gram-stained, and subcultured for further examination, in accordance with routine procedures.

Results

A specimen of sputum was taken from each of 67 subjects of various ages and both sexes, clinically diagnosed as bronchitic. Specimens were examined

by both cultural and immunofluorescence methods as soon as possible. The results are given in table II and show that more positive samples were obtained by immunofluorescence examination, while substantiating, in all but one case, the results of cultural methods. Control reactions were all negative.

Discussion

The results recorded show that immunofluorescence provides a rapid and sensitive method of detecting a potentially important pathogen in bronchitic sputum, where rapid diagnosis provides the possibility of early chemotherapy that might otherwise be delayed.

Those cases in which a positive immunofluorescent diagnosis of *Ps. aeruginosa* was made, in the absence of a positive culture, are explicable either by cultural failure or as false positives. The former might be caused by inadequate methods, transfer of antibiotics with the inoculum, low or non-viability of the bacteria. Improved selective cultural methods should eliminate the discrepancy, but the work, as presented, is intended to provide a comparison between this method and the cultural techniques currently applied in a large number of hospital laboratories.

We wish to thank Dr. K. A. Bisset for his help and advice.

References

- Cartwright, N. J. and Jaynes, M. H. (1972). A densitometer attachment for gel electrophoretic measurements in the Unicam SP800 spectrophotometer. *Lab. Pract.*, **21**, 643-644 and 648.
- Davis, B. J. (1964). Disc electrophoresis. II. Method and application to human serum proteins. *Ann. N.Y. Acad. Sci.*, **121**, 404-427.
- Fraser, R. G. and Paré, J. P. A. (1970). *Diagnosis of Diseases of the Chest*, vol. 1, p. 600. W. B. Saunders, Philadelphia, London and Toronto.
- Hale, W. L. and Bergquist, R. (1971). Chessboard analyses with antinuclear antibodies. *Ann. N.Y. Acad. Sci.*, **177**, 354-360.
- Jobbágy, A. and Király, K. (1966). Chemical characterisation of fluorescein isothiocyanate-protein conjugates. *Biochem. Biophys. Acta*, **124**, 166-175.
- Nairn, R. C. (1969). *Fluorescent Protein Tracing*, 3rd ed. Livingstone, Edinburgh and London.
- Nicholls, A. C., Pease, P. E., and Green, I. D. (1975). Agglutinin response to bacterial infection in acute exacerbations of chronic bronchitis. *J. clin. Path.*, **28**, 279-283.
- Ornstein, L. (1964). Disc-electrophoresis. I. Background and theory. *Ann. N.Y. Acad. Sci.*, **121**, 321-349.
- Ouchterlony, Ö. (1948). *In vitro* method for testing the toxin-producing capacity of diphtheria bacteria. *Acta path. microbiol. scand.*, **25**, 186-191.
- Perry, K. M. A. and Sellors, Sir T. H. (1963). *Chest Diseases*, vol. 2. Butterworth, London.

	<i>Pseudomonas aeruginosa</i>
Immunofluorescence located	14
Gram stain and culture	9
Location by both immunofluorescence, Gram stain, and culture	8
Negative for immunofluorescence but positive for Gram stain and culture	1
Positive for immunofluorescence but negative for Gram stain and culture	6

Table II A comparison of immunofluorescence and routine bacteriology

Total number of patients investigated—67

- Scheidegger, J. J. (1955). Une micro-méthode de l'immuno-électrophorèse. *Int. Arch. Allergy*, 7, 103-110.
- Sell, S. H. W., Saunders, R. S., and Cheatham, W. J. (1963). *Hemophilus influenzae* in respiratory infections. II. Specific serological antibodies identified by agglutination and immunofluorescent techniques. *Amer. J. Dis. Child.*, 105, 470-474.
- Thomason, B. M. and Cherry, W. B. (1963). Immuno-

- fluorescence techniques in the diagnosis of infections due to enteropathogenic *Escherichia coli*. *Rev. lat.-amer. Microbiol. Parasitol.*, 6, 63-76.
- Yager, R. H., Spertzel, R. O., Jaeger, R. F., and Tigertt, W. D. (1960). Domestic fowl—source of high titer *P. tularensis* serum for the fluorescent antibody technic. *Proc. Soc. exp. Biol. (N.Y.)*, 105, 651-654.

Reports and Bulletins prepared by the Association of Clinical Biochemists

The following reports and bulletins are published by the Association of Clinical Biochemists. They may be obtained from The Publishing Department, British Medical Journal (ACB Technical Bulletins), B.M.A. House, Tavistock Square, London WC1H 9JR. Overseas readers should remit by British Postal or Money Order.

SCIENTIFIC REPORTS (price £1.00/\$2.00 each)

- 3 Automatic Dispensing Pipettes: an assessment of 35 commercial instruments September 1967 P. M. G. BROUGHTON, A. H. GOWENLOCK, G. M. WIDDOWSON, and K. A. AHLQUIST
- 4 An Evaluation of five Commercial Flame Photometers suitable for the Simultaneous Determination of Sodium and Potassium March 1970 P. M. G. BROUGHTON and J. B. DAWSON

SCIENTIFIC REVIEWS (price £1.00/\$2.00 each)

- 1 The Assessment of Thyroid Function March 1971 F. V. FLYNN and J. R. HOBBS
- 2 Renal Function Tests Suitable for Clinical Practice January 1972 F. L. MITCHELL, N. VEALL, and R. W. E. WATTS
- 3 Biochemical Tests for the Assessment of Feto-placental Function May 1975 C. E. WILDE and R. E. OAKEY

TECHNICAL BULLETINS (price £1.00/\$2.00 each)

- 9 Determination of Urea by AutoAnalyzer November 1966 RUTH M. HASLAM
- 11 Determination of Serum Albumin by AutoAnalyzer using Bromocresol Green October 1967 B. E. NORTHAM and G. M. WIDDOWSON
- 13 An Assessment of the Technicon Type II Sampler Unit March 1968 B. C. GRAY and G. K. MCGOWAN
- 14 Atomic Absorption Spectroscopy: an outline of its principles and a guide to the selection of instruments May 1968 J. B. DAWSON and P. M. G. BROUGHTON
- 15 A Guide to Automatic Pipettes (2nd edition) June 1968 P. M. G. BROUGHTON
- 16 A Guide to Automation in Clinical Chemistry May 1969 P. M. G. BROUGHTON
- 17 Flame Photometers: a comparative list of 17 instruments readily available in Britain August 1969 P. WILDING
- 19 Spectrophotometers: a comparative list of low-priced instruments readily available in Britain May 1970 C. E. WILDE and P. SEWELL

- 20 Quantities and Units in Clinical Biochemistry June 1970 P. M. G. BROUGHTON

- 21 Filter Fluorimeters: A comparative list of 18 instruments September 1970 H. BRAUNSBURG and S. S. BROWN

- 22 Bilirubin Standards and the Determination of Bilirubin by Manual and Technicon AutoAnalyzer Methods January 1971 BARBARA BILLING, RUTH HASLAM, and N. WALD

- 23 Interchangeable Cells for Spectrophotometers and Fluorimeters September 1971 S. S. BROWN and A. H. GOWENLOCK

- 24 Simple Tests to Detect Poisons March 1972 B. W. MEADE *et al.*

- 25 Blood Gas Analysers May 1972 K. DIXON

- 26 Kits for Enzyme Activity Determination September 1972 S. B. ROSALKI and D. TARLOW

- 27 Assessment of Pumps Suitable for Incorporation into Existing Continuous Flow Analytical Systems November 1972 A. FLECK *et al.*

- 28 Routine Clinical Measurements of Transferrin in Human Serum September 1973 K. DIXON

- 29 Control Materials for Clinical Biochemistry (5th edition) September 1973 J. F. STEVENS

- 30 Notes on the Quality of Performance of Serum Cholesterol Assays September 1973 S. S. BROWN

- 31 Determination of Uric Acid in Blood and in Urine July 1974 R. W. E. WATTS

- 32 A Survey of Amino Acid Analysers Readily Available in the United Kingdom September 1974 J. E. CARLYLE and P. PURKISS

- 33 Definitions of some Words and Terms used in Automated Analysis November 1974 A. FLECK, R. ROBINSON, S. S. BROWN, and J. R. HOBBS

- 34 Measurement of Albumin in the Sera of Patients January 1975 LINDA SLATER, P. M. CARTER, and J. R. HOBBS

- 35 Investigation of the Validity of Temperature Correction Factors for Serum Aspartate and Alanine Transaminases March 1975 S. B. ROSALKI *et al.*