- Kaplow, L. S. and Beck, R. E. (1964). Leukocyte alkaline phosphatase response to surgical stress. *Surg. Gynec. Obstet.*, **119**, 97-102.
- Rutenburg, A. M., Rosales, C. L., and Bennett, J. M. (1965). An improved histochemical method for the demonstration of leukocyte alkaline phosphatase activity: clinical application. *J. Lab. clin. Med.*, **65**, 698-705.
- Trubowitz, S., Feldman, D., Benante, C., and Hunt, V. M. (1959). The alkaline phosphatase content of the human polymorphonuclear leukocyte in blood and marrow. *Amer. J. clin. Path.*, **31**, 483-486.
- Valentine, W. N., Follette, J. H., Hardin, E. B., Beck, W. S., and Lawrence, J. S. (1954). Studies on leukocyte alkaline phosphatase activity: relation to "stress" and pituitary-adrenal activity. *J. Lab. clin. Med.*, **44**, 219-228.
- Williams, D. M. (1975). Leucocyte alkaline phosphatase as a marker of cell maturity: a quantitative cytochemical and autoradiographic study. *Brit. J. Haemat.*, **31**, 371-379.
- Wyllie R. G. (1962). Neutrophil alkaline phosphatase. Response to acute inflammation. *Aust. Ann. Med.*, **11**, 118-125

A. S. D. SPIERS and
D. M. WILLIAMS

MRC Leukaemia Unit and
Department of Histopathology,
Royal Postgraduate Medical School,
Du Cane Road, London W12 0HS

Platelet Counts in Normal Pregnancy

We are pleased to be able to confirm the findings of Sejeny *et al* (*J. clin. Path.*, **28**, 812-813, 1975) that, using a Coulter Counter, the platelet count decreases progressively and significantly in normal pregnancy. Sejeny *et al* suggest, without adducing any evidence, that this decrease in count may be due to an increase in plasma volume. We, however, have evidence that does not prove, but strongly suggests, that this haemodilutional effect

is not nearly sufficient to account for the observed drop in platelet count.

We have studied about 30 patients in each trimester and 24 non-pregnant women. The results will be published elsewhere, but in brief we have found a significant progressive decrease in count and increase in the mean platelet volume so that the total platelet volume mass per ml (volume $\times$ numbers/ml) remained approximately constant. The concept of a constant total platelet mass has been reported before (O'Brien and Jamieson, 1974; Behrens, 1975). Additionally, we found marked shortening of the heparin thrombin clotting time of platelet-poor plasma. This may perhaps reflect the presence of platelet factor 4 liberated into the plasma as the result of thrombosis that occurs normally and extensively even in the healthy placenta.

From the table it will be seen that, as expected, the haemoglobin falls progressively through pregnancy. The packed cell volume also fell but there was no evidence of iron deficiency developing since the MCHC remained constant. Therefore it seems reasonable to assume that the fall in haemoglobin reflected a relative increase in plasma volume. Assuming that the original total platelet count remained constant, in this case the mean was $272 \times 10^9/l$, it is then possible to calculate what the count would have been if it had been diluted in the increased plasma volume as had the red cells. It will be seen that the observed count decreased far in excess of the calculated figure due to haemodilution. Thus very probably the total number of circulating platelets per unit volume decreases absolutely.

References

- Behrens, W. E. von (1975). Mediterranean macrothrombocytopenia. *Blood*, **46**, 199-207.
- O'Brien, J. R. and Jamieson, S. (1974). A relationship between platelet volume

and platelet number. *Thrombos. Diath. haemorrh. (Strutt.)*, **31**, 363-365.

Sejeny, S. A., Eastham, R. D., and Baker, S. R. (1975). Platelet counts during normal pregnancy. *J. clin. Path.*, **28**, 812-813.

J. R. O'BRIEN

Central Laboratory

St. Mary's General Hospital, Portsmouth

Amylase Assay by the Phadebas Method

In the past we have personally corrected a misconception among some colleagues concerning the assay of amylase by the Phadebas method but now that the erroneous notion has recently appeared in print it becomes necessary to respond likewise. Ojala and Harmoinen (1975), in the discussion on the methodology of the Phadebas amylase kit (Pharmacia AB), confirm the manufacturer's literature that the substrate does not remain suspended but sinks after the initial shaking at the beginning of incubation. They then go on to claim that 'a clearly higher and better reproduced amylase value was obtained by shaking the mixture by hand vigorously during the entire incubation'. We find no significant difference in activity or reproducibility of the assay whether the reaction tubes are (A) shaken or (B) remain unshaken during incubation. A mean amylase value = 501 ± 12 U/l, $n = 10$
B mean amylase value = 500 ± 12 U/l, $n = 10$.

Higher absorbances were encountered with the shaken tubes (A), as were noted originally by Ceska *et al* (1969), but this was paralleled by a higher absorbance of the blank.

The implication of the statement Ojala and Harmoinen is that while the substrate sediments the enzyme remains in solution in the supernatant. However, it can readily be shown by the experiments detailed below that the amylase is adsorbed on to the solid phase substrate during the initial mixing after addition of the tablet. A further series of reaction tubes containing the same serum was centrifuged immediately after addition of the tablet to the assay mixture. The supernatants were decanted into clean tubes, absorbances were noted, and incubation was carried out after the addition of a further tablet. The increase in absorbance in this series was minimal ($\Delta A < 0.01$). Simultaneously with this series, 4 ml of 0.9% saline was added to the residues in the original tubes and the assay procedure was continued. The

	Controls	1st Trimester	2nd Trimester	3rd Trimester
Haemoglobin (g/dl)	13.4	12.7	12.5	12.2
Packed cell volume	0.41	0.39	0.38	0.37
Haemoglobin per unit volume (MCHC)	32.7	32.6	32.9	33.0
Observed platelet count ($\times 10^9/l$)	272	249	243	210
Expected count due to haemodilution	—	257	253	247

Table Possible effect of haemodilution on platelet count: means