Monoclonal gammopathy (Waldenström's macroglobulinaemia) producing specific red cell antibody

O. A. OLUBOYEDE², O. BADEMOSI¹, A. DAVID-WEST², C. O. THOMAS²,
T. I. FRANCIS, AND L. LUZZATTO²

From the Departments of Medicine¹ and Haematology², University of Ibadan, Nigeria and University College Hospital, Ibadan, Nigeria

SYNOPSIS Two cases of Waldenström's macroglobulinaemia have been seen at University College Hospital, Ibadan in the last four years. Case 1 was a 30-year-old soldier who presented with splenomegaly and anaemia, was treated with chlorambucil, and had a complete remission sustained for over two years. Case 2 was a 58-year-old retired civil servant who presented with very severe anaemia and also splenomegaly, and died within three weeks of admission. Both patients had most of the typical features of Waldenström's disease, including retinal changes and serum IgM levels of 4200 and 5500 mg/dl respectively. In both cases an atypical cold antibody was detected in the course of blood cross-matching procedures. In case 1, the antibody agglutinated all adult and cord red cells tested, including the patient's own cells, to a titre of 8000 and above at 4°C. Surprisingly enough, when the patient went into remission and the serum IgM level had fallen to 400 mg/dl, this antibody was no longer detectable and has not reappeared two years later. In case 2, the antibody agglutinated all adult red cells tested to a titre of 2000 at 20°C but not the patient's own red cells. Since cord cells were agglutinated only to a titre of 4 to 20°C it was concluded that the patient had an alloantibody with I-specificity. Therefore in both these patients the monoclonal immunoglobulin produced by the neoplastic lymphoid cell clone had specific activity against red cell antigens.

Since Waldenström's original description of three cases of macroglobulinaemia in 1944, various facets of these groups of monoclonal gammopathies have been described and the literature has been extensively reviewed (Imhof, Baars, and Verloop, 1959; Waldenström, 1961; Cohen *et al*, 1964; McCallister *et al*, 1967; Waldenström, 1970). The scarcity of reports of primary Waldenström's macroglobulinaemia in Africa would tend to suggest that this condition does not occur. McFarlane and Nwokolo (1966) described a Nigerian woman with Waldenström-type macroglobulinaemia associated with rheumatoid arthritis. This report presents two cases of primary Waldenström's macroglobulinaemia seen in the University College Hospital, Ibadan over the last 10 years.

Case Reports

CASE 1 (table III)

An 18-year-old soldier was seen in the medical out-
Received for publication 11 August 1975

patient department of the University College Hospital (UCH), Ibadan on 28 November 1969. His illness had started three months earlier with progressive weakness, tiredness, and backache. He had been found to be anaemic and had been treated initially with buphenium hydroxy naphthoate (Alcopar) and iron in another hospital. Since he showed no response to medication, he was transfused with 4 units of packed cells over five days. He was then referred to UCH, Ibadan for further evaluation and treatment.

On admission, abnormal physical signs were pallor, mild icterus, a soft, slightly tender liver palpable 6 cm below the right costal margin, and a non-tender spleen palpable 2 cm below the left costal margin. There was no significant lymphadenopathy.

He developed photophobia, and funduscopy showed distention of retinal veins and bilateral fundal haemorrhages.

Investigations performed included: packed cell volume (PCV) 21%, total white cell count 3900/mm³; platelet count 305 000/mm³; reticulocyte count

his eight weeks in the hospital, the patient complained of blurring of vision, and petechiae were noticed on the face, lips, buccal mucosa, and trunk. Plasmapheresis was performed: the PCV after the procedure was 28%. Hepatosplenomegaly gradually decreased and the patient was discharged after three months in hospital. At the time of discharge, the fundal haemorrhages had resolved and vision was normal. The PCV was 37%. The patient returned to his duties as a soldier. Two years after discharge, his PCV was 38% and the macroglobulin level was 4.5 mg per ml.

CASE 2 (table III)

A 58-year-old farmer was admitted on 20 August 1973 with an 18-month history of chest pain and progressive lethargy. Between 1969 and 1970, he had been treated for anaemia but did not respond to treatment.

Abnormal physical signs were gross pallor and a tender spleen palpable 2 cm below the left costal margin. On fundoscopy he had retinal haemorrhages, with tortuosity and varicosities of retinal capillaries producing the so-called fundus paraprroteinaemicus.

Investigations performed were as follows: PCV 12%, WBC 9600/mm³, platelets 128000/mm³. Sternal marrow biopsy showed erythroid hyperplasia with marked lymphocytosis. The total serum proteins were 7.7 g/dl, albumin 2.1 g/dl, and globulin 5.6 g/dl. Electrophoresis of the serum on cellulose acetate membrane showed an intense and homogeneous band in the gamma globulin region which proved to be macroglobulin. Quantitation of the IgM gave 61 mg/ml (method of Fahey and McKelvey). This IgM had only kappa light chains.

Further tests showed that the patient had a strong cold antibody with a specificity of anti-I (table II). However, the antibody did not agglutinate the patient's own red cells at room temperature and no autoagglutination was seen in the blood film. It was assumed that this antibody was an alloantibody and the patient's phenotype was adult-i but the necessary antiserum to prove this assumption was not available.

Despite repeated blood transfusions, plasmapheresis (thrice on three consecutive days) and chlorambucil, the patient died three weeks after admission. Permission for necropsy was refused.

Discussion

Waldenström's macroglobulinaemia is characterized by a marked increase in IgM and bone marrow abnormalities characterized by lymphocytosis (Dameshek, 1966). The other features like haemolysis, haemorrhages, and the hyperviscosity syndrome

are secondary manifestations of this monoclonal gammopathy. The clinical features of our patients satisfy all the criteria necessary for the diagnosis of Waldenström's macroglobulinaemia.

Macroglobulinaemia is characteristically a disease of old people, and the youngest age of onset of 107 patients reviewed by Imhof *et al* (1959) was 32 years.

It is not quite clear from the available literature whether an early age of onset is more often associated with a worse prognosis as regards failure of response to treatment and rapid termination. The mild hepatitis noted on admission in case 1 was probably the result of serum hepatitis following initial blood transfusion. Post-hepatitis remission was dramatic, and the patient has remained in remission for two years—up to the time of writing. It is tempting to implicate the hepatitis as a cause of remission. Similarly, reported cases of complete clinical and haematological remission following urethane treatment and severe hepatitis in multiple myeloma (London, 1955) and remission of macroglobulinaemia following serum hepatitis in a 59-year-old woman (Wolf *et al*, 1972) have been documented. Osserman and Takatsuki (1965) also described the case of a 75-year-old man with a transient IgG plasma cell dyscrasia with type K Bence Jones proteinuria who experienced a remission of the plasmacytic dyscrasia following cessation of sulphafurazole diethanolamine (Gantrisin) treatment and an episode of viral hepatitis. The mechanism by which viral hepatitis could cause a remission in primary macroglobulinaemia is unknown. One might suggest a direct effect where the viral antigen has destroyed the immunocompetent cells or that the viral antigen increased the resistance of the patients to the abnormal clone of plasma cells producing the abnormal immunoglobulins. Also, in our case 1, there was no evidence of an active haemolytic process which may have responded to treatment with steroids and chlorambucil. He showed no evidence to incriminate chronic active hepatitis (Sherlock, 1966) or subacute hepatitis (Cohen *et al*, 1963) as being the cause of his macroglobulinaemia.

In many chronic infections which may cause secondary macroglobulinaemia, like the tropical splenomegaly syndrome (Sagoe, 1970), trypanosomiasis (Mattern and Lobez 1964), kala azar, collagen or neoplastic disease, the IgM seldom forms a distinct and homogeneous band (figure). Also, the absolute values of IgM seldom exceed 40-50 mg/ml.

Treatment aims at reducing the plasma concentration of the circulating macroglobulins which are responsible for some of the clinical manifestations (MacKay *et al*, 1956; Bayrd *et al*, 1965). Removal of the abnormal proteins by plasmapheresis (Schwab and Fahey, 1960; Lawson *et al*, 1968) is a useful

Figure Case 1 The electrophoretic pattern of the serum both before and after treatment compared with the pattern of the serum from a normal Nigerian donor and the pattern of the serum from a patient with a polyclonal increase in IgM suffering from tropical splenomegaly syndrome (TSS).

emergency measure. Blood transfusions for the correction of anaemia, steroids, and cytotoxic drugs like chlorambucil or cyclophosphamide are useful in reducing the rate of synthesis of the macroglobulins. Prolonged continuous treatment with chlorambucil has been advocated by several workers (Bayrd *et al*, 1965; McCallister *et al*, 1967). Our first patient is very unusual in that a complete remission, lasting just over two years, has followed a short course of chlorambucil.

The two patients presented are the only cases of Waldenström's macroglobulinaemia on record in the University College Hospital, Ibadan. The first patient's serum contained an autoantibody with properties similar to those causing 'cold agglutinin disease (CAD)', although the clinical manifestations of Raynaud phenomena were not observed, probably because of the relatively high environmental temperature. CAD and Waldenström's disease are both so rare that the possibility of their coexisting in this patient by chance alone need not be entertained. Rather, the evidence presented clearly indicates that they are causally related. Indeed, Waldenström's macroglobulinaemia can be defined as a lymphoma in which the monoclonal neoplastic cells secrete an immunoglobulin of the IgM class. It may happen that the IgM produced has specificity against a particular red cell antigen.

If the antigen concerned is present on the patient's red cells, the syndrome of CAD will result. Hence we conclude that CAD is a special variant of Waldenström's disease, which is a special variant of lymphoma. We may venture to make a prediction, by

analogy, regarding another type of immunoglobulin-producing neoplasm, namely multiple myeloma (MM). It is possible that eventually a case of MM will be found, secreting an IgG immunoglobulin also having red cell antibody specificity. In such a case, if the corresponding antigen were present on the patient's red cells, a syndrome of *warm-type* autoimmune haemolytic anaemia would complicate the disease.

In the second patient, the antigen against which the antibody was directed appeared to be absent from the patient's own red cells. However, the presence of this antibody made it impossible to obtain compatible blood for the patient because the corresponding antigen (I) was present on the red cells of all adult donors tested.

References

- Bayrd, E. D., Hagedorn, A. B., and McGuckin, W. F. (1965). Macroglobulinaemia, its recognition and treatment. *J. Amer. med. Ass.*, **193**, 724-726.
- Cohen, R. J., Bohannon, R. S., and Wallerstein, R. O. (1964). Waldenström's macroglobulinaemia. A study of ten cases. *Amer. J. Med.*, **41**, 274-284.
- Cohen, S., Solomon, A., Paronetto, F., and Popper, H. (1963). Subacute hepatitis in Waldenström's macroglobulinaemia. *Amer. J. Med.*, **34**, 256-263.
- Dacie, J. V. (1965). Haemolytic anemia. *Ann. N. Y. Acad. Sci.*, **124**, 415-421.
- Dameshek, W. (1966). What is Waldenström's macroglobulinaemia? *Acta med. scand.*, Suppl., **445**, pp. 163-170.
- Fahey, J. L. and McKelvey, E. M. (1965). Quantitative determination of serum immunoglobulins in antibody-agar plates. *J. Immunol.*, **94**, 84-90.
- Imhof, J. W., Baars, H., and Verloop, M. C. (1959). Clinical and haematological aspects of macroglobulinaemia Waldenström. *Acta med. scand.*, **163**, 349-366.
- Lawson, N. S., Nosanchuk, J. S., Oberman, H. A., and Meyers, M. C. (1968). Therapeutic plasmapheresis in treatment of patients with Waldenström's macroglobulinaemia. *Transfusion, (Philad.)*, **8**, 174-178.
- London, R. E. (1955). Multiple myeloma: report of a case showing unusual remission lasting two years following severe hepatitis. *Ann. intern. Med.*, **43**, 191-201.
- McCallister, B. D., Bayrd, E. D., Harrison, E. G., Jr., and McGuckin, W. F. (1967). Primary macroglobulinemia: review with a report of thirty-one cases and notes on the value of continuous chlorambucil therapy. *Amer. J. Med.*, **43**, 394-434.
- McFarlane, H. and Nwokolo, C. (1966). Waldenström type macroglobulinaemia in a Nigerian with rheumatoid arthritis. *J. clin. Path.*, **19**, 603-605.
- MacKay, I. R., Eriksen, N., Motulsky, A. G., and Volwiler, W. (1956). Cryo- and macroglobulinemia; electrophoretic, ultracentrifugal and clinical studies. *Amer. J. Med.*, **20**, 546-587.
- Mancini, G., Carbonara, A. O., and Heremans, J. F. (1965). Immunochemical quantitation of antigens by single radial immunodiffusion. *Immunochemistry*, **2**, 235-254.
- Mattern, P. and Lobez, D. (1964). Techniques et intérêt épidémiologique du diagnostic de la trypanosomiase humaine Africaine par la recherche de la B₂ macroglobuline dans le sang et dans le L.C.-R. *Ann. Inst. Pasteur*, **107**, 3, 415-421.

- Osserman, E. F. and Takatsuki, K. (1965). Considerations regarding the pathogenesis of the plasmacytic dyscrasias. *Ser. haemat. (Scand. J. Haemat., Suppl.)*, **4**, 28-49.
- Sagoe, A. S. (1970). Tropical splenomegaly syndrome: long-term proguanil therapy correlated with spleen size, serum IgM and lymphocyte transformation. *Brit. med. J.*, **3**, 378-382.
- Schwab, P. J. and Fahey, J. L. (1960). Treatment of Waldenström's macroglobulinemia by plasmapheresis. *New. Engl. J. Med.*, **263**, 574-579.
- Sherlock, S. (1966). Waldenström's chronic active hepatitis. *Acta med. scand. Suppl.*, **445**, pp. 426-433.
- Waldenström, J. (1944). Incipient myelomatosis or 'essential' hyperglobulinaemia with fibrinogenopenia—a new syndrome. *Acta med. scand.*, **117**, 216-247.
- Waldenström, J. (1961). Studies on conditions associated with disturbed gamma globulin formation (gammopathies). *Harvey Lect.*, **56**, 211-231.
- Waldenström, J. (1970). Benigne und maligne monoklonale Hypergamma + globulinaemia (Gammopathie). *Schweiz. med. Mschr.*, **100**, 290-296.
- Wolf, R. E., Riedel, L. O., Levin, W. C., and Ritzmann, S. E. (1972). Remission of macroglobulinemia coincident with hepatitis. *Arch. intern. Med.*, **130**, 392-395.

The February 1976 Issue

THE FEBRUARY 1976 ISSUE CONTAINS THE FOLLOWING PAPERS

Hepatitis B surface antigen (HBsAg) in the liver of patients with hepatitis: a comparison with serological detection M. B. RAY, V. J. DESMET, J. FEVERY, J. DE GROOTE, A. F. BRADBURN, AND J. DESMYTER

Distribution patterns of hepatitis B surface antigen (HBsAg) in the liver of hepatitis patients M. B. RAY, V. J. DESMET, J. FEVERY, J. DE. GROOTE, A. F. BRADBURN, AND J. DESMYTER

Clinicopathological spectrum of late postpartum renal failure: two contrasting cases P. M. FORD, D. A. LEVISON, P. F. DOWN, AND J. B. MCCONNELL

A review and selection of simple laboratory methods used for the study of glycosaminoglycan excretion and the diagnosis of the mucopolysaccharidoses C. A. PENNOCK

Diagnostic potential of lysosomal hydrolases in body cavity effusions MALCOLM MCCORMACK

Re-evaluation of heat precipitation method for plasma fibrinogen estimation: effect of abnormal proteins and plasma viscosity LEOPOLD DINTENFASS AND SYLVIA KAMMER

Red cell aplasia following prolonged D-penicillamine therapy J. L. GOLLAN, SONAY HUSSEIN, A. V. HOFFBRAND, AND SHEILA SHERLOCK

Significance of subnormal red-cell folate in thalassaemia S. C. TSO

Weber-Christian panniculitis and auto-immune disease: a case report T. G. ALLEN-MERSH

A microplate enzyme-immunoassay for toxoplasma antibody A. VOLLER, D. E. BIDWELL, A. BARTLETT, D. G. FLECK, M. PERKINS, AND B. OLADEHIN

Comparison of four techniques for the routine diagnosis of *Trichomonas vaginalis* infection P. R. MASON, HEATHER SUPER, AND P. J. FRIPP

A comparison of two commercial methods for the identification of the Enterobacteriaceae—API 20E and the Enterotube—with conventional methods L. J. HAYEK AND G. W. WILLIS

Activity and interaction of trimethoprim and sulphamethoxazole against *Escherichia coli* D. GREENWOOD AND F. O'GRADY

Technical methods

Cone-biopsy histology report R. A. MCINROY, H. R. GRIMSHAW, AND P. HARRISON

Reliability of enzymatic lysozyme determinations T. TEPPER, S. MEIJER, AND GERA BROUWER-PIJPEN

Letters to the Editor

Book reviews

Copies are still available and may be obtained from the PUBLISHING MANAGER, BRITISH MEDICAL ASSOCIATION, TAVISTOCK SQUARE, LONDON WC1H 9JR, price £2.00, including postage