Carriage of group D streptococci in the human bowel

C. J. NOBLE

From the Area Public Health Laboratory, Dulwich Hospital, East Dulwich Grove, London SE22 8QF, UK

SUMMARY Sixty faecal samples, 39 from adults and 21 from neonates, were investigated by means of a simple scheme to isolate and identify any group D streptococci present. A number of differences were found between the two groups. Group D streptococci were absent from 43% of the neonates compared with only 13% of the adults; *Streptococcus bovis* was commoner in the neonates (23.8%) than in the adults (5%), and *Streptococcus faecium* was not isolated from any of the infant samples although it was common in the adult samples (25%). The viable counts of *Strep. faecium* were found to be, on average, 100-fold lower than those of *Streptococcus faecalis*.

The methods and results are discussed with reference to the clinical significance of group D streptococci, especially in bacterial endocarditis and carcinoma of the colon.

Group D streptococci form part of the normal gut flora of man and animals, but the distribution of the various species within the group is still imperfectly understood (Medrek and Barnes, 1962). Most studies show that *Streptococcus faecalis* and *Streptococcus faecium* are common human gut commensals, while *Streptococcus bovis* and other species are less frequently present. Recently, attention has been focused on *Strep. bovis*, which has been shown to be an important cause of endocarditis. In a three-year study of streptococcal isolates, Parker and Ball (1976) showed that *Strep. bovis* was associated with 17.3% of 317 cases of endocarditis, and that in the over-55 age group this organism was associated with nearly one-quarter of the cases.

More recently, Klein *et al.* (1977) have shown that the incidence of gut carriage of *Strep. bovis* is greatly increased in patients with carcinoma of the colon. Gross *et al.* (1975) devised a relatively simple scheme for identifying and speciating group D streptococci, which was shown by Facklam (1976) to give an accuracy of speciation of 98.5% in one large series.

The present study is an investigation of group D streptococci isolated from faecal samples from three groups of patients—hospital neonates, adult inpatients, and adult outpatients—at Dulwich Hospital. The streptococci were identified by means of

the above-mentioned scheme, and viable counts were performed.

Material and methods

Sixty faecal samples were examined: 21 were obtained from hospital neonates aged 6-7 days, 10 from adult inpatients, and 29 from adult outpatients.

The transit time for domiciliary specimens was a maximum of five hours, but hospital specimens were received more promptly. On arrival in the laboratory all specimens were refrigerated at 4°C and processed within three hours. A pea-sized lump of each sample was placed in a weighed sterile container of glycerol broth, homogenised on a vortex mixer, and stored for up to one month at -20°C. Frozen specimens were thawed at room temperature, and serial dilutions were made in Ringer's solution up to a dilution of 10⁻⁴ (Hewitt and Rigby, 1976). All the dilutions were immediately inoculated on to *M. enterococcus* Agar, a selective medium for group D streptococci (Becton, Dickinson & Co Ltd) in 0.1-ml volumes, by means of Pasteur pipettes calibrated to deliver 0.02 ml drops, and were spread with a glass rod. All dilution plates were incubated aerobically at 37°C for 48 hours when counts of each colonial type were made at the most convenient dilution. The minimum count detectable with this system is 100 organisms per gram of faeces. Counts were made on the

Received for publication 31 May 1978

assumption that each sample weighed 1 g, and a correction was made later for the actual weight of sample taken.

Single colonies were picked and inoculated on Horse Blood Agar plates (Oxoid) for purity. Subcultures were made from the pure cultures on to 40% bile aesculin slopes (prepared in the laboratory from Oxoid constituents) and to Robertson's Cooked Meat Broths (laboratory prepared from Lab M constituents).

Strains that grew on 40% bile aesculin, blackened the medium after 48 hours' incubation, were catalase-negative, and were Gram-positive oval cocci in pairs or chains in broth culture, were presumed to be group D streptococci and were submitted to further tests. Strains that failed to satisfy these criteria were discarded. All isolates were tested for their ability to (1) utilise pyruvate, (2) hydrolyse arginine, and (3) grow in 6.5% sodium chloride broth. The scheme of identification is set out in the Figure and is that described by Gross *et al.* (1975) and Facklam (1974).

UTILISATION OF PYRUVATE

Formula of medium:

Difco yeast extract 2.5 g
Difco tryptone 5.0 g
Dipotassium hydrogen phosphate 2.5 g
Sodium of pyruvate salt 5.0 g
Bromothymol blue 0.02 g
Agar 6.0 g
Distilled water 500 ml.

The medium was prepared as a slope in a bijou bottle. A positive reaction was indicated by the development of a yellow coloration after 48 hours' incubation at 37°C (Facklam, 1976).

SALT TOLERANCE

Five millilitres of broths containing 6.5% sodium chloride were inoculated heavily and incubated at 37°C for 48 hours. They were examined for turbidity with the naked eye; broths that were not turbid were Gram stained and, if no organisms were seen, the test was repeated (Cowan, 1974).

HYDROLYSIS OF ARGININE

Ammonia production was indicated by the development of a brown coloration on the addition of 0.25 ml of Nessler's Reagent to 5 ml of medium after five days' incubation at 37°C (method of Niven *et al.* (1942), see Cowan (1974)).

In addition to the tests described above, carbohydrate fermentation tests were performed on those isolates thought to be either *Strep. faecium* or *Strep. bovis*. Hiss' 1% Serum Sugars with Andrade's Indicator were inoculated and incubated for 48

Figure Scheme for identifying and speciating group D streptococci.

hours at 37°C. Fermentation of mannitol, sucrose, and arabinose distinguishes *Strep. faecium* from *Strep. faecium* var *durans*, and fermentation of lactose, inulin, and raffinose distinguishes *Strep. bovis* from *Streptococcus equinus*.

Finally, all the presumptive group D isolates were submitted to Lancefield grouping. The isolates were cultured in 50-ml volumes of glucose broth (Southern Group Laboratory) at 37°C for 18 hours. The supernatant was removed and the deposit autoclaved to extract the group D antigen (Hamilton, 1972). Group D antiserum was obtained from Wellcome Biological Reagents, Beckenham, Kent. Precipitation tests were performed by the layering method (Williams, 1958) in capillary tubes, and positive reactions occurring within five minutes were recorded. Extracts that failed to group were retested at a dilution of 1 in 2, 1 in 4, and 1 in 8 in order to overcome any prozone effect due to antigen excess. A minority of isolates failed to group by this procedure and were submitted to counter-current immunoelectrophoresis by the method of McIntyre (1978).

A number of lyophilised strains of group D streptococci were obtained from the National

Table 1 *List of control strains*

<i>NCTC strains</i>	
8133	<i>Strep. bovis</i>
8307	<i>Strep. faecium</i> var <i>durans</i>
775	<i>Strep. faecalis</i>
2705	<i>Strep. faecalis</i> var <i>liquefaciens</i>
5957	<i>Strep. faecalis</i> var <i>zymogenes</i>
7171	<i>Strep. faecium</i>
<i>Streptococcal Reference Laboratory strains</i>	
A	<i>Strep. bovis</i>
B	<i>Strep. faecalis</i>
C	<i>Strep. faecium</i>
D	<i>Strep. bovis</i>

Collection of Type Cultures and some further isolates from the Streptococcal Reference Laboratory. These strains were used to provide positive and, where appropriate, negative controls for the biochemical tests (Table 1).

Results

All the control strains were tested alongside the faecal isolates and gave appropriate results for (1) hydrolysis of aesculin in 40% bile, (2) tolerance of 6.5% sodium chloride, (3) pyruvate reduction, and (4) arginine hydrolysis. One control *Strep. bovis* failed to ferment inulin on repeated testing. Serological grouping was performed with the following results:

<i>Number of strains tested</i>	<i>Group D status</i>
5 controls	Confirmed
46 faecal strains	43 confirmed

Three faecal isolates failed to group with the D antiserum by precipitation or immunoelectrophoresis.

On the basis of the scheme described under Material and methods, isolates were speciated into *Strep. faecalis*, *Strep. faecium*, *Strep. avium*, *Strep. bovis*, and *Strep. equinus*. No *Strep. faecium* var *durans* was isolated, and strains that could not be identified by this method were classified as unidentified group D streptococci.

Table 2 *Species of group D streptococci isolated from faeces of hospital neonates, adult inpatients, and adult outpatients*

<i>Patient group</i>	<i>No. of faeces examined</i>	<i>No. of faecal specimens containing various species of group D streptococci</i>						
		<i>Strep. faecalis</i>	<i>Strep. faecium</i>	<i>Strep. avium</i>	<i>Strep. bovis</i>	<i>Strep. equinus</i>	<i>Not identified</i>	<i>Any group strep.</i>
Neonates	21	10 (47.6)	0	0	5 (23.8)	0	1 (4.8)	12 (57.1)
Adult inpatients	10	8 (80)	3 (30)	1 (10)	0	0	1 (10)	9 (90)
Adult outpatients	29	14 (48.2)	12 (41.3)	2 (6.9)	2 (6.9)	2 (6.9)	6 (20.7)	25 (86.2)

Percentages are given in parentheses.

Table 2 shows the relative frequency of isolation of the various species of group D streptococci from the three groups of patients, and Table 3 shows the distribution of viable counts in the 43 specimens examined. The viable counts for the neonates and adult groups are combined, as insufficient data are available for separate tables.

One or more species of group D streptococci were isolated from 46 of the 60 samples studied. Seventeen samples yielded two different species, and two samples yielded three species.

Table 4 shows the selectivity of *M. enterococcus* Agar for group D streptococci.

Table 3 *Viable counts of various species of group D streptococci from all groups of patients (expressed as organisms per gram of faeces)*

<i>Species</i>	<i>No. of isolates falling in each log range</i>							<i>No. of samples studied</i>
	10^8	10^7	10^6	10^5	10^4	10^3	10^2	
<i>Faecalis</i>	1	5	4	5	1	2	1	19
<i>Faecium</i>	0	1	0	3	2	2	2	10
<i>Avium</i>	0	0	0	1	1	0	0	2
<i>Bovis</i>	0	1	2	1	0	0	0	4
<i>Equinus</i>	0	0	0	1	1	0	0	2
Unidentified	0	2	0	1	1	2	0	6

NB 1 *Strep. faecium* not isolated from neonates.

2 *Strep. bovis* counts all on neonates.

3 *Strep. faecalis* counts for adults and neonates not significantly different.

Table 4 *Selectivity of M. enterococcus Agar for group D streptococci*

Total number of isolates from all specimens	80
Total number of group D isolates	67
Total number of non group D isolates	13
% of non group D isolates	16.25

Discussion

A variety of group D streptococci may be isolated from human faeces (see Table 2). In this study, one or two species were commonly isolated from a

single sample. The predominant species was *Strep. faecalis*, which was cultured from 53% of the samples studied. This was followed by *Strep. faecium* (25%) and *Strep. bovis* (11.6%). Other group D streptococci were much less common. Thirteen per cent of the samples produced strains that could not be identified by means of the scheme described. Of these eight unidentified strains, all but one belonged to Lancefield group D. Five of them gave a positive arginine reaction but failed to grow in 6.5% sodium chloride broth or to utilise pyruvate.

In this study, 16% of all the isolates from *M. enterococcus* Agar were presumed not to be group D streptococci because they failed to blacken bile aesculin medium. The bile aesculin test is a reliable presumptive test of group D status, as shown by Facklam *et al.* (1974), who tested 920 strains of group D streptococci and found that 99% blackened the medium. Gross *et al.* (1975) found that 99.7% of 844 strains of group D streptococci were bile aesculin positive.

Where mixed cultures of group D and non group D isolates occurred, the presence of the latter did not make counting difficult. All colonial variants were submitted to bile aesculin testing, as it is not possible to distinguish group D streptococci on *M. enterococcus* Agar by colonial morphology alone. These findings are in agreement with those of Pavlova *et al.* (1972), who found that 18.4% of their isolates from faeces, sewage, and food samples, plated on *M. enterococcus* Agar, were not group D streptococci, 6.2% being either *Streptococcus mitis* or *Streptococcus salivarius*. Raibaud *et al.* (1961) reported overgrowth of lactobacilli when pig caecal samples were plated on this medium.

According to Pavlova *et al.* (1972), the yield of group D streptococci from faecal samples is less with *M. enterococcus* Agar than with the thallose acetate medium of Barnes (1956), 3.3×10^5 organisms per gram as compared with 8.3×10^6 organisms per gram. However, Burkwall and Hartman (1964) found that the range of percent recovery of group D streptococci from frozen foods plated on *M. enterococcus* Agar was 70-500%, assuming 100% recovery from Barnes' medium. Pavlova *et al.* (1972) used small inocula, 0.1-ml volumes of 10^{-4} faecal dilutions, and did not report any relative inhibition.

My findings are broadly in agreement with those of previous studies (Sabbaj *et al.*, 1971; van der Wiel-Korstanje and Winkler, 1975). There were, however, several differences between the flora of the neonates and of the adults that have not previously been recorded. *Strep. faecium* was isolated from 15 of the 39 adult faeces (38.5%)

whereas it was not isolated from any of the 21 neonatal samples. *Strep. bovis* was more frequently isolated from the neonatal group, 5 out of 21 (23.8%) compared with 2 out of 39 (5%) for the adults. Nine of the neonatal samples produced no group D streptococci (43%) compared with only five of the adult samples (13%). The adult group produced a wider range of streptococci, but this may have been merely a reflection of the larger number of adult specimens examined (39:21).

In over half of the samples yielding *Strep. faecalis*, viable counts of the organism were within the range 10^6 - 10^9 organisms per gram, whereas in the samples yielding *Strep. faecium*, viable counts of the organism were in all cases below 10^6 organisms/g, and two-thirds of samples had counts below 10^5 organisms/g.

I acknowledge the help and advice given by Dr M. T. Parker and Dr C. Dulake in the preparation of this paper.

References

- Barnes, E. M. (1956). Methods for the isolation of faecal streptococci (Lancefield Group D) from bacon factories. *Journal of Applied Bacteriology*, **19**, 193-203.
- Burkwall, M. K., and Hartman, P. A. (1964). Comparison of direct plating media for the isolation and enumeration of enterococci in certain frozen foods. *Applied Microbiology*, **12**, 18-23.
- Cowan, S. T., ed. (1974). *Cowan and Steel's Manual for the Identification of Medical Bacteria*, 2nd edition. Cambridge University Press, London.
- Facklam, R. R., Padula, J. F., Thacker, L. G., Wortham, E. C., and Sconyers, B. J. (1974). Presumptive identification of group A, B, and D streptococci. *Applied Microbiology*, **27**, 107-113.
- Facklam, R. R. (1976). A review of the microbiological techniques for the isolation and identification of streptococci. *CRC Critical Reviews in Clinical Laboratory Sciences*, **6**, 287-317.
- Gross, K. C., Houghton, M. P., and Senterfit, L. B. (1975). Presumptive speciation of *Streptococcus bovis* and other group D streptococci from human sources by using arginine and pyruvate tests. *Journal of Clinical Microbiology*, **1**, 54-60.
- Hamilton, W. J. (1972). Streptococcal grouping: heat extraction of C substance from beta haemolytic streptococci. *Medical Laboratory Technology*, **29**, 385-388.
- Hewitt, J. H., and Rigby, J. (1976). Effect of various milk feeds on numbers of *Escherichia coli* and *Bifidobacterium* in the stools of newborn infants. *Journal of Hygiene*, **77**, 129-139.
- Klein, R. S., Recco, R. A., Catalano, M. T., Edberg, S. C., Casey, J. I., and Steigbigel, N. H. (1977). Association of *Streptococcus bovis* with carcinoma of the colon. *New England Journal of Medicine*, **297**, 800-802.

- McIntyre, M. (1978). Detection of capsulated *Haemophilus influenzae* in chest infections by counter current immunoelectrophoresis. *Journal of Clinical Pathology*, **31**, 31-34.
- Medrek, T. F., and Barnes, E. M. (1962). The influence of the growth medium on the demonstration of a group D antigen in faecal streptococci. *Journal of General Microbiology*, **28**, 701-709.
- Niven, C. F. Jr., Smiley, K. L., and Sherman, J. M. (1942). The hydrolysis of arginine by streptococci. *Journal of Bacteriology*, **43**, 651-660.
- Parker, M. T., and Ball, L. C. (1976). Streptococci and aerococci associated with systemic infection in man. *Journal of Medical Microbiology*, **9**, 275-302.
- Pavlova, M. T., Brezenski, F. T., and Litsky, W. (1972). Evaluation of various media for isolation, enumeration and identification of fecal streptococci from natural sources. *Health Laboratory Science*, **9**, 289-298.
- Raibaud, P., Caulet, M., Galpin, J. V., and Mocquot, G. (1961). Studies on the bacterial flora of the alimentary tract of pigs. II. Streptococci: selective enumeration and differentiation of the dominant group. *Journal of Applied Bacteriology*, **24**, 285-306.
- Sabbaj, J., Suttar, V. L., and Finegold, S. M. (1971). Comparison of selective media for isolation of presumptive group D streptococci from human feces. *Applied Microbiology*, **22**, 1008-1011.
- Van der Wiel-Korstanje, J. A. A., and Winkler, K. C. (1975). The faecal flora in ulcerative colitis. *Journal of Medical Microbiology*, **8**, 491-501.
- Williams, R. E. O. (1958). Laboratory diagnosis of streptococcal infections. *Bulletin of the World Health Organisation*, **19**, 153-176.

Requests for reprints to: Dr C. J. Noble, Department of Medical Microbiology, King's College Hospital Medical School, Denmark Hill, London, SE5 8RX, UK.

Reports and Bulletins prepared by the Association of Clinical Biochemists

The following reports and bulletins are published by the Association of Clinical Biochemists. They may be obtained from The Publishing Department, British Medical Journal (ACB Technical Bulletins), B.M.A. House, Tavistock Square, London WC1H 9JR. Overseas readers should remit by British Postal or Money Order.

SCIENTIFIC REVIEWS (price £1.00/\$2.00 each)

- 1 The assessment of thyroid function March 1971
F. V. FLYNN and J. R. HOBBS
- 2 Renal function tests suitable for clinical practice
January 1972 F. L. MITCHELL, N. VEALL, and R. W. E. WATTS
- 3 Biochemical tests for the assessment of fetoplacental
function May 1975 C. E. WILDE and R. E. OAKLEY
- 4 Test of exocrine pancreatic function March 1977
A. H. GOWENLOCK

TECHNICAL BULLETINS (price £1.00/\$2.00 each)

- 22 Bilirubin standards and the determination of bilirubin
by manual and technicon AutoAnalyzer methods
January 1971 BARBARA BILLING, RUTH HASLAM, and N.
WALD
- 23 Interchangeable cells for spectrophotometers and
fluorimeters September 1971 S. S. BROWN and A. H.
GOWENLOCK
- 24 Simple tests to detect poisons March 1972 B. W.
MEADE *et al.*
- 25 Blood gas analysers May 1972 K. DIXON
- 26 Kits for enzyme activity determination September
1972 S. B. ROSALKI and D. TARLOW
- 27 Assessment of pumps suitable for incorporation into
existing continuous flow analytical systems November
1972 A. FLECK *et al.*

- 28 Routine clinical measurements of transferrin in
human serum September 1973 K. DIXON
- 29 Control materials for clinical biochemistry (5th
edition) September 1973 J. F. STEVENS
- 30 Notes on the quality of performance of serum
cholesterol assays September 1973 S. S. BROWN
- 31 Determination of uric acid in blood and in urine
July 1974 R. W. E. WATTS
- 32 A survey of amino acid analysers readily available in
the United Kingdom September 1974 J. E. CARLYLE and
P. PURKISS
- 33 Definitions of some words and terms used in auto-
mated analysis November 1974 A. FLECK, R. ROBINSON,
S. S. BROWN, and J. R. HOBBS
- 34 Measurement of albumin in the sera of patients
January 1975 LINDA SLATER, P. M. CARTER, and J. R.
HOBBS
- 35 Investigation of the validity of temperature correction
factors for serum aspartate and alanine transaminases
March 1975 S. B. ROSALKI *et al.*
- 36 Factors influencing the assay of creatinine November
1975 J. G. H. COOK
- 37 A survey of enzyme reaction rate analysers readily
available in the United Kingdom July 1977 R. A.
SAUNDERS and R. F. BURNS
- 38 Transport of specimens for clinical chemistry an-
alysis November 1977 P. WILDING, J. F. ZILVA, and
C. E. WILDE
- 39 A scheme for the evaluation of diagnostic kits May
1978 P. H. LLOYD