Urinary infection with coagulase-negative staphylococci in a teaching hospital

T. L. SHRESTHA¹ AND J. H. DARRELL

From the Department of Bacteriology, Royal Postgraduate Medical School, Hammersmith Hospital, Du Cane Road, London W12, UK

SUMMARY In the eight-month period of study of all urine samples processed in our routine laboratory, only 85 out of 12 152 specimens yielded a bacteriologically significant growth of either *Staphylococcus epidermidis* or micrococci. Their growth on MacConkey medium was strictly comparable to that on cysteine lactose electrolyte-deficient (CLED) media. Most micrococci isolated were from urine samples of non hospitalised women patients, were resistant to a novobiocin (5 µg) disc, and belonged to Baird Parker type 3. *Staph. epidermidis* came mainly from postoperative surgical in-patients. Their antibiotic sensitivity patterns are variable whereas micrococci are fully sensitive to all urinary antibiotics. We agree that the use of a novobiocin (5 µg) disc for provisional identification of micrococci and *Staph. epidermidis* is simple and practical for a busy routine diagnostic laboratory. The use of more extensive systems to biotype these organisms in a routine laboratory is not practical and not relevant to patient management.

The undoubted role of catalase-positive, coagulase-negative Gram-positive cocci, *Staphylococcus epidermidis*, and *Micrococcus* species in causing urinary tract infection has gradually become accepted in recent years. These organisms did not appear to occur commonly in our routine practice, though other workers have reported an incidence ranging from complete absence (Arneil *et al.*, 1970) to 30% of isolates (Gillespie *et al.*, 1978). We therefore decided to collect and identify all these organisms occurring in urine samples reaching our diagnostic laboratory from inside and outside the hospital in order to establish their frequency and to assess the suitability of our methods for isolation and identification.

Material and methods

Urine samples received in the routine laboratory are cultured without delay or stored for up to 3 hours at 2-4°C on a CHEMLAB Bench Cooler. The well-mixed unspun urine is streaked onto a MacConkey plate (Oxoid CM76) using a calibrated loop delivering 0.003 ml of urine. For the purposes of this investigation duplicate cultures were prepared

¹Present address: Pathology Department, Royal Infirmary, Sunderland.

Received for publication 4 September 1978

on plates of Cysteine Lactose Electrolyte Deficient (CLED) Agar, (Oxoid CM301). These were prepared in an exactly comparable manner to the McConkey plates, four specimens being cultured on one plate. A growth of 300 colonies or more of an organism is considered bacteriologically significant. Those yielding 30 colonies or less are discarded.

A wet preparation of the spun deposit (centrifuged for 10 minutes at 2000 rpm) is examined microscopically for formed elements, leucocytes, red blood cells, and bacteria. Those samples which have got pus cells, bacteria, or red cells require primary sensitivity tests, to sulphafurazole (200 µg), trimethoprim (2.5 µg), ampicillin (25 µg), nitrofurantoin (50 µg), and cephaloridine (25 µg). These tests are carried out on lysed blood agar plates ((Iso-Sensitest) Oxoid CM471 with 5% horse blood lysed with saponin) using Stokes' method and *Escherichia coli* NCTC 10418 as the control organism. After overnight incubation at 37°C all the cultures on CLED medium were read by one of us (TLS), and the results were compared with those of the routine cultures set up at the same time on MacConkey medium. Cultures yielding a bacteriologically significant growth of Gram-positive cocci, as defined above, were tested for coagulase production by the tube method of Gillespie (1943); catalase production was tested by the slide method

(Cruickshank *et al.*, 1975). Catalase-positive, coagulase-negative strains were biotyped using the methods of Baird-Parker (1963). In addition, strains were tested for sensitivity to a 5 µg novobiocin disc (Mitchell, 1968) as a diagnostic sensitivity test. These tests were carried out in parallel on Mueller-Hinton agar (Oxoid CM337), Wellcotest sensitivity test agar (Wellcome Reagents Limited, CM53), and Iso-Sensitest agar (Oxoid CM471) without added blood. Sensitive organisms gave a zone of inhibition measuring 8 mm or more from the edge of the disc.

Results

During the eight-month period October 1977-May 1978, our diagnostic laboratory processed 12 152 urine specimens. They were received from hospital ward patients, from those attending outpatient clinics, and from general practitioners. Of these specimens, 1186 gave significant bacteriuria due to all kinds of bacteria. Only 85 samples yielded a bacteriologically significant growth of Gram-positive, coagulase-negative, catalase-positive cocci (Table 1). Of these 85 samples, 17 were from non-hospitalised patients, all but one of them being female. The rest of the specimens were from hospitalised patients, that is, inpatients and outpatients (Table 2). Cultures on MacConkey agar and CLED medium gave strictly comparable results. Each of the 85 strains was recovered on both media, giving approximately equal numbers of colonies of the same size.

Of the 85 strains tested, 15 only were found to be micrococcus type 3, now re-classified as *Staphylococcus saprophyticus* biotype 3 on the basis of guanine and cytosine content of DNA (Buchanan and Gibbons, 1974). All these isolates were resistant to a 5 µg novobiocin disc.

No other micrococcus biotypes were encountered

Table 1 Urine specimens, October 1977-May 1978

Total number processed in the laboratory	12 152
Significant bacteriuria due to all kinds of bacteria	1186 (9.76%)
Significant bacteriuria due to coagulase-negative Gram-positive cocci out of all significant bacteriuria	85 (7.16%)

Table 2 Staphylococcal urinary infection in three groups of patients

	Hospital inpatients		Hospital outpatients		Non-hospitalised* patients	
<i>Staph. epidermidis</i> (<i>Staph. albus</i>)	Male	40	Male	4	Male	1
	Female	16	Female	3	Female	6
		56		7		7
<i>Staph. saprophyticus</i> (Micrococcus type 3)	Male	2	Male	—	Male	0
	Female	3	Female	—	Female	10
		5		0		10
Totals		61		7		17
						85

*General practitioner patients and patients attending casualty or antenatal clinic.

in this study. Ten of the 15 micrococci occurred in the group of patients who had never been hospitalised, that is, samples submitted by family practitioners, from the casualty department, and from the antenatal clinic. The other five occurred in hospitalised patients, accounting for only 8% of isolates in this group (Table 2).

All the hospital outpatient isolates were of *Staph. epidermidis* and occurred in patients who had recently left hospital (Table 2).

All micrococci were found to be sensitive to all antimicrobial agents tested in the primary sensitivity tests. The pattern of sensitivities is different with staphylococci and very variable (Table 2).

Discussion

The slow recognition of coagulase-negative staphylococci, especially micrococci (*Staph. saprophyticus*), as urinary pathogens in hospital practice is hardly surprising. In our series, they account for only 7.16% of the significant isolates. We are able to confirm the findings of other workers (Mitchell, 1968; Kerr, 1973; Maskell, 1974; Meers *et al.*, 1975; Sellin *et al.*, 1975; Pead *et al.*, 1977) that they account for a significant proportion of primary urinary infections, Gram-positive cocci occurring in women outside hospital, whereas in a hospital population their contribution is trivial. They do not commonly cause postoperative infections in males; these frequently follow surgery and are caused by *Staph. epidermidis*. The micrococci isolated by us were all type 3, now reclassified as *Staph. saprophyticus* biotype 3.

We were able to show that all strains of *Staph. epidermidis* and micrococci (*Staph. saprophyticus*) isolated by us from urine samples grew well on our routine MacConkey media. Hence, in our view, there is no need to include a special medium such as CLED when screening urines in order to isolate these organisms, provided the MacConkey medium in use has similar selective properties.

The biochemical methods of Baird-Parker (1963) are relatively time-consuming for the routine laboratory. However, they may be useful in associating

Table 3 Sensitivity patterns of micrococci type 3 and other coagulase-negative staphylococci from urine

Organism	Proportion of strains sensitive to:				
	Sulphonamide	Trimethoprim	Ampicillin	Nitrofurantoin	Cephaloridine
<i>Staphylococcus</i>	63/69	37/69	37/66*	65/67*	61/64*
<i>Micrococcus</i> type 3	14/14	14/14	14/14	14/14	14/14

*Low total numbers due to failure to perform test.

certain biotypes with specific types of infection. Care has to be taken in their correct performance. In particular, we had difficulty in interpreting some oxidation-fermentation tests when using the medium as described by Baird-Parker. Clear results were obtained when the concentration of the indicator-bromocresol purple was reduced to 0.002%, and soft paraffin was used to seal the anaerobic tubes in preference to sterile liquid paraffin (Chalmers, 1972). In contrast, novobiocin (5 µg) resistance is a simple test for provisional identification of micrococci (Mitchell, 1968). It gave comparable results irrespective of the medium used, provided lysed blood was not incorporated. Its addition had the effect of stimulating growth and giving reduced zones with some staphylococcal isolates.

In spite of the fact that not all biotypes of micrococci are novobiocin-resistant (Pead *et al.*, 1977), it appears still to be a useful screening test for routine laboratory use (Sellin *et al.*, 1975) to differentiate *Staph. epidermidis* and *Staph. saprophyticus*. Mitchell (1968) isolated four biotypes of *Staph. epidermidis*—II, IV, V, and VI—from urines, as did Pead *et al.* (1977). The same four types were isolated in this study. The only difference between our observations and those of others was that biotype VI was found to be commoner than biotype V in our series. These biotypes were isolated from hospital patients only in our study. Most of them came from surgical wards, and surgical intervention in and around the urinary tract seems to predispose to infection with these organisms. Most of the patients involved were males, unlike the micrococcal infections that occurred in females.

Although the numbers are small, it appears that at present micrococcus type 3 remains a fully antibiotic-sensitive organism. When it occurs as a cause of primary urinary infection it is likely to respond to any of the commonly prescribed urinary antibacterial antibiotics.

Staph. epidermidis associated with postoperative urinary tract infection, however, has a very variable sensitivity pattern. This, plus the fact that these infections occur in relation to urological surgery and the possible failure to establish urinary drainage, make these infections more difficult to manage.

Reviewing the sensitivity patterns of all our isolates (Table 3), it is surprising that so many strains of both micrococci and *Staph. epidermidis* are fully sensitive *in vitro* to sulphonamides and equally surprising that so many of the staphylococci are resistant to trimethoprim. This may reflect heavy usage of cotrimoxazole in hospital patients. In view of the complicated nature of many of the postoperative staphylococcal infections, use of sulphonamides alone would hardly recommend itself as a treatment likely to succeed. The fact that half our strains were resistant to ampicillin is not now surprising, and all strains are, of course, resistant to nalidixic acid. In view of the relative resistance to mecillinam, the place of this antibiotic in treating staphylococcal urinary infections must remain in doubt in spite of the high urinary levels attained. This means that the oral treatment of first choice for such infections would have to be an oral cephalosporin, with nitrofurantoin as a possible alternative.

We thank all the technical staff working in the urine section for their co-operation, Harry Edwards and his staff for preparation of media, and Miss A. Kirk for typing this manuscript.

References

- Arneil, G. C., McAllister, T. A., and Kay, P. (1970). Detection of bacteriuria at room-temperature. *Lancet*, **1**, 119-121.
- Baird-Parker, A. C. (1963). A classification of micrococci and staphylococci based on physiological and biochemical tests. *Journal of General Microbiology*, **30**, 409-427.
- Buchanan, R. E., and Gibbons, N. E. (Eds.) (1974). *Bergey's Manual of Determinative Bacteriology*, 8th edition. Williams and Wilkins, Baltimore.
- Chalmers, A. (1972). A modification of the oxidation/fermentation test for the classification of Micrococcaceae. *Medical Laboratory Technology*, **29**, 379-384.
- Cruikshank, R., Duguid, J. P., Marmion, B. P., and Swain, R. H. A. (1975). *Medical Microbiology*, Volume II, 12th edition. Churchill Livingstone, Edinburgh, London and New York.
- Gillespie, E. H. (1943). The routine use of the coagulase

- test for staphylococci. *Monthly Bulletin of the Emergency Public Health Laboratory Service*, **2**, 12-22.
- Gillespie, W. A., Sellin, M. A., Gill, P., Stephens, M., Tuckwell, L. A., and Hilton, A. L. (1978). Urinary tract infection in young women, with special reference to *Staphylococcus saprophyticus*. *Journal of Clinical Pathology*, **31**, 348-350.
- Kerr, H. (1973). Urinary infection caused by Micrococcus subgroup 3. *Journal of Clinical Pathology*, **26**, 918-920.
- Maskell, R. (1974). Importance of coagulase-negative staphylococci as pathogens in the urinary tract. *Lancet*, **1**, 1155-1158.
- Meers, P. D., Whyte, W., and Sandys, G. (1975). Coagulase-negative staphylococci and micrococci in urinary tract infections. *Journal of Clinical Pathology*, **28**, 270-273.
- Mitchell, R. G. (1968). Classification of *Staphylococcus albus* strains isolated from the urinary tract. *Journal of Clinical Pathology*, **21**, 93-96.
- Pead, L., Crump, J., and Maskell, R. (1977). Staphylococci as urinary pathogens. *Journal of Clinical Pathology*, **30**, 427-431.
- Sellin, M. A., Cooke, D. I., Gillespie, W. A., Sylvester, D. G. H., and Anderson, J. D. (1975). Micrococcal urinary-tract infections in young women. *Lancet*, **2**, 570-572.

Requests for reprints to: Dr J. H. Darrell, Department of Bacteriology, Royal Postgraduate Medical School, Hammersmith Hospital, Ducane Road, London W12 0HS, UK.

Reports and Bulletins prepared by the Association of Clinical Biochemists

The following reports and bulletins are published by the Association of Clinical Biochemists. They may be obtained from The Publishing Department, British Medical Journal (ACB Technical Bulletins), B.M.A. House, Tavistock Square, London WC1H 9JR. Overseas readers should remit by British Postal or Money Order.

SCIENTIFIC REVIEWS (price £1.00/\$2.00 each)

- 1 The assessment of thyroid function March 1971
F. V. FLYNN and J. R. HOBBS
- 2 Renal function tests suitable for clinical practice
January 1972 F. L. MITCHELL, N. VEALL, and R. W. E. WATTS
- 3 Biochemical tests for the assessment of fetoplacental
function May 1975 C. E. WILDE and R. E. OAKEY
- 4 Test of exocrine pancreatic function March 1977
A. H. GOWENLOCK

TECHNICAL BULLETINS (price £1.00/\$2.00 each)

- 22 Bilirubin standards and the determination of bilirubin
by manual and technicon AutoAnalyzer methods
January 1971 BARBARA BILLING, RUTH HASLAM, and N. WALD
- 23 Interchangeable cells for spectrophotometers and
fluorimeters September 1971 S. S. BROWN and A. H. GOWENLOCK
- 24 Simple tests to detect poisons March 1972 B. W. MEADE *et al.*
- 25 Blood gas analysers May 1972 K. DIXON
- 26 Kits for enzyme activity determination September
1972 S. B. ROSALKI and D. TARLOW
- 27 Assessment of pumps suitable for incorporation into
existing continuous flow analytical systems November
1972 A. FLECK *et al.*

- 28 Routine clinical measurements of transferrin in
human serum September 1973 K. DIXON
- 29 Control materials for clinical biochemistry (5th
edition) September 1973 J. F. STEVENS
- 30 Notes on the quality of performance of serum
cholesterol assays September 1973 S. S. BROWN
- 31 Determination of uric acid in blood and in urine
July 1974 R. W. E. WATTS
- 32 A survey of amino acid analysers readily available in
the United Kingdom September 1974 J. E. CARLYLE and
P. PURKISS
- 33 Definitions of some words and terms used in auto-
mated analysis November 1974 A. FLECK, R. ROBINSON,
S. S. BROWN, and J. R. HOBBS
- 34 Measurement of albumin in the sera of patients
January 1975 LINDA SLATER, P. M. CARTER, and J. R. HOBBS
- 35 Investigation of the validity of temperature correction
factors for serum aspartate and alanine transaminases
March 1975 S. B. ROSALKI *et al.*
- 36 Factors influencing the assay of creatinine November
1975 J. G. H. COOK
- 37 A survey of enzyme reaction rate analysers readily
available in the United Kingdom July 1977 R. A. SAUNDERS and R. F. BURNS
- 38 Transport of specimens for clinical chemistry an-
alysis November 1977 P. WILDING, J. F. ZILVA, and
C. E. WILDE
- 39 A scheme for the evaluation of diagnostic kits May
1978 P. H. LLOYD