Meningitis caused by *Pseudomonas paucimobilis*

V. HAJIROUSSOU¹, B. HOLMES², J. BULLAS¹, AND C. A. PINNING²

From ¹Manor Hospital, Moat Road, Walsall, West Midlands WS2 9PS
and the ²National Collection of Type Cultures, Central Public Health Laboratory,
Colindale, London NW9 5HT, UK

SUMMARY This appears to be the first report of meningitis due to *Pseudomonas paucimobilis* and the first report of a clinically significant isolate of this species in the UK. Characteristics by which the species may be recognised are given, and attention is drawn to the possible confusion of *Ps. paucimobilis* with other yellow-pigmented pseudomonads and *Flavobacterium* species.

Pseudomonas paucimobilis is a species described only recently (Holmes *et al.*, 1977). The strains on which the description of the species was based had been recovered largely from human clinical specimens and the hospital environment, but none was known to be the causative agent of infection. Although the clinical significance of *Ps. paucimobilis* remained unknown we now report what is, as far as we know, the first case of meningitis due to this species.

Case report

A 39-year-old male epileptic was admitted on 24 June 1978. He had had epilepsy for three years and was currently treated with phenobarbitone 60 mg and phenytoin 100 mg three times a day. He complained of headache for two days and had started to have convulsions on the day of admission. This was controlled with a single dose of 20 g diazepam intravenously. However, he remained very drowsy 6 hours later, and the axillary temperature was found to be 39.2°C (103°F). Clinical examination showed neck stiffness and a positive Kernig's sign.

A lumbar puncture was performed and turbid fluid obtained, giving the following results: white cells $0.2 \times 10^9/l$ (200/mm³), 95% lymphocytes, 5% neutrophils; protein 0.4 g/l (40 mg/100 ml), glucose 3.7 mmol/l (67.2 mg/100 ml); the blood glucose level was 4.7 mmol/l (85 mg/100 ml). Gram and Ziehl-Neelsen films were initially reported as negative. However, 48 hours later repeat examination of the original Gram-stained film revealed numerous Gram-negative rods. Other tests included haemoglobin 15.4 g/dl, white cell count $12.8 \times 10^9/l$ (12 800/mm³), 82% neutrophils showing toxic granulation and a shift to the left. A chest film was normal.

In view of the initial findings, tuberculous meningitis had to be considered and, because of the clinical state of the patient, treatment was started immediately with streptomycin 1 g daily intramuscularly, rifampicin 600 mg daily orally, and isoniazid 600 mg daily intramuscularly. The patient rapidly improved and became apyrexial within 30 hours.

Treatment was continued for four days and he was discharged symptom-free on the tenth day.

BACTERIOLOGY

Cultures of cerebrospinal fluid and blood, taken before the start of treatment, were incubated aerobically at 37°C on blood agar. After 48 hours cultures of both specimens yielded a moderately heavy pure growth of a yellow-pigmented, non-fermentative, Gram-negative, rod-shaped bacterium which could not be identified by routine laboratory tests. The isolates were sensitive to rifampicin.

Three isolates, two from cerebrospinal fluid and one from blood, were submitted to the National Collection of Type Cultures for computer-assisted identification. There, a set of 68 characteristics were determined for each isolate using methods described previously (Holmes *et al.*, 1975). In these tests the three isolates yielded identical results (Table), thereby indicating that the isolates represented a single strain. On the results of these characteristics, in conjunction with an unpublished probability matrix, the isolates were identified as *Ps. paucimobilis*. The isolates proved typical of the species except in their ability to produce an alkaline reaction on Christensen's citrate medium and in their inability to produce acid from salicin in ammonium salt sugar medium.

Cultures of swabs taken from the nose, throat, ear, and axilla of staff examining specimens in the laboratory, and from the houseman who performed the lumbar puncture and took blood for culture

Table Characteristics of the three isolates of *Pseudomonas paucimobilis* examined

<i>Isolates positive in:</i>			
Acid from the following ammonium salt sugars:	lactose maltose raffinose sucrose trehalose xylose Aesculin hydrolysis	Alkali production on Christensen's citrate agar Catalase production Cytochrome oxidase production Deoxyribonuclease production Growth at 37°C Growth at room temperature (18°C-22°C) Growth on β -hydroxybutyrate	Poly- β -hydroxybutyrate inclusion granules Production of yellow pigment Tween 20 hydrolysis Tween 80 hydrolysis Tyrosine hydrolysis β -galactosidase production
<i>Isolates negative in:</i>			
Acid from the following ammonium salt sugars:	Acid from glucose (10 g/100 ml) Acid from lactose (10 g/100 ml) Adonitol dulcitol glycerol inositol mannitol rhamnose salicin sorbitol	Growth at 42°C Growth on cetrimide agar Growth on MacConkey agar Growth on Simmons' citrate agar Hugh & Leifson O-F test Hydrogen sulphide production† Indole production KCN tolerance Lysine decarboxylase Malonate utilisation Motility‡	Nitrate reduction Nitrite reduction Opalescence on lecithovitellin agar Ornithine decarboxylase Phenylalanine deamination Pigment production on tyrosine agar Reduction of selenite (0.4 g/100 ml) Starch hydrolysis Urease production 3-ketolactose production
Acid from PWS glucose	Gas from PWS glucose Gelatinase production* Gluconate oxidation Growth at 5°C		

PWS = peptone water sugar.

*By stab and plate methods.

†By both lead acetate paper and triple sugar iron agar methods.

‡At both 37°C and room temperature (18°C-22°C).

from the patient described above, proved negative for *Ps. paucimobilis*.

Discussion

Two of the strains on which the original description of *Ps. paucimobilis* was based were representative of a group that had been designated group IIk, biotype 1 by workers at the Center for Disease Control, Atlanta, USA (Tatum *et al.*, 1974). This implies that group IIk, biotype 1 and *Ps. paucimobilis* are the same taxon, and thus, even before the description of *Ps. paucimobilis*, isolates belonging to the species were being recognised in human clinical material, principally from blood, environmental sources, spinal fluid, urine, and various wounds and abscesses (Tatum *et al.*, 1974). Although they would have been labelled unclassified non-fermenters when first received, after several isolates had been collected, strains were recognised as belonging together in a distinct taxon, provisionally as group IIk, biotype 1, but later as *Ps. paucimobilis*. However, because initial isolates of the taxon were labelled only as unclassified non-fermenters, clinical details were not requested on receipt of the strains, and consequently none was known to be implicated as a cause of infection. Also group IIk, biotype 1 was little recognised outside the USA, but since the naming of *Ps. paucimobilis* the species has become more widely known and recognised in clinical laboratories, and clinically significant isolates have been recognised: from a leg ulcer in Australia (Peel *et al.*, 1979) and from a septicaemia in the USA (Slotnick *et al.*, 1979). The case herein reported appears to be the

first of a clinically significant isolate of this species in the UK and the first report of this species as a cause of meningitis.

Ps. paucimobilis, because of its yellow pigment and because motility is difficult to demonstrate (room temperature in a hanging-drop preparation is recommended, and even then only a very low proportion of the cells may be actively motile), is most likely to be confused with *Flavobacterium* spp. However, *Flavobacterium* spp. usually grow on MacConkey agar, produce caseinase and gelatinase, and show resistance to carbenicillin and gentamicin, characters rarely, if ever, seen in *Ps. paucimobilis*. Yellow-pigmented strains of *Ps. cepacia*, *Ps. maltophilia*, and *Ps. stutzeri* may also be encountered occasionally, but, as well as all growing on MacConkey agar, they can be further distinguished from *Ps. paucimobilis* by production of caseinase and growth on Simmons' citrate in the case of *Ps. cepacia*, production of caseinase, gelatinase, and decarboxylation of lysine in *Ps. maltophilia*, and by the wrinkled colonies and nitrate reduction (to nitrogen gas) in *Ps. stutzeri*.

The role played by *Ps. paucimobilis* in infections is still uncertain although it may be responsible for various opportunistic infections. However, our patient had been previously healthy, had not been on any antibacterial therapy, and had had no recent contact with hospitals. The association of *Ps. paucimobilis* with moist sites suggests that the species is water-borne, and it is therefore possible that initial entry into the body was through the gastrointestinal tract. Failure to recover the organism from staff involved in the collection and examination of the patient specimens reinforces our belief that

Ps. paucimobilis was the causative agent of meningitis in this patient.

We thank Dr J. Litchfield for permission to report this case and Dr D. Trash for his help.

References

- Holmes, B., Lapage, S. P., and Malnick, H. (1975). Strains of *Pseudomonas putrefaciens* from clinical material. *Journal of Clinical Pathology*, **28**, 149-155.
- Holmes, B., Owen, R. J., Evans, A., Malnick, H., and Willcox, W. R. (1977). *Pseudomonas paucimobilis*, a new species isolated from human clinical specimens, the hospital environment, and other sources. *International Journal of Systematic Bacteriology*, **27**, 133-146.
- Peel, M. M., Davis, J. M., Armstrong, W. L. H., Wilson, J. R., and Holmes, B. (1979). *Pseudomonas paucimobilis* from an ulcer on the leg of a Japanese seaman. *Journal of Clinical Microbiology*, (in press).
- Slotnick, I. J., Hall, J., and Sacks, H. (1979). *Pseudomonas paucimobilis* septicaemia. Report of a case. *American Journal of Clinical Pathology*, (in press).
- Tatum, H. W., Ewing, W. H., and Weaver, R. E. (1974). Miscellaneous Gram-negative bacteria. In *Manual of Clinical Microbiology*, 2nd edition, edited by E. H. Lennette, E. H. Spaulding, and J. P. Truant, pp. 270-294. American Society for Microbiology, Washington, DC.

Requests for reprints to: Dr V. Hajiroussou, Manor Hospital, Moat Road, Walsall WS2 9PS, UK.

Reports and Bulletins prepared by the Association of Clinical Biochemists

The following reports and bulletins are published by the Association of Clinical Biochemists. They may be obtained from The Publishing Department, British Medical Journal (ACB Technical Bulletins), B.M.A. House, Tavistock Square, London WC1H 9JR. Overseas readers should remit by British Postal or Money Order.

SCIENTIFIC REVIEWS (price £1.00/\$2.00 each)

- 1 The assessment of thyroid function March 1971
F. V. FLYNN and J. R. HOBBS
- 2 Renal function tests suitable for clinical practice
January 1972 F. L. MITCHELL, N. VEALL, and R. W. E. WATTS
- 3 Biochemical tests for the assessment of fetoplacental
function May 1975 C. E. WILDE and R. E. OAKEY
- 4 Test of exocrine pancreatic function March 1977
A. H. GOWENLOCK
- 5 Assay of cholinesterase in clinical chemistry March
1979 ELSIE SILK, J. KING, and MARY WHITTAKER

TECHNICAL BULLETINS (price £1.00/\$2.00 each)

- 22 Bilirubin standards and the determination of bilirubin
by manual and technicon AutoAnalyzer methods
January 1971 BARBARA BILLING, RUTH HASLAM, and N.
WALD
- 23 Interchangeable cells for spectrophotometers and
fluorimeters September 1971 S. S. BROWN and A. H.
GOWENLOCK
- 24 Simple tests to detect poisons March 1972 B. W.
MEADE *et al.*
- 25 Blood gas analysers May 1972 K. DIXON
- 26 Kits for enzyme activity determination September
1972 S. B. ROSALKI and D. TARLOW
- 27 Assessment of pumps suitable for incorporation into
existing continuous flow analytical systems November
1972 A. FLECK *et al.*

- 28 Routine clinical measurements of transferrin in
human serum September 1973 K. DIXON
- 29 Control materials for clinical biochemistry (5th
edition) September 1973 J. F. STEVENS
- 30 Notes on the quality of performance of serum
cholesterol assays September 1973 S. S. BROWN
- 31 Determination of uric acid in blood and in urine
July 1974 R. W. E. WATTS
- 32 A survey of amino acid analysers readily available in
the United Kingdom September 1974 J. E. CARLYLE
and P. PURKISS
- 33 Definitions of some words and terms used in auto-
mated analysis November 1974 A. FLECK, R. ROBINSON,
S. S. BROWN, and J. R. HOBBS
- 34 Measurement of albumin in the sera of patients
January 1975 LINDA SLATER, P. M. CARTER, and J. R.
HOBBS
- 35 Investigation of the validity of temperature correction
factors for serum aspartate and alanine transaminases
March 1975 S. B. ROSALKI *et al.*
- 36 Factors influencing the assay of creatinine November
1975 J. G. H. COOK
- 37 A survey of enzyme reaction rate analysers readily
available in the United Kingdom July 1977 R. A.
SAUNDERS and R. F. BURNS
- 38 Transport of specimens for clinical chemistry an-
alysis November 1977 P. WILDING, J. F. ZILVA, and
C. E. WILDE
- 39 A scheme for the evaluation of diagnostic kits May
1978 P. H. LLOYD