this antiphagocytic action in guinea-pigs and rats is due to the large amounts of exopolysaccharide produced by these bacteria. Hence, the ability of mucoid bacteria to escape the normal alveolar macrophage-mediated clearance mechanisms of the lung appears to be due primarily to a 'barrier' effect of the slime. In addition, it has been shown that slime from mucoid type *Ps. aeruginosa* will inhibit phagocytic killing of some bacteria by rabbit polymorphonuclear leucocytes.⁶

If it is possible to inhibit production of the exopolysaccharide or to destroy it after formation, normal clearance mechanisms may be able to operate. An alternative is to stimulate macrophage activity so that phagocytosis is able to take place in the presence of the slime. We think that these findings are important in the study of the mucoid form of *Ps. aeruginosa* in chronic respiratory disease. The results may partly explain the difficulty in clearing mucoid organisms from the lung and the increase in severity of cystic fibrosis associated with the presence of this organism.

While it has long been recognised that capsules can confer virulence on some bacteria, there are no reports of previous studies demonstrating this using alveolar macrophages and the mucoid form of *Ps. aeruginosa*.

This may be explained by the difficulty that exists in working with the mucoid form of this organism, as spontaneous reversion to the non-mucoid form occurs *in vitro*.⁷ Selection of the colony from solid media is important, particularly when non-mucoid forms are present in the culture. Finally, when long-term *in vitro* exposure to macrophages is attempted, spontaneous reversion to non-mucoid forms allows phagocytosis to take place, giving the false impression that mucoid forms have been engulfed.

RW RUHEN¹

PG HOLT²

JM PAPADIMITRIOU³

Departments of ¹Microbiology and

³Pathology,

University of Western Australia
Queen Elizabeth II Medical Centre,
Nedlands 6009, Western Australia

²Princess Margaret Children's Research
Foundation,
Subiaco 6008, Western Australia

References

- ¹ Doggett RG, Harrison GM, Carter RE. Mucoid *Pseudomonas aeruginosa* in

patients with chronic illnesses. *Lancet* 1971;1:236-7.

- ² Burns MW. Significance of *Pseudomonas aeruginosa* in sputum. *Br Med J* 1973;3:283.

- ³ Doggett RG, *et al.* An atypical *Pseudomonas aeruginosa* associated with cystic fibrosis of the pancreas. *J Pediatr* 1966;68:215-21.

- ⁴ Burns MW, May JR. Bacterial precipitins in serum of patients with cystic fibrosis. *Lancet* 1968;1:270-2.

- ⁵ Holt PG. Alveolar macrophages. I. A simple technique for the preparation of high numbers of viable alveolar macrophages from small laboratory animals. *J Immunol Methods* 1979;27:189-98.

- ⁶ Schwarzmans S, Boring JR. Antiphagocytic effect of slime from a mucoid strain of *Pseudomonas aeruginosa*. *Infect Immun* 1971;3:762-7.

- ⁷ Govan JRW. Mucoid strains of *Pseudomonas aeruginosa*: The influence of culture medium on the stability of mucus production. *J Med Microbiol* 1975;8:513-22.

Correlation between two commercial streptococcal grouping kits

In recent years, a number of rapid streptococcal grouping kits have become commercially available. We have evaluated one, the Streptosec test (Organon Teknika), which has recently been marketed, by comparing it with another, the Streptex test (Wellcome Reagents), a method in current use in our laboratory that has been shown to be satisfactory and reliable.¹

Streptosec is a coagglutination method employing antibodies to the streptococci (groups A, B, C, and G) bound to Cowan type I staphylococci. The four reagents are stained and dried on to wells on a white plastic tile.

Both tests were carried out according to the manufacturers' instructions as follows:

Streptex A heavy suspension of beta-haemolytic streptococci was made in 0.4 ml (400 μ l) of extraction enzyme in a test tube. The suspension was incubated at 56°C for 1 hour and then centrifuged at 1200 g for 10 minutes. Using a pasteur pipette, 1 drop of the clear supernatant was added to each of the six circles on the glass tile provided with the kit. One drop of each well-mixed streptococcal latex suspension (groups A, B, C, D, F, and G) was added to the appropriate circle on the tile and mixed with the bacterial extract with a wooden stick. The slide was rocked gently for up to 2 minutes and examined for agglutination.

Streptosec Three to four streaks of growth were taken from a dense culture of beta-haemolytic streptococci on blood agar with a wire loop and suspended in 250 μ l of Todd Hewitt broth. The suspension was mixed for a few seconds in a vortex mixer; 50 μ l of suspension was placed next to the blue spot in each reaction area of the tile. The liquid and dried reagent were mixed using a plastic spatula (provided with the kit) and spread over the reaction area. The plate was rocked gently for 60 seconds and examined for coagglutination. When cross reactions occurred, the suspension was trypsinised and retested.

Eighty-four fresh clinical isolates of beta haemolytic streptococci found to be of group A, B, C, or G by the Streptex method were grouped by the Streptosec method. With Streptex 23 were group A, 29 group B, 8 group C, and 24 group G. The same results were obtained with Streptosec. Thus, in our hands, Streptosec provided a reliable alternative for the grouping of beta haemolytic streptococci belonging to groups A, B, C, and G.

CS AUBREY

M WALLIS

NA SIMMONS

Department of Clinical Bacteriology
and Virology,
Guy's Hospital,
London SE1 9RT.

Reference

- ¹ Facklam RR, Cooksey RC, Wortham EC. Evaluation of commercial latex agglutination reagents for grouping streptococci. *J Clin Microbiol* 1979;10:641-6.

Immunoperoxidase staining

Recently, my MLSOs have been anxious to develop immunoperoxidase methods, and I have discovered that they were ordering the reagent diaminobenzidine tetrahydrochloride, which was described by Mason *et al.* (*J Clin Pathol* 1980;33:609-16). This reagent, of course, has been mentioned in previous papers.

The formula of this reagent suggests that it is related to the carcinogenic hydrocarbons, some of which, particularly benzidine, are prohibited for laboratory use. There is no indication either in the papers or in the catalogues of reagents that these substances are carcinogenic. In the most recent publication, *Safety in Pathology Laboratories*,¹ benzidine is included among the carcinogens, and it is recommended that methods using these chemicals should be discontinued (HM

Letters to the Editors

(69) 57 and 74); it is also stated that when these reagents must be used the Chester Beattie Code of Practice should be rigidly followed.

I suggest that papers reporting studies in which these reagents are used should state that these are carcinogenic hydrocarbons. The chemical firms supplying these materials should indicate in their catalogues that these are carcinogenic hydrocarbons and that the Chester Beattie Code of Practice should be followed in their use.

SYLVIA W DAVIES
Department of Pathology,
St Catherine's Hospital,
Church Road,
Birkenhead,
Merseyside L42 0LQ

Reference

- ¹ *Safety in Pathology Laboratories*, Department of Health and Social Security 1972 Dd 908849 K200 5/72 (8967) HMSO, 1972;44.

Historic laboratory apparatus

The photograph shows a paraffin lamp; the coil-like objects on top are hollow, cone-shaped receptacles of copper which fit closely into one another. This paraffin lamp was found in an old laboratory in Valetta, Malta, where, at the beginning of this century, much of the research work that led to the discovery of the *Micrococcus melitensis* (now called *Brucella melitensis*)

in the blood of the goat took place under the aegis of the Commission for the Investigation of Mediterranean Fever.

Happily, this old laboratory has now been restored (25 June 1980) by the expert and devoted work of the eminent Maltese medical historian, Dr Paul Cassar, who would appreciate very much the views and suggestions of your readers as to what could possibly be the nature and uses of this ancient? DIY item of laboratory equipment.

ETHELWALD E VELLA
Colonel,
Royal Army Medical College,
Millbank,
London SW1P 4RJ

Stability of heparin in intravenous fluids

Administration of heparin by continuous intravenous infusions has been shown to be as effective as intermittent intravenous injection in the treatment of thromboembolism, but major bleeding is a more frequent complication with intermittent injection of heparin than with continuous infusion.^{1,2} Nevertheless, reports that heparin may be unstable in intravenous fluids continue to cast doubt on continuous infusion being the optimal method of administration. Jacobs *et al.*³ and Okuno and Nelson,⁴ using different assays of heparin, reported a loss of its potency in commonly used intravenous fluids, even within a few hours, and measurements with anti-Xa assay apparently showed

erratic behaviour. These observations require confirmation since they carry significant implications regarding heparin administration in the management of venous thromboembolism.

The stability of heparin in intravenous fluids has been re-examined using four assays which measure most known aspects of heparin activity. The activated partial thromboplastin time method, the protamine sulphate titration technique, the metachromatic assay, and an anti-Xa assay were used, and details of these methods have been reported previously.⁵ Heparin (mucous) from Allen and Hanburys (Glaxo Australia) batch No. 251324 (5000 units/ml) was used in these studies; 5 ml of heparin was added to 1 litre glass bottles of saline, 5% dextrose (Abbott Laboratories), and Hartmann's solution (Travenol Laboratories). It was found that the initial volume of fluid in these bottles ranged from 1001 to 1069 ml (± 1 ml). Thus the final concentration of heparin was 24 ± 1 units/ml. The bottles were stored at room temperature ($22 \pm 2^\circ\text{C}$), and samples were removed with a syringe as required.

As can be seen from the Table, no loss was detected in the potency of heparin in intravenous fluids for up to 24 hours by either the chemical or biological assays. Contrary to the suggestion of Okuno and Nelson,⁴ sensitivity of the method of assay was found to have no relation to the stability of heparin. In a separate series of experiments, the pH was monitored continuously with a strip-chart recorder connected to a Radiometer PHM61 pH meter

Potency of heparin in intravenous fluids with time as a percentage of the initial potency

Intravenous fluid	pH after heparin addition	Time after heparin addition (hours)	Anti-Xa assay	Protamine sulphate titration	Metachromatic assay	APTT
Hartmann's solution	5.92	0	100	100	100	100
		3	103	100	104	102
		6	104	100	97	98
		24	104	100	100	98
Saline	5.40	0	100	100	100	100
		3	98	100	96	101
		6	104	100	100	103
		24	101	100	102	102
5% Dextrose	5.95	0	100	100	100	100
		3	94	100	102	101
		6	97	100	99	97
		24	100	100	100	95

Results are the mean of six measurements. The pH of the final mixture did not show a change with time.