- ⁷ Rosen PP, Snyder RE, Robbins G. Specimen radiography for nonpalpable breast lesions found by mammography: procedures and results. *Cancer* 1974;**34**:2028-33.
- ⁸ Fisher ER, Posada M, Ramos H. Evaluation of mammography based upon correlation of specimen mammograms and histopathologic findings. *Am J Clin Pathol* 1974;**62**:60-72.
- ⁹ Gallagher HS. Breast specimen radiography. *Am J Clin Pathol* 1975;**64**:749-55.
- ¹⁰ Koehl RH, Snyder RE, Hutter RVP, Foote FW. The incidence and significance of calcifications within operative breast

specimens. *Am J Clin Pathol* 1970;**53**:3-14.

- ¹¹ Egan JF, Saylor CB, Goodman MJ. A technique for localising occult lesions. *CA*. 1976;**26**:32-7.

Requests for reprints to: Dr GWH Stamp, Department of Pathology, University of Liverpool, Duncan Building, Royal Liverpool Hospital, Prescott Street, Merseyside L69 3BX, England.

Letters to the Editor

Histological criteria for childhood coeliac disease

I am writing to confirm the findings by Dr MN Marsh on high mitotic indices¹ in small intestinal mucosal biopsy of patients with gluten-sensitive enteropathy.

I would like to emphasise that in our series a high mitotic index has been one of the most useful pointers in differentiating coeliac disease from other causes of "flat mucosa," including that of cow's milk intolerance. The three important morphological criteria used by us in childhood coeliac disease are as follows: (a) a flat mucosa with elongated crypts (the latter is absent in most cases of cow's milk intolerance), giving the mucosa an appearance of a normal or occasionally increased thickness (> 550 μ m); (b) infiltration of the surface enterocytes by an increased number of lymphocytes (emperipolesis) and a mitotic index of the lymphocytes of greater than 0.3%; (c) abnormal fat absorption pattern, characterised by accumulation of fat blobs in the supranuclear spaces of the surface enterocytes, which is easily detected by any special fat stain.

F RAAFAT

Queen Elizabeth Hospital for Children,
Hackney Road,
London E2 8PS

Reference

- ¹ Marsh MN. Studies of intestinal lymphoid tissue. IV—The predictive value of raised mitotic indices among jejunal epithelial lymphocytes in the diagnosis of gluten-sensitive enteropathy. *J Clin Pathol* 1982;**35**:517-25.

The in vitro responses of *Bacteroides fragilis* to Moxalactam, Cefotaxime, Cefmetazole, Josamycin and erythromycin

Although the sensitivity of *Bacteroides fragilis* to Moxalactam, Cefotaxime, Cef-

metazole, Josamycin and erythromycin has been described little has been reported about the bactericidal effect of these antibiotics against this organism. Such an effect may be of considerable importance when the treatment of highly susceptible patients—such as the immunosuppressed—is being considered. Furthermore, little is known about the in vitro and in vivo acquisition of resistance of *B fragilis* to these drugs. The results of our observations may, therefore, be of interest.

We have studied the susceptibility of two clinically isolated strains of *B fragilis* ss *fragilis* (strains A and B) and one of *B fragilis* ss *vulgatus* (strain C) to Moxalactam, Cefotaxime, Cefmetazole, Josamycin and erythromycin. Tubes of brucella-enriched broth medium containing serial dilutions of the various antimicrobial drugs were inoculated with approximately 10^7 CFU/ml of the strains under test. Anaerobic incubation at 35°C was carried out over 24 h. Subcultures were made using a standard loop, to blood agar. The minimum inhibitory concentration (MIC) was defined as the minimum antibiotic concentration preventing bacterial growth detectable by the naked eye after 24 h incubation. The minimum bactericidal concentration (MBC) was arbitrarily defined as the minimum concentration of the drug producing a 99.9% reduction of the inoculum.

The MIC and MBC values for Moxalactam, Cefotaxime, Cefmetazole, Josamycin and erythromycin against the three strains

of *Bacteroides* tested are shown in the Table. Cefotaxime showed poor activity against the three strains when compared with the other four drugs. Moxalactam, Cefotaxime and Cefmetazole were bactericidal at concentrations equal to or only slightly higher than their inhibitory values. Although sterilisation of the cultures was not achieved—even with concentrations 30–500 times higher than the particular MIC—the average reduction was from 10^7 CFU/ml to 10^3 CFU/ml. The effect of Josamycin and erythromycin was entirely bacteriostatic failing to reduce the initial inoculum at concentrations 30–100 times higher than their MICs.

Resistance to the antibiotics studied was not induced in the *Bacteroides* strains following a single exposure at concentrations below, equal to, or above the MICs. Resistance following a single exposure, although uncommon, has been shown to occur with clindamycin¹ and in *Bacteroides* isolated from faeces of patients receiving this drug.²

In summary, considerable bactericidal activity against *B fragilis* has been demonstrated by Moxalactam, Cefotaxime and Cefmetazole although each drug produced "persisters"; these are defined as (bacterial) cells which survive exposure to ostensibly lethal concentrations of bactericidal antibiotics but whose progeny remain fully sensitive to the drug.³ *B fragilis* organisms demonstrating such phenotype resistance may play an important part in

Minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC) of five antimicrobials against *Bacteroides fragilis* (mg/l)

	Moxalactam		Cefotaxime		Cefmetazole		Josamycin		erythromycin	
	MIC	MBC	MIC	MBC	MIC	MBC	MIC	MBC	MIC	MBC
B fragilis A	2	4	64	256	4	4	0.25	>32	0.25	>32
B fragilis B	2	2	32	256	4	16	0.5	>32	1	>32
B fragilis C	2	4	32	512	4	4	0.12	>32	0.12	>32