- ⁶ Stamm AM, Polt SS. False negative cryptococcal antigen test. *JAMA* 1980; **244**:1359.
- ⁷ Gordon MA, Lapa E. Elimination of rheumatoid factor in the latex test for cryptococcosis. *Am J Clin Pathol* 1974; **61**:488-94.
- ⁸ Dolan CT. Specificity of the latex-cryptococcal antigen test. *Am J Clin Pathol* 1972; **58**:358-64.
- ⁹ Markenson JA, McDougal JS, Tsairis P, Lockshin MD, Christian CL. Rheumatoid meningitis: a localised immune process. *Ann Intern Med* 1979; **90**:786-9.
- ¹⁰ Bartfield H. The incidence and significance of seropositive tests for rheumatoid factor in non-rheumatoid diseases. *Ann Intern Med* 1960; **52**:1059-66.

Immunocytochemistry of gastric mucosal blood groups

We read with interest the findings of the detailed study by Kapadia *et al*¹ on the immunocytochemistry of gastric mucosal blood groups.

We studied gastric carcinoma and agree that only a minority of these cases show loss of blood group antigens.² We were staining the alcohol-soluble glycolipid blood group substance present on the endothelial cells of all subjects, regardless of secretor status like Kovarik *et al*³ and Davidsohn *et al*⁴ 1971 and suppose that Kapadia *et al* were staining the water-soluble glycoprotein component—no mention is made of small vessel staining. We found changes in some tumours involving loss of A or B substance and persistence of H (loss of terminal residue). The peroxidase technique is more sensitive and we are presumably staining a different antigen but would suggest that non-secretors may express certain blood group antigens in gastric mucosa. The exploration of the hypotheses suggested on page 334¹ will require a comprehensive study of all types of A, B, H antigens.

CL BERRY
Department of Morbid Anatomy,
The London Hospital,
London E1 1BB

References

- ¹ Kapadia A, Feizi T, Jewell D, Keeling J, Slavin G. Immunocytochemical studies of blood group A, H, I and i Antigens in gastric mucosae of infants with normal gastric histology and of patients with gastric carcinoma and chronic benign peptic ulceration *J Clin Pathol* 1981; **34**:320-37.

- ² Slocombe GW, Berry CL, Swettenham KV. The variability of blood group antigens in gastric carcinoma as demonstrated by the immunoperoxidase technique. *Virchows Archiv A* 1980; **387**:289-300.
- ³ Kovarik S, Davidsohn I, Stejskal R. ABO antigens in cancer: detection with the mixed cell agglutination reaction. *Arch Pathol* 1968; **86**:12-21.
- ⁴ Davidsohn I, Ni Ly, Stejskal R. Tissue isoantigens A, B and H in carcinoma of the stomach. *Arch Pathol* 1971; **92**:456-64.

Dr Kapadia and colleagues reply as follows:

Our study (see reference 1 above) was largely concerned with blood group A, H, I and i activities of the mucus secreting epithelial cells of the stomach. The predominant material showing immunofluorescence would be glycoproteins. The conclusions were based on reactivities of the tissue sections with one each of the following antisera: anti-A, anti-H, anti-I(Ma), anti-I(Step) and anti-i(Den). Precise measurement of the amounts of immunoreactive material is not possible in tissue sections and it is likely that the staining reactions reflect the fine specificities and the affinities of antibodies in a given antiserum.

Additional problems in quantitation arise concerning glycolipids carrying the blood group antigens. These are extracted to varying degrees by formalin-fixation and paraffin-embedding procedures. It is possible that the proportion of glycolipids is increased in the tumour cells. Thus the staining reactions with endothelial and tumour cells reflect material that survives the extraction procedures.

T FEIZI
A KAPADIA
G SLAVIN
Clinical Research Centre,
Division of Communicable Diseases,
Watford Road, Harrow,
Middlesex, HA1 3UJ

Prognostic value of measurement of elastosis in breast carcinoma

We are impressed by the correlation obtained by AJ Robertson and colleagues (July 1981)¹ between visual estimation of breast cancer elastosis, and that generated by Quantimet 720 analysis of the same sections of the material.

However, we are concerned that lack of homogeneity of elastosis in breast lesions may be responsible for the apparent absence of prognostic value. The topo-

graphical distribution of elastic fibres in benign^{2,3} and malignant⁴ breast lesions exhibits a focal, discontinuous character. In an earlier study⁵ of breast cancer elastic we found poor correlation between gravimetric assay of insoluble elastin in tumour samples and visual estimation of elastosis in histological preparations from elsewhere in the same carcinomas. Furthermore, even at microscopic level there are at least two types of tumour elastosis. One occurs in the vicinity of endogenous⁶ ducts, blood vessels and interlobular stroma, and the other, around neoplastic⁷ invasive carcinoma cells.

Before concluding that elastosis bears no relation to breast cancer prognosis, we feel that the elements of elastotic lesions should be examined individually and in their topographical context.

JD DAVIES
K BARNARD
Department of Pathology,
University of Bristol,
Bristol BS8 1TD

References

- ¹ Robertson JD, Brown RA, Cree IA, MacGillivray JB, Slidders W, Beck JB. Prognostic value of measurement of elastosis in breast carcinoma. *J Clin Pathol* 1981; **34**:738-843.
- ² Davies JD. Causes of mammary-duct obliteration and hyperelastosis. *Senologia (Paris)* 1976; **1**:84.
- ³ Davies JD, Haines JF. Duct obliteration in sclerosing adenosis. *Proc VI Congr Europ-Soc Pathol* 1977; **6**:177.
- ⁴ Billroth T. Untersuchungen über den feineren Bau und die Entwicklung der Brustdrüsengeschwülste. *Virchows Archiv* 1860; **18**:51-81.
- ⁵ Harris PL, Burn IJ, Barnard K, McCullagh KG, Partridge SM. Elastosis in breast cancer. *Clin Oncol* 1978; **4**:102-3.
- ⁶ Davies JD, Barnard K, Young EW. Peroxidase-antiperoxidase immunolocalisation of newly formed elastin in paraffin-embedded breast lesions. *Proc Path Soc Gt Brit* 1981; **143**:20.
- ⁷ McCullagh KG, Barnard K, Davies JD, Partridge SM. Newly synthesised elastin is associated with neoplastic epithelial cells in human mammary carcinoma. *Experientia* 1980; **36**:1315-6.

Book reviews

Recent Advances in Haematology. No 3. Ed AV Hoffbrand. (Pp 352; illustrated; £16.) Churchill Livingstone. 1982.