Platelet and coagulation factors in proliferative diabetic retinopathy

DQ BORSEY,* CV PROWSE,† RS GRAY,* J DAWES,‡ K JAMES,\$ RA ELTON, ||
BF CLARKE*

*From the *Diabetic and Dietetic Department; †Blood Transfusion Service, Royal Infirmary, Edinburgh; ‡MRC Immunoassay Team, Edinburgh; §Department of Surgery and ||Medical Computing and Statistics Unit, University of Edinburgh, Scotland*

SUMMARY Plasma β -thromboglobulin, platelet factor 4, fibrinogen, fibrinopeptide A, antithrombin III, factor VIII related antigen, α_2 -macroglobulin, platelet count, and total glycosylated haemoglobin were measured in three well matched groups of subjects: non-diabetic controls, diabetics without retinopathy, and diabetics with proliferative retinopathy. β -thromboglobulin and platelet factor 4 concentrations were significantly higher in the diabetics with retinopathy than in the controls and platelet factor 4 was also increased in the diabetics without retinopathy compared with controls. Fibrinogen concentration was raised in diabetics without retinopathy compared with controls, diabetics with retinopathy compared with controls, and diabetics with retinopathy compared with those without. Fibrinopeptide A concentration did not differ significantly between groups. Antithrombin III levels were increased in diabetics with retinopathy compared with controls, and in diabetics with retinopathy compared with those without. Factor VIII related antigen values were higher in both the diabetic groups when compared with the controls. Fibrinopeptide A concentration correlated with both β -thromboglobulin and platelet factor 4 in each of the three groups.

Haemostatic abnormalities in diabetes have been shown, although a hypercoagulable state has not been confirmed. These changes in platelet and coagulation function may be secondary to the development of microvascular disease and their role in the pathogenesis of retinopathy remains uncertain.

Patients with diabetes mellitus have significantly increased morbidity and mortality as a consequence of specific microvascular disease, which results in conditions such as retinopathy and nephropathy. Although the precise cause of these vascular complications remains uncertain, evidence is accumulating that an imbalance of the haemostatic mechanisms may be entailed in their initiation or propagation. Various abnormalities of platelet function and of coagulation in diabetes have been identified and recently reviewed.¹ The contradictory findings and conclusions of different research groups may be partially attributed to the heterogeneity of their diabetic patients, the poor matching of control groups, the non-standardisation of coagulation and platelet assays, and the uncertain importance of abnormal *in vitro* tests.

We have measured plasma concentrations of β -thromboglobulin, platelet factor 4, fibrinogen, fibrinopeptide A, antithrombin III, factor VIII related antigen, α_2 -macroglobulin, platelet count, and total glycosylated haemoglobin in three well matched groups of subjects: non-diabetic controls, insulin dependent diabetics without retinopathy, and insulin dependent diabetics with proliferative retinopathy.

Patients and methods

PATIENTS

We studied 20 healthy non-diabetic controls, 20 insulin dependent diabetics without ophthalmoscopic evidence of retinopathy, and 20 insulin dependent diabetics with proliferative retinopathy. Subjects in each group were individually matched for age, sex and, where applicable, duration of diab-

Table 1 Details of the subjects studied

	Normal subjects	Diabetics	
		without retinopathy	with retinopathy
No of subjects	20	20	20
% male	60	60	60
Age (yr)	35 ± 9	34 ± 9	35 ± 11
Duration of diabetes (yr)	—	21 ± 7	19 ± 7
Daily insulin dose (units)	—	62 ± 18	68 ± 29
% ideal body weight*	—	101 ± 11	100 ± 11

Values given as mean ± 1 SD.

*Metropolitan Life Assurance Company Tables 1960.

etes. Details of the subjects studied are given in Table 1. All diabetics had presented with the clinical features of type I diabetes before the age of 35 and been treated with insulin since diagnosis.

All subjects had normal serum creatinine concentrations and absence of proteinuria on testing with Albustix. None had chronic liver disease or was receiving drugs (for example, oral contraceptives, salicylates) known to interfere with haemostasis. Only two of the diabetics studied had signs or symptoms, or both, of macrovascular disease (one angina and the other peripheral vascular disease) and they both had retinopathy.

METHODS

All diabetic fundi were examined ophthalmoscopically through dilated pupils by one observer (BFC). Proliferative retinopathy was defined as the presence of neovascularisation. A single, 2 h postprandial venous blood sample was withdrawn into a polypropylene syringe, with minimal venous stasis, from resting subjects. The samples were analysed as detailed below:

β -thromboglobulin—radioimmunoassay of samples collected into anticoagulant mixture containing prostaglandin E₁.^{2,3}

Platelet factor 4—radioimmunoassay.⁴

Fibrinopeptide A—radioimmunoassay of samples collected into aprotinin-heparin.⁵

Fibrinogen (factor I)—clot turbidity assay of Ellis and Stransky.⁶

Antithrombin III

(a) **Antithrombin III activity**—bioassay with chromogenic substrate using a commercially available kit (Boehringer-Mannheim Ltd).

(b) **Antithrombin III antigen**—electroimmunoassay method of Laurell using commercial antisera (Behringwerke).⁷

Factor VIII related antigen—electroimmunoassay method of Laurell using commercial antisera (Behringwerke).⁷

α_2 -macroglobulin—standard Mancini gel diffusion method.⁸ Standard antigen in four different concent-

rations was included on every plate. The standard was a human α_2 -macroglobulin reference obtained from Meloy Laboratories Inc (Springfield, Virginia, USA). α_2 -macroglobulin was measured only in the diabetic subjects.

Glycosylated haemoglobin—measured by a modification of the short column chromatographic method of Kynoch and Lehmann.⁹

Platelet count—electronically determined using the Thrombocounter (Coulter Ltd).

The assay methods for β -thromboglobulin, platelet factor 4, fibrinopeptide A, antithrombin III activity, antithrombin III antigen, and factor VIII related antigen gave within batch coefficients of variation between 4 and 6% with between batch coefficients of variation of 8–12%. The between batch coefficient of variation for glycosylated haemoglobin was 3.5%. Fibrinogen levels and platelet count were determined by non-batch methods with coefficients of variation of 5% and 3% respectively. Previous studies by one of us have shown that the assay procedure for α_2 -macroglobulin is highly reproducible ($100 \pm 2.2\%$).⁸ Further details of the above assays are available in the quoted references.

Laboratory staff were unaware of the identities of the samples.

STATISTICAL METHODS

Non-parametric methods were used to analyse the data in view of the presence of skew and outliers in at least some of the measurements. Median and range were used as summary statistics within groups, paired measurements were tested between groups by Wilcoxon's signed rank tests, and associations between measurements within groups by Kendall rank correlation coefficients.

Results

Table 2 details the medians and ranges of the measured platelet and coagulation factors within the three groups and also the results of a comparison of paired measurements between groups.

β -thromboglobulin concentrations were higher ($p < 0.01$) in the diabetics with retinopathy than in the non-diabetic controls, and the platelet factor 4 values were significantly increased in both the diabetic groups when compared with the controls (diabetics with retinopathy $p < 0.01$; diabetics without retinopathy $p < 0.05$). A highly significant correlation between β -thromboglobulin and platelet factor 4 were seen in each of the three groups (controls and diabetics with retinopathy $p < 0.001$; diabetics without retinopathy $p < 0.01$).

The fibrinogen concentrations were higher in the

Table 2 Platelet and coagulation factors (median and range) in the three study groups and a comparison of paired measurements between groups

	Non-diabetic controls	Diabetics		Controls v diabetics without retinopathy	Controls v diabetics with retinopathy	Diabetics with retinopathy v those without
		without retinopathy	with retinopathy			
β -thromboglobulin (ng/ml)	31.5 (19.1-68.5)	44.9 (22.2-96.3)	51.5 (24.1-127.5)	NS	†	NS
Platelet factor 4 (ng/ml)	10.4 (10.0-52.0)	17.5 (8.5-72.6)	19.6 (6.7-63.7)	*	†	NS
Fibrinogen (g/l)	1.70 (1.04-2.93)	2.46 (1.56-3.46)	2.93 (1.85-5.70)	†	‡	*
Fibrinopeptide A (ng/ml)	1.64 (1.60-19.20)	2.82 (2.00-16.60)	2.80 (2.00-17.80)	NS	NS	NS
Antithrombin III activity (%)	103 (82-119)	104 (92-122)	116 (92-132)	NS	‡	†
Antithrombin III antigen (%)	100 (76-126)	100 (87-121)	113 (87-126)	NS	*	*
Factor VIII related antigen (%)	98 (72-140)	252 (126-460)	267 (142-400)	‡	‡	NS
α_2 -macroglobulin (mg%)	—	378 (192-641)	480 (187-696)	—	—	NS
Platelet count ($\times 10^9/l$)	226 (106-284)	255 (104-420)	267 (126-350)	*	NS	NS
Glycosylated haemoglobin (%)	6.6 (5.6-8.3)	10.7 (8.3-14.1)	11.7 (10.2-15.9)	‡	‡	*

NS = not significant.

* $p < 0.05$, † $p < 0.01$, ‡ $p < 0.001$.

two diabetic groups than in the controls and also in the diabetics with retinopathy than in those without retinopathy.

There was no significant difference in fibrinopeptide A concentrations between the three groups. Within each group, however, fibrinopeptide A cor-

related significantly with β -thromboglobulin ($p < 0.05$) and platelet factor 4 (controls and diabetics without retinopathy $p < 0.05$; diabetics with retinopathy $p < 0.01$) (Figure).

Antithrombin III (activity and antigen) levels were significantly higher in the diabetics with retinopathy than in either of the other groups. There was a strong correlation between antithrombin III activity and antigen in each of the three groups (controls $p < 0.01$; diabetic groups $p < 0.001$). The concentration of α_2 -macroglobulin was higher in the diabetics with retinopathy, than in those without but this difference was not significant. α_2 -macroglobulin correlated with antithrombin III antigen only in the diabetics with retinopathy. There was no correlation between α_2 -macroglobulin and antithrombin activity.

Concentrations of factor VIII related antigen were higher in both the diabetic groups than in the non-diabetic controls ($p < 0.001$).

The platelet count was higher in the non-retinopathic diabetics than in the control subjects ($p < 0.05$), but no other significant differences existed between groups.

Total glycosylated haemoglobin was higher in the diabetics with retinopathy than in those without ($p < 0.05$). The relation between glycosylated haemoglobin and the haemostatic factors studied was considered in each of the three groups: the only significant correlations were with fibrinogen in the control group ($p < 0.001$), with platelet factor 4 in the uncomplicated diabetics ($p < 0.05$), and with fibrinopeptide A and α_2 -macroglobulin in the diabetics with retinopathy ($p < 0.05$).

Correlation of platelet factor 4 with fibrinopeptide A. Normals: $r = 0.45$, $p < 0.05$; no diabetic retinopathy (No DR): $r = 0.36$, $p < 0.05$; diabetic retinopathy (DR): $r = 0.44$, $p < 0.05$.

Discussion

The published work provides a confusing picture of the importance of haemostatic abnormalities in the development of diabetic retinopathy. The poor matching of study groups, the non-standardisation of assays, the uncertain relevance of *in vitro* tests to *in vivo* function, and the inclusion of patients with coincident nephropathy and macroangiopathy have all contributed to this confusion. The subjects in our groups were individually matched for age, sex, and, where applicable, duration of diabetes, and all diabetics with nephropathy were excluded. In contrast to many previous studies, our two diabetic groups were chosen to represent the extremes of the "retinopathy spectrum"—namely, those without retinopathy and those with proliferative retinopathy. Unfortunately, the exclusion of patients with asymptomatic large vessel disease remains a problem since it necessitates unacceptable extensive and invasive investigation.

The platelet specific proteins β -thromboglobulin and platelet factor 4 are released together from α granules during platelet aggregation, and increased plasma levels may reflect enhanced *in vivo* platelet activation. Their precise biological functions remain uncertain. β -thromboglobulin may reduce prostacyclin generation in the vessel wall¹⁰ and thus allow a more sustained platelet-endothelial interaction. Platelet factor 4 has heparin neutralising activity¹¹ and may neutralise any natural heparin like inhibitors of coagulation. It has been suggested that intensive release of platelet factor 4 modifies the surface properties of the platelet membrane and thereby accelerates the rate of formation of the fibrin network between platelets.¹² A close correlation between the plasma levels of platelet factor 4 and β -thromboglobulin has been well described.^{4 13–15}

Both our diabetic groups had higher β -thromboglobulin concentrations than the controls, although this was significant only in patients with retinopathy. Both normal^{16 17} and increased^{2 15 18–20} plasma β -thromboglobulin concentrations have been reported in diabetes. The lack of a significant difference in β -thromboglobulin concentrations between the diabetics with retinopathy and those without is in keeping with a previous report from Edinburgh,² but contrasts with work from other centres.^{15 18–21} The inconsistent findings of previous studies may be partially attributed to the use of different anticoagulant mixtures, sampling techniques, and assay methods² and to the inclusion of diabetics with nephropathy. The latter is important since plasma β -thromboglobulin (but not platelet factor 4) concentrations are increased in the

Borsey, Prowse, Gray, Dawes, James, Elton, Clarke

presence of renal impairment.^{14 20}

The platelet factor 4 results were, as expected, similar to those for β -thromboglobulin, although both diabetic groups had significantly higher levels than the non-diabetics. Platelet factor 4 concentration has been previously reported as both normal^{17 22} and increased in diabetics.^{15 23}

Fibrinopeptide A, which is cleaved from fibrinogen by thrombin, is a sensitive indicator of thrombin formation²⁴ and the lack of a significant difference in concentrations between our three groups is evidence against activation of the coagulation system in diabetes. Our finding that fibrinopeptide A concentrations correlated with both β -thromboglobulin and platelet factor 4 concentrations suggests that thrombin may initiate the platelet release reaction. A close association between thrombin formation and the platelet release reaction in normal subjects has been previously noted *in vitro*²⁵ and a correlation between fibrinopeptide A and β -thromboglobulin in diabetics has been reported by Burrows.²¹

Fibrinogen is the immediate precursor of fibrin, influences plasma viscosity,²⁶ and is a necessary cofactor for ADP and epinephrine induced platelet aggregation.²⁷ It is generally agreed that fibrinogen concentrations are increased in diabetic patients when compared with controls^{28–30} and several authors have, as in our study, shown higher concentrations in diabetics with retinopathy than in those without.^{29–31} This increase in fibrinogen concentration may be non-specific, perhaps in response to chronic vascular injury, rather than reflecting a hypercoagulable state.¹ The higher concentrations of other acute phase reactants (such as haptoglobin, β -lipoprotein, and caeruloplasmin) in the serum of diabetic subjects³² lends support to this view. Our study has failed to confirm a reported correlation between glycosylated haemoglobin and fibrinogen in diabetics.³³

Antithrombin III is the major inhibitor of the activated coagulant enzymes XIIa, XIa, IXa, Xa, and thrombin. The risk of thrombo-embolic disease in patients with congenital antithrombin III deficiency is significantly increased.³⁴ In a hypercoagulable state antithrombin III would be consumed and low circulating levels expected. Our findings of raised antithrombin III levels in those diabetics with retinopathy was therefore somewhat surprising. There have been reports of low,^{35–37} normal,^{38 39} and raised^{22 29 40} levels of antithrombin III in diabetes. Discrepancies between functional and immunological assays may account for some of these inconsistencies, although we found a strong correlation between antithrombin III activity and antigen. What then is the explanation of our findings? The raised

levels of antithrombin III may simply represent part of an acute phase reaction occurring subsequent to the development of retinopathy. Alternatively, an increase in antithrombin III may occur as a defence response to enhanced intravascular production of thrombin via a feedback mechanism increasing synthesis or decreasing catabolism of antithrombin III.⁴¹

α_2 -macroglobulin is a further inhibitor of coagulation proteases which accounts for about a quarter of the total functional antithrombin III activity in the blood. Indeed, Elder *et al*⁴⁰ attributed the raised levels of antithrombin III activity in their diabetics to an increase in α_2 -macroglobulin concentrations. We found no correlation between antithrombin III activity and α_2 -macroglobulin. Although it is generally accepted that α_2 -macroglobulin levels are higher in diabetics,^{31 42 43} the relation between retinopathy and α_2 -macroglobulin has remained uncertain because of other variables including age, duration of diabetes, metabolic control and nephropathy, which may influence α_2 -macroglobulin levels.⁴²⁻⁴⁴ Previous studies, which often failed to consider these factors, have reported both higher α_2 -macroglobulin concentrations in diabetics with retinopathy^{31 43} and comparable levels in retinopathies and non-retinopathies.⁴² Our results are in keeping with those of Gray *et al*,⁴⁵ who found no significant difference in α_2 -macroglobulin levels between diabetics with retinopathy and those without when all the above variables were taken into account.

Factor VIII related antigen is a glycoprotein, higher oligomers of which are active in promoting platelet adhesion. It is found only in endothelial cells (where it is synthesised), platelets, and megakaryocytes.⁴⁶ Von Willebrand's factor appears to be a property of factor VIII related antigen⁴⁷ and these factors are associated in plasma with factor VIII coagulant activity. Our finding of increased factor VIII related antigen levels in diabetic patients is in keeping with previous reports^{33 48 49} and could result from enzymic degradation in the circulation of the factor VIII polymer and/or from greater endothelial synthesis and release of factor VIII related antigen.^{50 51} The lack of raised plasma fibrinogen A levels supports the latter explanation since in an enzymatically active environment one would expect increased thrombin formation. We found no significant difference in factor VIII related antigen values between diabetics with retinopathy and those without, although higher levels have been previously reported in diabetics with vascular disease.^{48 50 52}

This study has shown certain platelet and coagulation abnormalities in diabetes, but the presence of a hypercoagulable state has not been confirmed. Sev-

eral of our findings may be explained on the basis of a non-specific stress reaction in diabetics, possibly secondary to chronic vascular damage. Further studies, particularly of a prospective nature, are clearly indicated to determine whether the observed changes are causal or secondary to the vascular complications of diabetes mellitus.

References

- Jones RL, Peterson CM. Haematologic alterations in diabetes mellitus. *Am J Med* 1981;70:339-52.
- Borsey DQ, Dawes J, Fraser DM, Prowse CV, Elton RA, Clarke BF. Plasma beta-thromboglobulin in diabetes mellitus. *Diabetologia* 1980;18:353-7.
- Bolton AE, Ludlam CA, Moore S, Pepper DS, Cash JD. Three approaches to the radioimmunoassay of human β -thromboglobulin. *Br J Haematol* 1976;33:233-8.
- Bolton AE, Ludlam CA, Pepper DS, Moore S, Cash JD. A radioimmunoassay for platelet factor 4. *Thromb Res* 1976;8:51-8.
- Kockum C. Radioimmunoassay of fibrinogen A: Clinical applications. *Thromb Res* 1976;8:225-36.
- Ellis BC, Stransky A. A quick and accurate method for the determination of fibrinogen in plasma. *J Lab Clin Med* 1961;58:477-88.
- Laurell CB. Quantitative estimation of proteins by electrophoresis in agarose gel containing antibodies. *Anal Biochem* 1966;15:45-52.
- James K, Johnson G, Fudenberg HH. The quantitative estimation of α_2 -macroglobulin in normal, pathological and cord sera. *Clin Chim Acta* 1966;14:207-14.
- Borsey DQ, Fraser DM, Gray RS, Elton RA, Smith AF, Clarke BF. Glycosylated haemoglobin and its temporal relationship to plasma glucose in non-insulin dependent (Type 2) diabetes mellitus. *Metabolism* 1982;31:362-6.
- Hope W, Chesterman CN, Dusting GJ, Smith I, Morgan FJ, Martin TJ. Inhibition by β -thromboglobulin of prostacyclin (PGI₂) formation in arterial endothelial cells. *Thromb Haemost* 1979;42:8.
- Van Creveld S, Paulsen MMP. Significance of clotting factors in blood-platelets in normal and pathological conditions. *Lancet* 1951;ii:242-4.
- Niewiarowski S, Lipinski B, Farbiszewski R, Poplowski A. The release of platelet factor 4 during platelet aggregation and the possible significance of this reaction in haemostasis. *Experientia* 1968;24:343-4.
- Dawes J, Smith RC, Pepper DS. The release, distribution and clearance of human B-thromboglobulin and platelet factor 4. *Thromb Res* 1978;12:851-61.
- Kurihara Y, Nakayama H, Nakagawa S. Plasma β -thromboglobulin and platelet factor 4 in renal insufficiency. *Thromb Res* 1980;18:557-60.
- Betteridge DJ, Zahavi J, Jones NAG, Shine B, Kakkar VV, Galton DJ. Platelet function in diabetes mellitus in relationship to complications, glycosylated haemoglobin and serum lipoproteins. *Eur J Clin Invest* 1981;11:273-7.
- Campbell IW, Dawes J, Fraser DM, *et al*. Plasma β -thromboglobulin in diabetes mellitus. *Diabetologia* 1977;26:1175-7.
- Kazmier FJ, Bowie EJ, O'Fallon WM, Zimmerman BR, Osmundson PJ, Palumbo PJ. A prospective study of peripheral occlusive arterial disease in diabetes IV: Platelet and plasma functions. *Mayo Clin Proc* 1981;56:243-53.
- Preston FE, Ward JD, Marcola BH, Porter NR, Timperley WR, O'Malley BC. Elevated β -thromboglobulin levels and circula-

- ing platelet aggregates in diabetic microangiopathy. *Lancet* 1978;1:238-9.
- ¹⁹ Burrows AW, Chavin SI, Hockaday TDR. Plasma-thromboglobulin concentrations in diabetes mellitus. *Lancet* 1978;1:235-7.
 - ²⁰ Rasi V, Ikkala E, Hekali R, Myllylä G. Factors affecting plasma β -thromboglobulin in diabetes mellitus. *Med Biol* 1980;58:269-72.
 - ²¹ Burrows AW. Beta-thromboglobulin in diabetes: relationships with blood glucose and fibrinopeptide A. *Horm Metab Res* 1981; suppl II: 22-5.
 - ²² Corbella E, Miragliotta G, Masperi R, Villa S, Bini A, de Gaetano G, Chiumello G. Platelet aggregation and antithrombin III levels in diabetic children. *Haemostasis* 1979;8:30-7.
 - ²³ Chimielewski J, Farbiszewski R. Platelet factor 4 (PF4) release during human platelet aggregation in diabetic patients. *Thromb Diath Haemorrh* 1970;24:203-5.
 - ²⁴ Nossel HL, Ti M, Kaplan KL, Spanondis K, Soland T, Butler VP. The generation of fibrinopeptide A in clinical blood samples: evidence for thrombin activity. *J Clin Invest* 1976;58:1136-44.
 - ²⁵ Prowse CV, Vigano S, Borse DQ, Dawes J. The release of beta-thromboglobulin from platelets during the clotting of whole blood. *Thromb Res* 1980;17:433-42.
 - ²⁶ Hoare EM, Barnes AJ, Dormandy JA. Abnormal blood viscosity in diabetes mellitus and retinopathy. *Biorheology* 1976;13:21-5.
 - ²⁷ Solum NO, Stormorken H. Influence of fibrinogen on the aggregation of washed human blood platelets induced by adenosine diphosphate, thrombin, collagen and adrenaline. *Scand J Clin Lab Invest* 1965;17 (suppl 84): 170-82.
 - ²⁸ Mayne JM, Bridges JM, Weaver JA. Platelet, plasma fibrinogen and factor VIII levels in diabetes mellitus. *Diabetologia* 1970;6:436-40.
 - ²⁹ Fuller H, Keen JH, Jarrett RJ, *et al.* Haemostatic variables associated with diabetes and its complications. *Br Med J* 1978;ii:964-6.
 - ³⁰ Cederholm-Williams SA, Dornan TL, Turner RC. The metabolism of fibrinogen and plasminogen related to diabetic retinopathy in man. *Eur J Clin Invest* 1981;11:133-8.
 - ³¹ Almer LA, Pandolfi M. Fibrinolysis and diabetic retinopathy. *Diabetes* 1976;25 (suppl 2): 807-9.
 - ³² Jonsson A, Wales ZK. Blood glycoprotein levels in diabetes mellitus. *Diabetologia* 1976;12:245-50.
 - ³³ Coller BS, Frank RN, Milton RC, Gralnick HR. Plasma co-factors of platelet function: correlation with diabetic retinopathy and haemoglobins Ala-c. *Ann Intern Med* 1978;88:311-6.
 - ³⁴ Marciniak E, Farley CH, DeSimone PA. Familial thrombosis due to antithrombin III deficiency. *Blood* 1974;43:219-31.
 - ³⁵ Banerjee RN, Sahni AL, Kumere U, Arza N. Antithrombin III deficiency in maturity onset diabetes mellitus and atherosclerosis. *Thromb Diath Haemorrh* 1974;31:339-45.
 - ³⁶ Monnier L, Follae G, Mirouze J. Antithrombin III deficiency in diabetes mellitus: influence on vascular degenerative complications. *Horm Metab Res* 1978;10:470-3.
 - ³⁷ Sowers JR, Tuck ML, Sowers DK. Plasma antithrombin III and thrombin generation time: correlation with haemoglobin A_{1c} and fasting serum glucose in young diabetic women. *Diabetes Care* 1980;3:655-8.
 - ³⁸ Zucker ML, Gomperts ED, Russel D, *et al.* Antithrombin functional activity after saturated and unsaturated fatty meals and fasting in normal subjects and some disease states. *Thromb Res* 1979;15:37-48.
 - ³⁹ Gandolfo GM, DeAngelis A, Torresi MV. Determination of antithrombin III activities by different methods in diabetic patients. *Haemostasis* 1980;9:15-9.
 - ⁴⁰ Elder GE, Mayne EE, Daly JG, Kennedy AL, *et al.* Antithrombin III activity and other coagulation changes in proliferative diabetic retinopathy. *Haemostasis* 1980;9:288-96.
 - ⁴¹ Calvalho A, Ellman L. Activation of the coagulation system in polycythemia vera. *Blood* 1976;47:669-78.
 - ⁴² Ganrot PO, Gydel K, Ekelund H. Serum concentration of alpha₂-macroglobulin, haptoglobin and alpha₁-antitrypsin in diabetes mellitus. *Acta Endocrinol* 1967;55:537-44.
 - ⁴³ James K, Merriman J, Gray RS, Duncan LJP, Herd R. Serum α_2 -macroglobulin levels in diabetes. *J Clin Pathol* 1980;33:163-6.
 - ⁴⁴ Ganrot PO, Schersten B. Serum α_2 -macroglobulin concentration and its variation with age and sex. *Clin Chim Acta* 1967;15:113-20.
 - ⁴⁵ Gray RS, James K, Merriman J *et al.* Alpha₂-macroglobulin and proliferative retinopathy in Types I Diabetes. *Horm Metab Res* 1982;14:389-92.
 - ⁴⁶ Jaffe EA, Deykin D, Salzman E, Schnippe L. Endothelial cells and biology of Factor VIII. *N Engl J Med* 1977;296:377-83.
 - ⁴⁷ Von Willebrand's disease (Editorial). *Br Med J* 1976;ii:715-6.
 - ⁴⁸ Pandolfi M, Almer LO, Holmberg L. Increased von Willebrand-antihaemophilic factor A in diabetic retinopathy. *Acta Ophthalmol* 1974;52:823-8.
 - ⁴⁹ Gensini GF, Abbate R, Favilla S, Neri Serneri GG. Changes of platelet function and blood clotting in diabetes mellitus. *Thromb Haemost* 1979;42:983-93.
 - ⁵⁰ Boneu B, Abbal M, Plante J, Bierme R. Factor VIII complex and endothelium damage. *Lancet* 1975;i:1430.
 - ⁵¹ Giustolisi R, Musso R, Russo M, Cantania N, Lombardo T, Cacciola E. Possible evidence for increased Factor VIII antigen synthesis in vascular endothelium of diabetic subjects. *Thromb Haemost* 1982;47:293.
 - ⁵² Lufkin EG, Fass DN, O'Fallon WM, Bowie EJW. Increased von Willebrand factor in diabetes mellitus. *Metabolism* 1979;28:63-6.

Requests for reprints to: Dr DQ Borse, Ninewells Hospital, Ninewells, Dundee DD1 9SY, Scotland.