Letters

- ² Newman GR, Jasani B, Williams ED. The visualisation of trace amounts of diaminobenzidine (DAB) polymer by a novel gold-sulphide-silver method. *J Microsc* 1983;132:RP1-2.
- ³ Newman GR, Jasani B, Williams ED. Amplification of trace amounts of DAB in immunocytochemistry. *J Pathol* 1984;142:A6-7.
- ⁴ Siegesmund VA, Yorke DE, Drogen R. A quantitative immunoperoxidase procedure employing energy dispersive x ray analysis. *J Histochem Cytochem* 1979;27:1226-30.
- ⁵ Litwin JA. Transition metal-catalysed oxidation of 3,3'-diaminobenzidine (DAB) in a model system. *Acta Histochem* 1982;71:111-7.
- ⁶ Hsu SM, Soban E. Color modification of diaminobenzidine (DAB) precipitation by metallic ions and its application for double immunohistochemistry. *J Histochem Cytochem* 1982;30:1079-82.
- ⁷ Timm F. Zur Histochemie der Schwermetalle: Das Sulfid-silberverfahren. *Zeitschrift Fur Die Gesamte Gerichtliche Medizin (Berlin)* 1958;46:706-11.
- ⁸ Danscher G. Histochemical demonstration of heavy metals. *Histochemistry* 1981;71:1-16.
- ⁹ Gallyas F, Gorcs T, Merchenthaler I. High-grade intensification of the end-product of the diaminobenzidine reaction for peroxidase histochemistry. *J Histochem Cytochem* 1982;30:183-4.
- ¹⁰ Holgate CS, Jackson P, Cowen PN, Bird CC. Immunogold-silver staining: new method of immunostaining with enhanced sensitivity. *J Histochem Cytochem* 1983;31:938-44.
- ¹¹ Chan VTW, Fleming KA, McGee JO'D. Detection of subpicogram quantities of specific DNA sequences on blot hybridization with biotinylated probes. *Nucleic Acids Res* 1985;13:8083-91.

Distinguishing lymphoma and small cell anaplastic carcinoma of the thyroid by immunocytochemistry

I read with interest the report by Burt *et al*¹ regarding the problem of differential diagnosis between lymphoma and small cell anaplastic carcinoma of the thyroid. A similar study has been performed in our department in Leicester on a smaller number of cases (19). In addition to thyroglobulin and epithelial membrane antigen antisera, our study also included a cytokeratin antibody (CAM 5.2). It was found that this antibody was more sensitive for detecting epithelial malignancies than epithelial membrane antigen. A combination of common leucocyte antigen and CAM 5.2 resolved the differential diagnosis in all but two of the cases. The use of this or a related cytokeratin antibody, together with common leucocyte antigen is thus suggested for this diagnostic problem. Epithelial membrane antigen, apart from re-

duced sensitivity compared with cytokeratin, suffers from the additional disadvantage of being reactive in a proportion of lymphomas.²

CH KENDALL
Department of Pathology,
Clinical Sciences Building,
Leicester Royal Infirmary,
PO Box 65,
Leicester LE2 7LX

References

- ¹ Burt AD, Kerr DJ, Brown IL, Boyle P. Lymphoid and epithelial markers in small cell anaplastic thyroid tumours. *J Clin Pathol* 1985;38:893-6.
- ² Gatter KC, Alcock C, Heryet A, Mason DY. Clinical importance of analysing malignant tumours of uncertain origin with immunohistological techniques. *Lancet* 1985;i:1302-5.

T lymphocyte numbers and serum E rosette inhibitory substance

It is well established that T lymphocyte numbers are diminished in certain disease conditions. These include protein calorie malnutrition^{1,2} plasmodium falciparum infection,^{3,4} measles infection,^{5,6} and systemic lupus erythematosus.⁷ Others are HBs Ag positive chronic hepatitis,⁸ rheumatoid arthritis,⁹ and cancer.¹⁰ Sera from most of these patients inhibit both in vitro phytohaemagglutinin transformation of lymphocytes,^{5,11} and E rosette formation by normal human lymphocytes.¹²⁻¹⁷

There is circumstantial evidence to suggest that the presence of E rosette inhibitory substance (probably an immune complex) could be partly responsible for the diminished number of E rosettes that are recorded in these patients. Treatment with levamisole considerably increases the number of E rosette forming T lymphocytes in vivo and in vitro in several disease conditions, in which patients commonly possess circulating immune complexes and low E rosetting lymphocytes.^{10,18-20} We also observed recently that, in common with children who had protein calorie malnutrition, children with malaria or measles infections had increased titres of circulating immune complexes, serum E rosette inhibitory substance(s), and diminished numbers of circulating E rosettes.¹⁶

It is therefore necessary for workers carrying out studies of lymphocyte subpopulations to be cautious in interpreting findings of diminished E rosettes. The percentage of E rosettes observed in such cases

may not represent the total circulating E rosette numbers present. From our experience a test for the presence of E rosette inhibitory substance(s) in such patients is useful, as there may be some circulating T lymphocytes that are not capable of forming E rosettes in vitro, probably as a result of the previous binding of their surface receptors to inhibitory substances in vivo.¹⁴

References

- ¹ Chandra RK. Lymphocyte subpopulations in human malnutrition: cytotoxic and suppressor cells. *Paediatrics* 1977;59:423-7.
- ² Salimonu LS, Johnson AOK, Williams AIO, Adeleye Iyabode G, Osunkoya BO. Lymphocyte subpopulations and antibody levels in immunized malnourished children. *Br J Nutr* 1982;48:7-14.
- ³ Wyler DJ. Peripheral lymphocyte subpopulations in human falciparum malaria. *Clin Exp Immunol* 1976;23:471-6.
- ⁴ Ade-Serrano NA, Osunkoya BO. Circulating T and B lymphocytes in acute falciparum malaria. *Niger Med J* 1977;2:251-4.
- ⁵ Whittle HC, Dosseter J, Oduloju A, Bryceson ADM, Greenwood BM. Cell mediated immunity during natural measles infection. *J Clin Invest* 1978;62:678-84.
- ⁶ Bhaskaram P, Rays Reddy V. Effect of measles on cell mediated immunity. *Indian J Med Res* 1983;77:83-6.
- ⁷ Hurd ER, Giuliano VJ. The effect of cytophosphamide on B and T lymphocytes in patients with connective tissue diseases. *Arthritis Rheum* 1975;18:67-75.
- ⁸ De Horatius RJ, Strickland RG, Williams RC. T and B lymphocytes in acute and chronic hepatitis. *Clin Immunol Immunopathol* 1974;2:353-60.
- ⁹ Yu DTY, Pette JB. Cellular immunological aspects of rheumatoid arthritis. *Semin Arthritis Rheum* 1974;4:25-52.
- ¹⁰ Akaza J, Yokoyama H, Umeda T, Nijima T. Restoration by levamisole of E rosette formation and its abrogation by autochthonous serum from patients with bladder cancer. *Cancer* 1979;1:97-100.
- ¹¹ Moore DL, Heyworth B, Brown J. Effect of autologous plasma on lymphocytes transformation in malaria and in protein-energy malnutrition. *Immunol* 1977;33:777-85.
- ¹² Salimonu LS. Immune responses and a serum inhibitory substance in malnourished children. Nigeria: University of Ibadan, 1980 pp 240. (PhD thesis.)
- ¹³ Salimonu LS, Osunkoya BO. Lymphocytes in protein calorie malnutrition. *Am J Clin Nutr* 1980;33:2699-700.
- ¹⁴ Salimonu LS, Johnson AOK, Williams AIO, Adeleye Iyabode G, Osunkoya BO. The occurrence and properties of E rosette inhibitory substance in the sera of malnourished children. *Clin Exp Immunol* 1982;47:626-34.
- ¹⁵ Salimonu LS. Soluble immune complexes, acute phase proteins and E rosette inhibitory sub-