Role of clostridial organisms in neutropenic enterocolitis

Neutropenic enterocolitis affects the ileocaecal region of the bowel and is characterised by intense thickening and oedema with mucosal necrosis. It is associated with agranulocytosis, often with *Clostridium septicum* infection, and commonly occurs in patients with acute leukaemia, many of whom have been treated with cytosine arabinoside. The strong association with *C septicum* has led to reports that this organism may have a primary causative role in the condition.^{1,2} We report two cases with other clostridial organisms which, we propose, may act as opportunistic invaders.

Case reports**CASE 1**

A 57 year old woman presented with acute myeloid leukaemia. Remission was achieved, followed by relapse and a second remission 10 months later. She again relapsed and received chemotherapy comprising rubidomycin, cytosine arabinoside, and 6-thioguanine. After this she developed intestinal obstruction, with a temperature of 37.5°C and a white cell count of $10.4 \times 10^6/l$, comprising 98% blasts and 2% lymphocytes, with complete agranulocytosis. She died shortly afterwards.

Postmortem examination, performed four hours after death, showed multiple gaseous spaces within the liver and spleen, with haemolytic staining of the inferior vena cava. There was obstructive thickening of the ileum 50 cm proximal to the ileocaecal valve, occupying a 20 cm segment of bowel. Blood cultures before and after death produced pure growths of *C welchii*, as did swabs from the affected area. Histologically there was intense transmural oedema and congestion, with patchy mucosal necrosis, but no evidence of leukaemic infiltration. Gram positive bacilli were numerous and immunofluorescence staining was negative for *C septicum* but positive for *C welchii*.

CASE 2

A 48 year old man treated 10 years previously with cyclophosphamide for a lymphocytic (T cell) lymphoma of the scalp presented with acute myelomonocytic leukaemia. Remission was obtained but he relapsed 13 months after diagnosis. Reinduction was attempted with intermediate dose cytosine arabinoside (0.5 g/m² twice daily). Following this he was asymptomatic with a white cell count of $0.1 \times 10^6/l$. He died unexpectedly with a mild fever but no localising evidence of infection.

Postmortem examination showed evidence of clostridial infection, together with pronounced thickening along a 10 cm segment of ascending colon, which was adherent to the inferior surface of the liver. Multiple mucosal ulcers were noted. Culture of swabs from colon, liver, and spleen produced a heavy growth of *C sordellii*. Histological examination showed mucosal ulceration with gross transmural oedema and congestion of the affected colon. Gram positive bacilli were clustered around the ulcerated areas and on the serosal surface. There was no leukaemic infiltrate.

Clostridial septicemia in patients with malignancy, particularly leukaemia, is well recognised. Wynne and Armstrong³ reported 15 episodes, 11 of which were due to *C welchii*. In all instances the portal of entry was the gastrointestinal tract, due to varying causes including recent anti-neoplastic chemotherapy or radiation, an invasive diagnostic procedure, or malignant disease of the bowel.

C sordellii is rarely regarded as a pathogen, and neither it or *C welchii* have been reported as the sole organism in association with neutropenic enterocolitis. We feel these cases strongly imply that this condition may be complicated by clostridial species other than *C septicum* and that they are behaving opportunistically in a patient with immune deficiency and a mucosal defect of the gastrointestinal tract, rather than that they have a primary causative role. The cause of the initial insult remains unclear, although neutropenic ulceration⁴ and damage by cytosine arabinoside⁵ are contenders, where there is no malignant infiltration of the bowel. Other possibilities are intramural haemorrhage due to thrombocytopenia and localised ischaemic damage.

Clinically, it is important to consider neutropenic enterocolitis in the appropriate clinical setting, and if suspected, antibacterial treatment to cover clostridial sepsis would seem advisable in view of the poor prognosis associated with the condition.

KM NEWBOLD*

MG LORD*

TP BAGLIN†

*Department of Pathology,

The Medical School,

University of Birmingham, B15 2TJ,

†Department of Haematology,

Queen Elizabeth Hospital,

Edgbaston,

Birmingham B15 2TH.

References

- 1 King A, Rampling A, Wight DGD, Warren DE. Neutropenic enterocolitis due to Clostridium

septicum infection. *J Clin Pathol* 1984;37:335-43.

- 2 Rifkin GD. Neutropenic enterocolitis and Clostridium septicum infection in patients with agranulocytosis. *Arch Intern Med* 1980;140:834-5.
- 3 Wynne JW, Armstrong D. Clostridial septicemia. *Cancer* 1972;29:215-21.
- 4 Geelhoed GW, Kane MA, Dale DC, Wells SA. Colon ulceration and perforation in cyclic neutropenia. *J Paediatr Surg* 1973;8:379-82.
- 5 Slavin RE, Dias MA, Saral R. Cytosine arabinoside induced gastrointestinal toxic alterations in sequential chemotherapeutic protocols. *Cancer* 1978;42:1747-59.

Serum alkaline phosphatase isoenzymes

The novel technique for quantitation of serum alkaline phosphatase (ALP) isoenzymes¹ is interesting and clearly permits identification of the source of the serum enzyme. Although we agree that knowledge of the alkaline phosphatase isoenzyme type provides clinically useful information, the clinical evaluation does not really provide sufficient evidence to justify this claim. It is likely that several other tests would be performed prior to the request for quantitative ALP isoenzyme determination; therefore it is essential to evaluate the additional contribution that this test would make to diagnosis, not its contribution in isolation. The authors state that several other biochemical variables were available for each patient but apparently fail to take these into account in their analysis.

We examined serum total ALP activity prospectively in 2884 acute medical admissions and found it to be increased (>1.15 times upper reference limit) in 204 patients (7.1%). Diagnoses were established by means of predetermined clinical, radiological, and histopathological criteria and patients were allocated to one of six diagnostic categories (table 1). Qualitative assessment of ALP isoenzyme type by polyacrylamide gel electrophoresis (PAGE) was available for 178 patients and was consistent with the clinical diagnosis in 70% (80% of those for whom the diagnosis was definitely established).

One hundred and fifty seven had a definite diagnosis and a PAGE result, and of these, 140 had a full complement of the following tests: haemoglobin, white cell count, bilirubin, total protein, albumin, globulin, aspartate transaminase (AST), alanine transaminase (ALT), gamma glutamyl transferase (GGT), as well as the initial ALP.

Table 1 Numbers of patients in diagnostic groups

Group	No of patients (%)
Liver disease: (Acute hepatitis, chronic liver disease, biliary tract disease, tumours and drugs)	63 (30.9)
Bone disease: (Metastases, Paget's disease, osteomalacia, ankylosing spondylitis and recent fractures)	30 (14.2)
Coexistent liver and bone disease	6 (2.9)
Transient: (Heart failure, acute infections, diabetic ketoacidosis, thyrotoxicosis)	66 (32.3)
Other causes: (Leukaemias, polycythaemia, lymphoma, hypernephroma)	9 (4.4)
Diagnosis unknown: (Failure to satisfy diagnostic criteria, failure to attend follow up)	31 (15.2)
Total	204

Stepwise logistic regression analysis was used to determine which variables, if any, were useful in deciding which patients had liver or bone disease or transient increase in ALP. The classification merit statistics for these analyses showed that in the diagnosis of liver disease the only significant variables were bilirubin and GGT (using 1 n GGT), whereas in the diagnosis of bone disease the significant variables were, in decreasing order of significance, PAGE, bilirubin, white cell count and ALT. With PAGE removed from the analysis, the significant variables became GGT, initial ALP, bilirubin, and white cell count. Reintroduction of PAGE produced a significant improvement in prediction ($p = 0.001$) (table 2).

Similarly, for patients initially classified as "not liver" on the basis of GGT and bilirubin, addition of PAGE produced a small but significant improvement in prediction ($p = 0.018$).

We conclude that knowledge of the result of a qualitative test of ALP isoenzymes

(PAGE) provides useful additional information in the biochemical diagnosis of raised serum total ALP but that this is confined to the diagnosis of bone disease and when the GGT and bilirubin estimations are unhelpful. It is in this context (real life) that evaluations of the clinical usefulness of new methods of ALP isoenzyme determination should be set. In addition, established electrophoretic techniques permit identification of the so called biliary enzyme and Ig-ALP complexes, neither of which will be detected by the methods based on enzyme inhibition.

SG PARKER*

SA CAREY†

AW SKILLEN‡

JUDY M SIMPSON**

Departments of Medicine*

and Biochemistry,†

Ashington Hospital, Northumberland.

Departments of Clinical Biochemistry‡

and Medical Statistics,**

The Medical School, Framlington Place,

University of Newcastle upon Tyne, NE2 4HH.

The authors reply:

The clinical information presented in our paper was intended to confirm that the type of raised serum ALP isoenzyme activity detected by our method was appropriate for the clinical conditions tabulated, not to evaluate its usefulness in differentiating liver and bone disease. In routine diagnosis we believe that ALP isoenzyme estimations are useful in two important situations:

(i) In the patient with an isolated raised total ALP activity—that is, all other routine laboratory tests being normal

(ii) In the patient who has both bone and liver disease and, in addition, rising total ALP activity, such as the patient with renal failure receiving dialysis (renal osteodystrophy) who has hepatitis (not an uncommon occurrence).

In both of these situations tests such as ALT, AST, GGT and bilirubin are of little use as they are all normal in the first case and usually all raised in the second case. Furthermore, in the second case only qualitative ALP isoenzyme estimations would indicate which disease is deteriorating. Thus, although we agree that investigations other than ALP isoenzymes provide the diagnosis in most of the patients with raised ALP activity, we believe in selected clinical situations ALP isoenzyme measurements are justified.

Regarding the information presented by Parker *et al* on the merit of ALP isoenzyme in differentiating liver and bone disease using unselected cases with raised ALP activity, it is difficult to comment on the data provided without additional information such as the reasons why their qualitative isoenzyme measurements disagree with the clinical diagnosis in 20% of cases. Could this have some influence on the final conclusions about the merit statistics for ALP isoenzymes?

We would also be interested to know how ALP isoenzyme measurements were interpreted and whether the so called biliary

Table 2 Standard classification merit statistics with and without PAGE

	All patients				Not liver group		
	Liver v not liver (%)	Bone v not bone (%)	Without PAGE	With PAGE	Bone v not bone (%)	Without PAGE	With PAGE
Diagnostic accuracy	70.2	87.9	81.6	86.5	84.9	82.8	86.0
True positive	68.8	80.8	65.2	79.2	78.3	76.2	79.2
True negative	71.1	89.6	84.7	88.0	87.1	84.7	88.4
Sensitivity	55.0	63.6	45.5	57.6	66.7	59.3	70.4
Specificity	81.5	95.4	92.6	95.4	92.4	92.4	92.4