Cell surface expression of β_2 -microglobulin (β_2 m) correlates with stages of differentiation in B cell tumours

R A JONES, C S SCOTT, D R NORFOLK, A N STARK, J A CHILD

From the Department of Haematology, Regional Radiotherapy Centre, Cookridge Hospital, Leeds

SUMMARY Cell surface β_2 -microglobulin (β_2 m) densities of malignant B cells were determined by enzyme immunoassay in 97 cases of immunologically defined lymphoproliferative disease. Absolute β_2 m densities were found to depend on disease category with the lowest levels found on cells from chronic lymphocytic leukaemia (mean = 5.6 ng/ 10^6 cells, n = 27); atypical chronic lymphocytic leukaemia (mean = 5.9 ng/ 10^6 cells, n = 8); and prolymphocytoid chronic lymphocytic leukaemia variant (mean = 6.0 ng/ 10^6 cells, n = 16). β_2 m densities for B non-Hodgkin's lymphoma (n = 14) and B prolymphocytic leukaemia (n = 17) cases were 8.1 and 10.0 ng/ 10^6 cells, respectively, and the highest densities were found on cells from "late-B cell" tumours (mean = 14.3 ng/ 10^6 cells). Plasma cells from cases of Ig secreting tumours expressed unexpectedly low β_2 m densities (mean = 9.3 ng/ 10^6 cells; n = 6).

β_2 -microglobulin (β_2 m) forms the invariant light chain of the HLA class I molecule¹ and is expressed, in non-covalent association with the 43000 dalton heavy chain,² on the surface of cells from many tissues.^{3,4} HLA antigens seem to play a part in regulating the immune response by major histocompatibility complex restriction of T cell mediated cytotoxicity,⁵ and β_2 m is required for the structural integrity of the HLA molecule.⁶⁻⁸ Defective expression of HLA has been reported in a variety of tumours,^{4,9} and it is possible that this impairs T cell mediated responses to tumour cells.^{10,11} β_2 m is shed into plasma as a result of normal membrane turnover¹² and is raised in many pathological conditions.¹³ Correlations between serum β_2 m and estimated tumour mass have been noted in chronic lymphocytic leukaemia¹⁴ and myeloma,¹⁵ and serum β_2 m is also closely correlated with prognosis in myeloma.¹⁶ The underlying importance of these observations has yet to be determined. Further insights may be gained by examining relations between serum concentration and cell surface expression. We present the findings of a vertical study into surface β_2 m expression in malignant lymphoid cells from patients with morphologically and immunologically classified B cell disorders.

Material and methods

Mononuclear cell fractions were obtained from 97 cases of lymphoproliferative disease by density centrifugation (Lymphoprep: Nyegaard) of edetic acid and anticoagulated peripheral blood or bone marrow. Cell viabilities exceeded 90% in all cases.

All patients had B cell disorders as defined by monoclonal antibodies to B cell determinants (Leu12: CD19; and B1: CD20) and surface or cytoplasmic immunoglobulin light chain restriction. Cases were classified as chronic lymphocytic leukaemia (CLL: n = 27), prolymphocytic leukaemia (PLL: n = 17), non-Hodgkin's lymphoma (NHL: n = 14), or leukaemic reticuloendotheliosis (LRE: n = 7) by conventional morphological and previously described immunological criteria.¹⁷ A further 24 cases were investigated, comprising two groups defined as (a) cases with morphology suggestive of typical chronic lymphocytic leukaemia which showed increased (>20%) FMC7-positive components or surface immunoglobulin (SIg) densities (CLL-atyp; n = 8); and (b) cases which were morphologically and immunologically consistent with chronic lymphocytic leukaemia in "prolymphocytoid transformation"¹⁸ (CLL-Pro; n = 16). In addition, plasmacytoid and plasma cells from eight cases of immunoglobulin-secreting tumours were also examined: Walden-

β_2m expression in B cell tumours

strom's macroglobulinaemia (WM; $n = 2$) and myeloma ($n = 6$).

Surface β_2m determinations were performed by immunoenzyme assay¹⁹ of fractionated lymphoid cells from cases with leu12 or SIg/CIg light chain restriction, or both, in at least 80% of cells, or a peripheral white cell count $> 100 \times 10^9/l$ (two cases).

Statistical comparisons of surface β_2m expression by malignant B cells from the various diagnostic categories were made using the non-parametric Mann-Whitney U test.

Results

The fig shows the results of β_2m expression by malignant B cells in the different diagnostic categories. Cases of LRE and WM were combined to form the "late-B" group for statistical analysis. The lowest β_2m densities were found in the chronic lymphocytic leukaemia (mean 5.6 ng/ 10^6 cells), CLL-pro (mean 6.0), and CLL-atyp (mean 5.9) groups. No significant differences in the expression of β_2m were found between the three CLL groups, and for the purpose of subsequent statistical analysis, the CLL-pro and CLL-atyp groups were combined. Absolute β_2m values exceeding 9.0 ng/ 10^6 cells were found in one of 27

of CLL, two of 24 of atyp-CLL/CLL-pro, five of 14 of non-Hodgkin's lymphoma, 13 of 17 of prolymphocytic leukaemia, and nine of nine of late-B cases. When compared with the CLL groups, significantly increased surface β_2m densities were found in non-Hodgkin's lymphoma (mean 8.1) at $p < 0.05$ and in prolymphocytic leukaemia (mean 10.0) and late-B (mean 14.3) at $p < 0.001$. Surface expression was further increased in late-B compared with that in prolymphocytic leukaemia ($p < 0.01$) and non-Hodgkin's lymphoma ($p < 0.001$). Plasma cells, however, expressed significantly less ($p < 0.01$) β_2m than late B cells (mean 9.3), with mean expression not significantly different from non-Hodgkin's lymphoma, indicating decreasing expression with terminal B cell differentiation.

Discussion

The use of immunological markers provided a means whereby the various stages of B cell maturation could be defined. By examining the immunological profile of B cell neoplasms, it was possible tentatively to establish their relative location in order of differentiation as CLL, CLL-Pro, PLL, LRE, WM, and myeloma. It is evident from the results in this

Expression of cell surface β_2m by lymphoid cells in B cell malignancies. (—) indicates mean value for each category.

study that the increasing expression of surface β_2m closely parallels this sequence, with the notable exception of myeloma. In this respect β_2m is similar to the membrane densities of FMC7, B1/B4, SIg and HLADr (Ia) determinants, which increase with maturation but are lost at the terminal plasma cell stage.²⁰⁻²³ Interestingly, the prolymphocytoid variants of CLL (CLL-Pro), in which PLL-like morphology is seen despite retention of many of the CLL phenotypic characteristics,^{18 23 24} showed surface β_2m expression similar to that of typical CLL.

The position of non-Hodgkin's lymphoma in the differentiation pathway is less clear, as development from centroblasts to centrocytes within active lymph nodes is not solely related to the generation of antibody secreting cells.^{25 26} Considerable variation in surface β_2m expression was found in our cases of non-Hodgkin's lymphoma, and it is considered that this reflects the immunological heterogeneity of this group.

The function of surface β_2m with respect to B cell differentiation is unknown. Cytotoxic T cells recognise antigen in combination with the class I HLA molecule on the target cell surface.²⁷ β_2m has also been found in association with non-HLA molecules, such as H-Y and T1 antigens, and a structure cross reacting serologically with β_2m is an integral part of the Qa antigen.²⁸⁻³⁰ In vitro experimental evidence suggests that non-HLA β_2m may be associated with trigger molecules of B cell function,^{31 32} and it has been suggested that β_2m may have a role in regulating the expression of certain cell membrane antigens during cellular differentiation.²⁸ If this is the case then it is not unreasonable to speculate that membrane β_2m density will vary as a function of cellular maturity and that its decline at the terminal plasma cell stage is consistent with loss of requirement for surface components entailed in control of differentiation. A common feature of disease progression in various malignancies is raised serum β_2m . Its measurement is considered to be of value as an indicator of disease activity in CLL and is a powerful prognostic indicator in myeloma.³³ The origin of increasing β_2m production in malignant disease progression has not been established, but an inverse relation has been shown between surface β_2m expression and rate of secretion in carcinoma cell lines.³⁴ Non-haemopoietic tumours often exhibit reduced β_2m expression, compared with their normal counterparts,^{34 9} and it has been proposed that this may provide a means of evading T cell cytotoxic mechanisms,³⁵ since HLA expression on target cells is required for T cell cytotoxicity.³⁶ Whether B cell neoplasms behave in similar fashion has not been determined, but it is tempting to speculate that the rise in serum β_2m seen in the progression of some B cell tumours to more clinically aggressive

forms³³ may be accompanied by a reduction in surface β_2m expression. Studies are currently in progress to examine β_2m surface expression and secretion rates in relation to serum concentrations in B cell malignancies.

References

- Peterson PA, Rask L, Lindblom B. Highly purified papain-solubilised HL-A antigens contain beta 2-microglobulin. *Proc Natl Acad Sci USA* 1974;**71**:35-9.
- Cresswell P, Turner MJ, Strominger JL. Papain solubilised HL-A antigens from cultured human lymphocytes contain two peptide fragments. *Proc Natl Acad Sci USA* 1973;**70**:1603-7.
- Fleming KA, McMichael A, Morton JA, Woods J, McGee J. Distribution of HLA class I antigens in normal human tissue and in mammary cancer. *J Clin Pathol* 1981;**34**:779-84.
- Natalie PG, Bigotti A, Nicotra MR, Viora M, Manfredi D, Ferrone S. Distribution of human class I (HLA-A, B, C) histocompatibility antigens in normal and malignant tissues of nonlymphoid origin. *Cancer Res* 1984;**44**:4679-87.
- McMichael AJ. HLA restriction of human cytotoxic T cells. *Springer Seminars in Immunopathology* 1980;**3**:3-22.
- Lancet D, Parham P, Strominger JL. Heavy chain of HLA-A and HLA-B antigens is conformationally labile: a possible role for beta 2-microglobulin. *Proc Natl Acad Sci USA* 1979;**76**:3844-8.
- Parham P, Barnstable CJ, Bodmer WF. Use of monoclonal antibody (W6/32) in structural studies of HLA-A, B, C antigens. *J Immunol* 1979;**123**:342-9.
- Yokoyama K, Geier SS, Uehara H, Natheson SG. Secondary structure of the murine histocompatibility alloantigen H-2K^b. Relationship between heavy chain, beta 2-microglobulin and antigenic reactivity. *Biochemistry* 1985;**24**:3002-6.
- Turbitt ML, Mackie RM. Loss of Beta 2-microglobulin from the cell surface of cutaneous malignant and premalignant lesions. *Br J Dermatol* 1981;**104**:507-13.
- Gomard E, Duprez V, Henin Y, Levy JP. H-2 region product as determinant in immune cytolysis of syngeneic tumour cells by anti-MSV T lymphocytes. *Nature* 1976;**260**:706-9.
- Schrier PI, Bernards R, Vaessen RT, Houweling A, van der Eb AJ. Expression of class I major histocompatibility antigens switched off by highly oncogenic adenovirus 12 in transformed rat cells. *Nature* 1983;**305**:771-5.
- Cresswell P, Stringer T, Strominger JL, Turner MJ, Grey HM, Kubo RT. Immunological identity of the small subunit of HL-A antigens and beta 2-microglobulin and its turnover on the cell membrane. *Proc Natl Acad Sci USA* 1974;**71**:2123-7.
- Plesner T. Immunochemical studies of human beta 2-microglobulin. *Allergy* 1980;**35**:627-37.
- Spati B, Child JA, Kerruish SM, Cooper EH. Behaviour of serum beta 2-microglobulin and acute phase reactant proteins in chronic lymphocytic leukaemia. *Acta Haematol* 1980;**64**:79-86.
- Bataille R, Magub J, Grenier J, Donnadio D, Sany J. Serum beta 2-microglobulin in multiple myeloma: relation to presenting features and clinical staging. *Eur J Cancer Clin Oncol* 1982;**18**:59-66.
- Norfolk D, Child JA, Cooper EH, Kerruish S, Milford Ward A. Serum beta 2-microglobulin in myelomatosis: Potential value in stratification and monitoring. *Br J Cancer* 1979;**39**:510-5.
- Scott CS, Limbert HJ, Mackarill ID, Roberts BE. Membrane phenotypic studies in B cell lymphoproliferative disorders. *J Clin Pathol* 1985;**38**:995-1001.
- Stark AN, Limbert HJ, Roberts BE, Jones RA, Scott CS. Prolymphocytoid transformation of CLL: a clinical and immunological study of 22 cases. *Leuk Res* 1986;**10**:1225-32.

- 19 Jones RA, Scott CS, Child JA. Quantitation of cell-surface beta 2-microglobulin using a comparative antibody inhibition assay. *J Immunol Methods* 1987;**96**:87–96.
- 20 Gordon J, Mellstedt H, Aman P, Biberfeld P, Bjorkholm M, Klein G. Phenotypes in chronic B-lymphocytic leukemia probed by monoclonal antibodies and immunoglobulin secretion studies: Identification of stages of maturation arrest and the relation to clinical findings. *Blood* 1983;**62**:910–7.
- 21 Melo JV, San Miguel JF, Moss VE, Catovsky D. The membrane phenotype of hairy cell leukemia: A study with monoclonal antibodies. *Semin Oncol* 1984;**11**:381–5.
- 22 Zola H, Moore HA, Hohmann A, Hunter IK, Nikoloutsopoulos A, Bradley J. The antigen of mature human B cells detected by the monoclonal antibody FMC7: studies on the nature of the antigen and modulation of its expression. *J Immunol* 1984;**133**:321–6.
- 23 Caligaris-Cappio F, Janossy G. Surface markers in CLL of B cell type. *Semin Hematol* 1985;**22**:1–12.
- 24 Scott CS, Limbert HJ, Roberts BE, Stark AN. Polymphocytoid variants of CLL: an immunological and morphological survey. *Leuk Res* 1986 (in press).
- 25 Lennert K, Mohri H, Stein H, Kaiserling E, Muller-Hermelink H. In: *Handbuch der speziellen Pathologischen Anatomie und Histologie*. Berlin: Springer-Verlag, 1978;33–45.
- 26 Jones EL. Antibody probes in diagnosis and classification of lymphomas. *J Pathol* 1983;**141**:259–86.
- 27 Thorsby E. The role of HLA in T-cell activation. *Hum Immunol* 1984;**9**:1–7.
- 28 Fellous M, Gunther E, Kemler R, *et al*. Association of H-Y male antigens with β^2m on human lymphoid and differentiated mouse teratocarcinoma cell-lines. *J Exp Med* 1978;**148**:58–70.
- 29 Trucco M, de Petris S, Garotta G, Ceppellini R. Quantitative analysis of cell-surface HLA structures by means of monoclonal antibodies. *Hum Immunol* 1980;**3**:233–43.
- 30 Cotner T, Mashimo H, Kung PC, Goldstein G, Strominger JL. Human T-cell antigens bearing a structural relationship to HLA antigens. *Proc Natl Acad Sci USA* 1981;**78**:3858–62.
- 31 Messner RP. B2m: and old molecule assumes a new look. *J Lab Clin Med* 1984;**104**:141–5.
- 32 Ringden O, Moller E. Use of mitogens for the functional characterisation of human lymphocyte subpopulations. *Scand J Immunol* 1976;**5**:125–34.
- 33 Child JA, Kushwaha MR. Serum beta 2-microglobulin in lymphoproliferative and myeloproliferative diseases. *Hematol Oncol* 1984;**2**:391–401.
- 34 Nilsson K, Evrin P, Welsh KI. Production of beta 2-microglobulin by normal and malignant human cell lines and peripheral lymphocytes. *Transplantation Review* 1974;**21**:53–84.
- 35 Sanderson AR, Beverley PC. Interferon, beta 2-microglobulin and immunoselection in the pathway to malignancy. *Immunology Today* 1983;**4**:211–13.
- 36 Zinkernagel RM, Doherty PC. MHC cytotoxic T cells: studies on the biological role of polymorphic major transplantation antigens determining T cell restriction—specificity, function and responsiveness. *Adv Immunol* 1979;**27**:51–177.

Requests for reprints to: Dr RA Jones, Department of Haematology, Regional Radiotherapy Centre, Cookridge Hospital, Leeds LS16 6QB, England.