Letters to the Editor

Coronary artery dissection

Coronary artery dissection is an unusual cause of sudden death and the diagnosis is rarely made in life. It may, however, be more common than the few reported cases suggest, as it may be confused with thrombosis at necropsy unless a careful examination of the occluded vessel is made. We report two cases, the first a housewife aged 47, who was found dead in bed by her husband shortly after complaining of chest and arm pain, and the second a housewife aged 35 who died suddenly after complaining of similar symptoms while eating in a restaurant.

In both of these cases the cause of death was coronary occlusion due to coronary artery dissection. In the first, the occlusion was thought to be thrombotic at necropsy and the correct diagnosis was made only on histological examination. The dissection was recognised at necropsy in the second case after careful inspection of the vessel, the appearance initially suggesting thrombosis. The vessels affected were, respectively, the left circumflex and right coronary arteries, and in each case the dissection occurred close to the origin of the vessel. The histological appearances were virtually identical,

the dissection occurring, not within the media as in dissecting aortic aneurysm but adjacent to the external elastic lamina. (figure). This is presumably related to the structure of the coronary vessels, which are muscular rather than elastic arteries. Neither of the arteries was atheromatous or showed convincing mucoid medial degeneration. No intimal tear was demonstrable in relation to the dissection in either vessel, and there was no evidence of dissection or intimal tearing in the aortic roots.

Coronary artery dissection is a rare condition judging by the small number of reported cases. It accounted for 28 of 130 cases of non-aortic dissection gleaned by Guthrie and Maclean in their study,¹ which also showed a strong female preponderance, perhaps because pregnancy is a predisposing factor. Nalbandian and Chason² reviewed nine cases, including two of their own, and concluded that there was no common aetiological factor, but they did note that in their series the affected artery was, in all except one case, the anterior descending branch of the left coronary.

Distinguishing solitary coronary dissection from acute thrombotic occlusion would rarely be possible in life, but the correct diagnosis, albeit after death, in a relatively young person is of some importance for other members of the family; premature

coronary artery disease due to familial hyperlipoproteinaemia can then be excluded.

N GUBBAY

BW CODLING*

*Pathology Department,
Cheltenham General Hospital,
Sandford Road,
Cheltenham GL53 7AN
*Pathology Department,
Gloucester Royal Hospital,
Great Western Road,
Gloucester GL1 3NN*

References

- 1 Guthrie W, Maclean H. Dissecting aneurysms of arteries other than the aorta. *J Clin Pathol* 1972;108:219-35.
- 2 Nalbandian RM, Chason JL. Intramural (Intramural) dissecting hematomas in normal or otherwise unremarkable coronary arteries. *Am J Clin Pathol* 1965;43:348-56.

Granulocyte markers cathepsin G and leucocyte elastase are rarely observed in Reed-Sternberg and Hodgkin's cells

Recently there has been an increasing search for reliable histopathological markers of Reed-Sternberg and Hodgkin's cells in Hodgkin's disease. Initial demonstration of certain proteins and enzymes, such as immunoglobulin light and heavy chains, muramidase, albumin, α_1 -antitrypsin, and α_1 -antichymotrypsin were variously taken as evidence that these cells were of B cell or histiocytic lineage, or that they were effete, end stage cells with "leaky" membranes, permitting ready ingress of macromolecules. In the absence of direct evidence of synthesis or active synthetic apparatus in vitro, however, many of the conclusions drawn were purely presumptive. The availability of reliable markers for Reed-Sternberg and Hodgkin's cells in tissue sections is of more than academic importance; this has been emphasised more recently with the use of T cell markers. The use of these markers has often shown that many specimens previously classified as Hodgkin's disease are, in fact, of T cell lineage.

Accordingly, more recently, interest has arisen in Reed-Sternberg and Hodgkin's cell markers detectable by monoclonal antibodies. These have included antibodies such as Ki-1¹ directed against a population of lymphoid cells, and more recently, a range of antibodies reactive with granulocyte

Dissection has occurred adjacent to external elastic lamina (arrow).