- 2 Montalban C, Martinez-Fernandez R, Calleja JL, *et al.* Visceral leishmaniasis (kala-azar) as an opportunistic infection in patients infected with the human immunodeficiency virus in Spain. *Rev Infect Dis* 1989;11:655-60.
- 3 Berenguer J, Moreno S, Cercenado E, Bernaldo de Quiros JCL, Garcia de la Fuente A, Bouza E. Visceral leishmaniasis in patients infected with human immunodeficiency virus (HIV). *Ann Intern Med* 1989;111:129-32.
- 4 Montalban C, Calleja JL, Erice A, *et al.* Visceral Leishmaniasis in patients infected with human immunodeficiency virus. *J Infect* 1990;21:261-70.
- 5 Badaró R, Falcoff E, Badaró FS, *et al.* Treatment of visceral leishmaniasis with pentavalent antimony and interferon gamma. *N Engl J Med* 1990;322:16-21.

Persistent polyclonal B lymphocytosis with Epstein-Barr virus antibodies and subsequent malignant pulmonary blastoma

Lawlor and colleagues have reported a case of persistent polyclonal B lymphocytosis (PPBL) and subsequent malignant pulmonary blastoma and suggest a possible aetiological role for Epstein-Barr virus (EBV).¹ This conclusion was based on a single EBV viral capsid antigen (VCA), IgG titre of 1/1280, for which the method was not given. We cannot agree with this conclusion.

EBV antibody titres are usually measured by indirect immunofluorescence and a great deal has been published which describes this serological screening in normal persons and patients with EBV related diseases. Antibodies to EBV are present in most adults worldwide.² The extent of antibody responses to EBV antigens varies with age and immune function, with the prevalence of high antibody titres increasing with age in healthy persons.³ Overrepresentation of high titres of VCA specific IgG antibodies, compared with appropriate controls, has been observed in a large variety of non-malignant and malignant diseases.² Patients with the EBV associated malignancies, such as Burkitt's lymphoma and nasopharyngeal carcinoma (NPC), have substantially higher titres and a broader spectrum of antibodies to EBV determined antigens. Titres obtained in various laboratories are not always directly comparable and it is unfortunate that the method was not given in Lawlor's study.

Results obtained in our laboratory of EBV serology in various patient groups are presented in the table. Although the geometric mean titres of anti-EBV VCA IgG are higher in infectious mononucleosis and NPC, where EBV has a specific and well documented role, it is clear from the considerable overlap in the range of titres in all groups that individual measurement is a poor indicator of EBV disease. It is thus unacceptable to use a single titre of antibodies to any one antigen as a basis for implicating EBV.

The profile of antibody response to a range of EBV antigens is more reliable and may

assist in identifying potential aetiological roles for the virus.

In Lawlor's case report a role for Epstein-Barr virus in the pathogenesis of PPBL and malignant pulmonary blastoma has not been adequately established.

We agree with Sumaya⁴ that there is an urgent need for a collaborative effort by diagnostic and research laboratories to standardise the limits of "normal" EBV serological responses.

DE WYATT
PV COYLE
*Regional Virus Laboratory,
Royal Victoria Hospital,
Grosvenor Road,
Belfast BT12 6BN.*

- 1 Lawlor E, Murray M, Sullivan F, *et al.* Persistent polyclonal B lymphocytosis with Epstein-Barr virus antibodies and subsequent malignant pulmonary blastoma. *J Clin Pathol* 1991; 44:341-2.
- 2 Henle W, Henle G. Seroepidemiology of the virus. In: Epstein MA, Achong BG, eds. *The Epstein-Barr virus*. New York: Springer-Verlag, 1979:61-78.
- 3 Schmaier KE, van der Horst CM, Klotman ME. Epstein-Barr virus and the elderly host. *Rev Infect Dis* 1989;11:64-73.
- 4 Sumaya CV. Epstein-Barr virus serologic testing: diagnostic indications and interpretations. *Pediatr Infect Dis* 1986;5:337-42.

Dr Lawlor comments:

I fully agree with Wyatt and Coyle that there may be considerable overlap between EBV titres in a variety of conditions and that a wider antibody profile to EBV antigens where available is desirable. I would also support their comments on the necessity of standardisation of methods of EBV serology, particularly immunofluorescence, the method used in this case.

A number of apparently normal subjects can have a raised IgG VCA titre for some years after infectious mononucleosis.¹ In this particular patient, a middle aged woman with a benign lymphoproliferative disorder who subsequently developed a rare pulmonary tumour, the occurrence of a raised titre IgG VCA in the absence of evidence of generalised non-specific immune stimulation² (CMV 1/8, herpes 1/32, Toxoplasma (1 gM) 1/64 and IgG titres at the lower limit of normal), while not a definite indicator of EBV disease, suggest that the role of EBV should be considered in the aetiology of one or both of these conditions.

- 1 Horwitz CA, Henle W, Henle G, Rudnick H, Latts E. Long term serological follow-up of patients for Epstein-Barr virus after recovery from infectious mononucleosis. *J Infect Dis* 1985;151:1150-3.
- 2 Holmes GP, Kaplan JE, Stewart JA, Hunt B, Pinsky MPH, Schonberger LB. A cluster of patients with a chronic mononucleosis-like syndrome. *JAMA* 1987;257:2297-307.

BOOK REVIEWS

All titles reviewed here are available from the BMJ Bookshop, PO Box 295, London WC1H 9TE. Prices include postage in the United Kingdom and for members of the British Forces Overseas, but overseas customers should add £2 per item for postage and packing. Payment can be made by cheque in sterling drawn on a United Kingdom bank, or by credit card (Mastercard, Visa or American Express) stating card number, expiry date, and your full name.

Markers for Neural and Endocrine Cells. Molecular and Cell Biology, Diagnostic Applications. Ed M Gratzl, K Langley. (Pp 222; 48 figs; £67.) VCH. 1991. ISBN 3-527-28167-3.

It is a stated hope of the editors that the book should act as a bridge between fundamental research workers and clinicians by reviewing the latest data on the molecular biology, biochemistry and cell biology of various marker proteins for neural and endocrine cells. In large measure this has been achieved by uniformity of style and presentation which have made the text attractive to both basic scientists and clinicians. The usefulness of this book is greatly enhanced by a liberal number of illustrations and a list of relevant up to date references.

For those who have a particular interest in the relation between the nervous and endocrine systems, this book is to be recommended.

This multi-author book comprises seven chapters divided into three main sections: ontogenetic relations; molecular and cell biology; and applications of marker substances useful for characterising neurones and endocrine cells.

While neurones and endocrine cells share many properties they also differ in important respects (chapter 1). For example, they do not all share a common embryological origin from the ectodermal germ layer: examples include the endocrine cells associated with the digestive tract.

Further characterisation of membrane proteins of synaptic vesicles has allowed major progress to be made in the understanding of the biogenesis and life cycles of the various types of synaptic vesicle and the development of new concepts for neuronal secretion in the context of established pathways of exocytosis in other cells (chapter 2).

Chapter 3 reviews the biochemical, cell biological, and molecular biological features of neuroendocrine granules, and the immunohistochemical data of these proteins are summarised and compared in normal and human tumours of neuroendocrine cells.

Chapter 4 concentrates on aspects of cellular expression and describes the likely functions of a family of related glycoproteins collectively known as neural cell adhesion molecules (NCAM). It is now apparent that NCAM is not confined in adults to neurones

Titres of anti-EBV VCA IgG antibody in four patient groups

Patient group (n =)	Geometric mean titre	Titre range
Infectious mononucleosis (21)	353.3	80-10,240
Other suspected viral illnesses (29)	232.7	<10- 1280
NPC (35)	1358.3	160-10,240
Other head and neck tumours (19)	297.5	20- 5120