Expression of 120 kilodalton protein and cytotoxicity in *Helicobacter pylori*

J E Crabtree, N Figura, J D Taylor, M Bugnoli, D Armellini, D S Tompkins

Abstract

Antral biopsy culture supernatants from 14 subjects with chronic gastritis, known to have IgA antibodies to the 120 kilodalton protein, showed positive recognition of this antigen in western blots against a cytotoxin positive strain of *Helicobacter pylori* but gave negative reactions with two cytotoxin negative strains. Control immunoblots with culture supernatants from 13 non-responders to the protein were all negative. This indicates a direct association between expression of the 120 kilodalton protein in *H pylori* strains and cytotoxicity.

Peptic ulceration in patients with chronic gastritis has recently been shown to be specifically associated with mucosal IgA recognition of a 120 kilodalton protein of *Helicobacter pylori*.¹ Additionally, local IgA recognition of this protein is associated with the activity of the gastritis (polymorph infiltration) and epithelial surface degeneration.¹ Broth culture supernatants of *H pylori*, which have cytotoxic capacity, inducing vacuolation of cell lines in tissue culture, have recently been shown to contain a protein of similar molecular weight (128-130 kilodaltons).^{2,3} As patients with peptic ulceration are more frequently infected with cytotoxin producing strains of *H pylori* than those with gastritis alone,⁴ the possibility exists that the mucosal IgA recognition of the 120 kilodalton protein¹ is linked to the cytotoxicity of colonising strains.

In this study the relation between 120 kilodalton antigen recognition and bacterial cytotoxin production was investigated. The mucosal IgA response of *H pylori* positive patients known to have a 120 kilodalton antigen response¹ was examined by western blotting supernatants of in vitro cultured antral biopsy specimens against *H pylori* strains of known cytotoxin potential.

Methods

Twenty seven patients (mean (SEM) age 52.9 (15.8)) with chronic gastritis, who had previously been shown to have a positive mucosal IgA response to *H pylori* by western blotting,¹ and four subjects with histologically normal gastric mucosa were investigated. Thirteen of the 27 patients with chronic gastritis had no IgA antibodies to the *H pylori* 120 kilodalton protein and 14 patients had a positive IgA response.

Supernatants of three day in vitro cultured antral biopsy specimens were western blotted for *H pylori* IgA antibodies, as previously described.^{1,5} Bacterial preparations used for western blotting were whole cell sonicates of broth cultured cytotoxin positive (CCUG 17874) and cytotoxin negative (G-21 and G-17-91) strains of *H pylori*. Protein loading for sodium dodecyl sulphate polyacrylamide gel electrophoresis, culture supernatant dilutions, and immunoblotting conditions were identical for each strain examined. Bacterial culture and broth culture filtrate preparation were as previously described.⁴ Broth culture filtrates were assessed for cytotoxicity against HeLa, Vero, and CHO cells in vitro using the methods of Figura *et al.*⁴

Strain G-21, in addition to being cultured as above, was also grown on plates of Columbia agar with 7% horse blood, 1% Vitox (Oxoid), and the Skirrow selective supplement and used for comparative analysis with broth cultured G-21 in western blots.

Results

Patients with chronic gastritis all showed a positive mucosal IgA response to *H pylori* in western blots using all three strains of *H pylori*. No positive IgA response was observed in subjects with histologically normal antral mucosae. All the 14 patients previously defined as having a positive response to the 120 kilodalton protein¹ recognised this protein in immunoblots against the cytotoxin positive strain (CCUG 17874) (fig 1A). The 120 kilodalton protein was not recognised by these 14 subjects when their culture supernatants were immunoblotted against the cytotoxin negative (G-21 and G-17-91) strains (fig 1B). These patients showed considerably reduced recognition of other *H pylori* proteins when their culture supernatants were immunoblotted against the cytotoxin negative strains (fig 1B).

Culture supernatants of the 13 patients with chronic gastritis, previously defined as having no mucosal IgA antibodies to the 120 kilodalton protein,¹ showed no recognition of this protein when immunoblotted against either the cytotoxin positive or negative strains of *H pylori* (figs 1A and B). In these patients, there was considerable homogeneity in antigens recognised in immunoblots against both the cytotoxin positive and negative strains of *H pylori*. No difference in IgA recognition of bacterial proteins in western blots of culture

Department of
Medicine, St James's
University Hospital,
Leeds LS9 7TF

J E Crabtree
J D Taylor

Department of
Microbiology,
Bradford Royal
Infirmary, Bradford
D S Tompkins

Istituto di Malattie
Infettive, Università di
Siena, Siena, Italy
N Figura

Sclavo Research
Centre, Siena, Italy
M Bugnoli
D Armellini

Correspondence to:
Dr J E Crabtree

Accepted for publication
12 December 1991

Figure 1 IgA immunoblots of gastric culture supernatants (A) Cytotoxin positive *H. pylori* strain CCUG 17874. (B) Cytotoxin negative *H. pylori* strain G-17-91. Lanes 1–10 in A and B are identical patients. Lane 1: control with normal gastric mucosa; lanes 2–10: patients with chronic gastritis. Lanes 2–5: no recognition of the 120 kilodalton protein. Lanes 6–10: recognition of the 120 kilodalton protein in A but not in B. Protein molecular weight standards are given in kilodaltons.

supernatants against plate and broth grown *H. pylori* strain G-21 was apparent.

Discussion

Previous studies on systemic IgG responses to *H. pylori* have shown that patients who do not recognise the 120 kilodalton protein have no evidence of this protein in their homologous isolates.⁷ The association of a secretory IgA response to the 120 kilodalton protein with peptic ulceration and active gastritis suggests that strains of *H. pylori* expressing this protein are more pathogenic.¹ Whether the 120 kilodalton protein is the prime factor involved in inducing pathogenicity or whether it is a marker for cytotoxicity is presently unclear.

As broth culture filtrates of *H. pylori* strains with in vitro cytotoxic potential^{3,4,6} contain a 128–130 kilodalton protein associated with cytotoxicity^{2,3} we considered it important to examine, by western blotting, the mucosal IgA recognition of the 120 kilodalton protein against bacterial strains of defined cytotoxic potential. Our study clearly shows that immu-

nological recognition of the 120 kilodalton protein by the secretory IgA response was only evident when culture supernatants were immunoblotted against the cytotoxin positive strain of *H. pylori*. There was no recognition of the 120 kilodalton protein when cytotoxin negative strains were used as the antigen preparation. The marked reduction in recognition of other proteins by the 120 kilodalton positive patients when immunoblotted against the cytotoxic negative strains indicates considerable antigenic differences between the cytotoxin positive and negative strains. Interestingly, the patients negative for antigens to the 120 kilodalton protein showed very similar patterns of immunoreactivity in western blots irrespective of the strain used as antigen.

Recent analysis of the biotype and macromolecular profiles of cytotoxin positive and negative strains of *H. pylori* has emphasised that this bacterium is a genomically heterogeneous species.⁸ A close association between motility and cytotoxin production was observed in this latter study.⁸ Although the current study associates the recognition of the 120 kilodalton protein with bacterial cytotoxicity, further studies with purified 120 kilodalton protein are required to delineate firstly whether this protein is responsible for the in vitro cytotoxicity and secondly the relevance of this in vitro phenomenon to pathogenicity in vivo.

This study was undertaken with financial support from the Yorkshire Regional Health Authority and Glaxo plc.

- 1 Crabtree JE, Taylor JD, Wyatt JJ, *et al.* Mucosal IgA recognition of *Helicobacter pylori* 120 kDa protein, peptic ulceration and gastric pathology. *Lancet* 1991;338:332–5.
- 2 Figura N, Bugnoli M, Guglielmetti P, *et al.* Antibodies to vacuolating toxin of *Helicobacter pylori* in dyspeptic patients. In: Pajares JM, ed. *Helicobacter pylori, gastritis and peptic ulcer disease*. Berlin: Springer Verlag (in press).
- 3 Cover TL, Dooley CP, Blaser MJ. Characterisation of and human serological response to proteins in *Helicobacter pylori* broth culture supernatants with vacuolising cytotoxin activity. *Infect Immun* 1990;58:603–10.
- 4 Figura N, Guglielmetti A, Rossolini A, *et al.* Cytotoxin production by *Campylobacter pylori* strains isolated from patients with peptic ulcers and from patients with chronic gastritis only. *J Clin Microbiol* 1989;27:225–6.
- 5 Crabtree JE, Mahony MJ, Taylor JD, Heatley RV, Littlewood JM, Tompkins DS. Immune responses to *Helicobacter pylori* in children with recurrent abdominal pain. *J Clin Pathol* 1991;44:768–71.
- 6 Leunk RD, Johnson PT, David BC, Kraft WG, Morgan DR. Cytotoxic activity in broth-culture filtrates of *Campylobacter pylori*. *J Med Microbiol* 1988;26:93–9.
- 7 Apel I, Jacobs E, Kist M, Bredt W. Antibody response of patients against a 120 kDa surface protein of *Campylobacter pylori*. *Zbl Bakt Hyg A* 1988;268:271–6.
- 8 Owen RJ, Bickely J, Moreno M, Costas M, Morgan DR. Biotype and macromolecular profiles of cytotoxin-producing strains of *Helicobacter pylori* from antral gastric mucosa. *FEMS Microbiol Letts* 1991;79:199–204.