Papers

Can histopathologists reliably diagnose molar pregnancy?

A J Howat, S Beck, H Fox, S C Harris, A S Hill, C M Nicholson, R A Williams

Abstract

Aims—To assess the degree of difficulty in diagnosing partial mole by analysing intraobserver and interobserver agreement among a group of pathologists for these diagnoses.

Methods—Fifty mixed cases of partial mole, complete mole, and non-molar pregnancy were submitted to seven histopathologists, two of whom are expert gynaecological pathologists; the other five were district general hospital consultants, one of whom works in Australia. These participants gave each slide a firm diagnosis of either partial mole, complete mole, or non-molar pregnancy. Some 12 months later, the slides were recoded and again submitted for a second diagnostic round to assess intraobserver as well as interobserver agreement. Standard histological criteria for each diagnostic category were circulated with the slides.

Results— κ statistics showed that complete mole could be reliably distinguished from non-molar pregnancy, but neither non-molar pregnancy nor complete mole could be easily differentiated from partial mole. In only 35 out of 50 cases was there agreement between five or more of the seven participants. Agreement between the expert gynaecological pathologists was no better than for others in the group. Interestingly, the intraobserver agreement for each pathologist was good to excellent.

Conclusions—These results imply that the reported histological criteria are either not being applied consistently or that they are lacking in practical use. An atypical growth pattern of trophoblast, rather than the polar accentuation seen in normal first trimester pregnancies, seems to be the important diagnostic histological feature for partial mole. Ploidy studies might also help with problem cases.

(J Clin Pathol 1993;46:599-602)

In daily practice one recurring problem for histopathologists is whether products of conception show molar features or merely hydropic change associated with fetal death.^{1 2} This is especially so for partial moles which

may have fetal parts and membranes as well as villi, trophoblast, and decidua. There are, however, histological criteria that are said to easily distinguish between complete mole, non-molar pregnancy, and partial mole.¹⁻⁴ The diagnosis of partial mole or complete mole is important, with the patient having to enter the follow up surveillance programme for persistent trophoblastic disease and a request for her not to become pregnant; this entails measurement of urinary β human chorionic gonadotrophin for six to 12 months.^{5 6}

This study was designed to test how good histopathologists are at differentiating complete mole, partial mole, and non-molar pregnancy, to assess the value of the recognised histological criteria.

Methods

Fifty mixed cases of non-molar pregnancy, partial mole, and complete mole were selected from the files at Royal Preston Hospital and the Jessop Hospital for Women. Slides were coded and submitted to the seven participants. Some 12 months later, the slides were recoded and submitted for a second round. Table 1 shows the histological criteria sent with the slides. Ploidy studies were not carried out on these cases.

The results were then statistically evaluated for intra- and interobserver agreement as follows:

Table 1 Histological criteria

Non-molar	
Grossly normal/few vesicles	
Often fetus/fetal parts	
Variable hydropic change	
Atrophic attenuated trophoblast	
Occasional syncytial sprouts	
Partial mole	
Normal volume of placenta	
Often fetus/fetal parts	
Small vesicles mixed with normal villi	
Variable hydropic change	
Variable trophoblast hyperplasia	
Circumferential proliferation	
Central cisternal degeneration	
Scalloping of villi with trophoblast "inclusions"	
Some villi more normal with blood vessels	
Some small fibrosed avascular villi	
Complete mole	
Bulky uterus > dates	
Bunch of grapes grossly	
Rarely fetal tissue (if ever)	
Swollen avascular villi	
Variable trophoblast hyperplasia	

Department of
Histopathology, Royal
Preston Hospital,
Preston PR2 4HG
A J Howat
C M Nicholson

Department of
Histopathology,
Doncaster Royal
Infirmary
S Beck

Department of
Pathological Sciences,
University of
Manchester
H Fox

Department of
Histopathology,
Staffordshire General
Infirmary
S C Harris

Department of
Histopathology, Jessop
Hospital for Women,
Sheffield
A S Hill

Department of
Pathology,
Wangaratta District
Base Hospital,
Wangaratta, Victoria,
Australia
R A Williams

Correspondence to:
A J Howat

Accepted for publication
9 February 1993

Table 2 Histological opinions

Slide No	Pathologist													
	A		B		C		D		E		F		G	
	Run 1	2	1	2	1	2	1	2	1	2	1	2	1	2
1	NM	NM	NM	PM	PM	PM	PM	NM	PM	PM	PM	NM	NM	PM
2	NM	NM	NM	NM	NM	NM	NM	PM	NM	NM	PM	PM	NM	NM
3	NM	NM	PM	PM	PM	PM	PM	PM	NM	NM	CM	PM	NM	NM
4	NM	NM	PM	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM
5	NM	PM	NM	PM	NM	NM	NM	NM	NM	NM	NM	NM	NM	PM
6	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM
7	CM	CM	PM	CM	CM	CM	CM	CM	CM	CM	CM	CM	PM	PM
8	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM
9	PM	PM	PM	PM	NM	NM	PM	NM	PM	PM	PM	NM	PM	PM
10	PM	PM	PM	PM	PM	PM	PM	PM	PM	CM	CM	PM	PM	PM
11	CM	CM	CM	CM	CM	CM	CM	PM	CM	CM	CM	CM	PM	PM
12	NM	NM	PM	PM	NM	NM	NM	PM	NM	NM	NM	NM	PM	PM
13	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM
14	NM	NM	NM	NM	NM	NM	PM	NM	NM	NM	PM	NM	NM	NM
15	PM	PM	PM	PM	NM	PM	PM	PM	PM	PM	PM	PM	PM	PM
16	NM	PM	PM	PM	NM	NM	NM	NM	PM	PM	NM	NM	PM	NM
17	CM	CM	PM	CM	CM	CM	CM	PM	CM	CM	CM	CM	CM	CM
18	PM	CM	CM	CM	PM	CM	PM	PM	PM	CM	CM	CM	PM	PM
19	CM	CM	CM	CM	CM	CM	CM	CM	CM	CM	CM	CM	CM	PM
20	NM	NM	NM	PM	NM	NM	NM	PM	PM	PM	NM	PM	PM	PM
21	PM	PM	NM	PM	NM	NM	NM	PM	NM	NM	PM	PM	PM	PM
22	NM	NM	NM	NM	NM	PM	NM	NM	NM	NM	NM	NM	NM	NM
23	CM	CM	CM	CM	CM	CM	CM	CM	PM	PM	PM	CM	CM	PM
24	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM	PM	PM	NM	NM
25	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM
26	NM	NM	NM	PM	NM	NM	NM	NM	PM	PM	NM	NM	PM	PM
27	CM	CM	CM	CM	CM	CM	CM	CM	CM	CM	CM	CM	CM	CM
28	NM	NM	NM	NM	PM	PM	PM	PM	NM	NM	PM	PM	PM	PM
29	PM	NM	PM	PM	NM	NM	PM	PM	PM	PM	PM	PM	NM	PM
30	CM	CM	CM	CM	CM	CM	CM	CM	CM	CM	CM	CM	PM	PM
31	CM	CM	CM	CM	CM	CM	CM	PM	PM	CM	CM	CM	CM	PM
32	NM	NM	NM	NM	NM	NM	NM	NM	NM	PM	NM	NM	NM	NM
33	NM	PM	NM	PM	NM	NM	PM	NM	NM	PM	NM	NM	PM	PM
34	CM	CM	PM	CM	CM	PM	CM	CM	CM	CM	CM	CM	PM	CM
35	NM	PM	PM	PM	NM	NM	NM	NM	NM	NM	NM	NM	NM	PM
36	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM
37	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM
38	PM	PM	PM	PM	PM	PM	PM	NM	NM	PM	NM	NM	PM	PM
39	CM	CM	CM	CM	CM	CM	CM	CM	PM	CM	CM	CM	CM	CM
40	NM	PM	PM	PM	NM	NM	PM	PM	NM	NM	PM	PM	PM	PM
41	CM	CM	PM	CM	CM	CM	CM	CM	PM	CM	CM	CM	CM	CM
42	CM	CM	CM	CM	CM	CM	CM	CM	PM	CM	CM	CM	CM	CM
43	PM	CM	PM	CM	CM	CM	PM	PM	PM	CM	CM	CM	PM	PM
44	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM
45	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM
46	CM	CM	CM	CM	CM	CM	CM	CM	CM	CM	CM	CM	CM	CM
47	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM
48	CM	CM	CM	CM	CM	CM	CM	CM	CM	CM	CM	CM	CM	CM
49	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM	NM
50	NM	NM	PM	NM	NM	NM	PM	NM	NM	NM	NM	NM	NM	NM

NM: non-molar pregnancy; CM: complete mole; PM: partial mole.

(1) Consensus diagnosis—defined if greater than, or equal to, five pathologists agreeing for both runs;

(ii) interobserver agreement— κ value⁷ calculated for each “pair” of pathologists:

$$\kappa = \frac{Po - Pe}{1 - Pe}$$

where Po = observed agreement

Pe = agreement expected by chance

values 1.0 = perfect agreement

> 0.75 = excellent agreement beyond chance

0.4–0.75 = fair to good agreement beyond chance

< 0.4 = poor agreement beyond chance

0 = chance agreement only

(iii) Intraobserver agreement—percentage agreement between two runs for each pathologist.

Table 3 Distribution of diagnosis

Diagnosis	Number of slides with consensus
Non-molar pregnancy	17
Partial mole	4
Complete mole	14

Results

Table 2 shows the answers for each run. Only 35 out of the 50 slides achieved consensus. Table 3 gives the distribution of these cases. Of the 15 cases not reaching consensus, two were problems of differentiating partial mole from complete mole (cases 18 and 43). The other 13 involved the decision between non-molar pregnancy and partial mole. There was no problem in differentiating complete mole from non-molar pregnancy.

Table 4 Kappa values

	B	C	D	E	F	G
A	.589	.690	.745	.670	.586	.595
B		.491	.658	.455	.417	.561
C			.773	.563	.578	.501
D				.618	.628	.599
E					.466	.459
F						.393

Table 5 Intraobserver agreement

	Agreement %
A	84
B	74
C	90
D	72
E	74
F	82
G	82

Figure 1 Complete mole showing circumferential trophoblast hyperplasia and swollen avascular villi.

Figure 2 Partial mole showing villi with central cysternal degeneration, scalloping of villi with a "Norwegian fjord" periphery, and mild trophoblastic hyperplasia.

Table 4 shows the κ values using data from run 1. Agreement varied between poor (pathologist F v G) to excellent (C v D). The values are for all 50 cases, including the 15 cases for which no consensus was established.

Table 5 shows the intraobserver agreement. The values are good to excellent, ranging from 72–90%.

Discussion

About 15% of established pregnancies spontaneously abort; dilatation and curettage is often done in these cases to remove any retained products of conception.^{1,2} When villi and trophoblast are present in the products of

Figure 3 Non-molar pregnancy showing villi with mild hydropic change and no clinically relevant trophoblast hyperplasia.

conception, the pathologist must exclude trophoblastic disease, especially complete mole and partial mole. Complete mole can be reliably distinguished from non-molar pregnancy. In two cases complete mole could not be easily differentiated from partial mole, but this is of little clinical importance as all molar pregnancies should enter the programme for detection of persistent trophoblastic disease.⁵ Unfortunately (but not surprisingly), our study has identified problems differentiating non-molar pregnancy from partial mole. On review, it is clear that many cases of non-molar pregnancy showed significant hydropic change; the slides were purposely selected to show this feature. Nevertheless, some pathologists would be happy to leave these women and allow them to become pregnant again without follow up; others would impose restrictions on fertility, insisting on urinary β human chorionic gonadotrophin follow up. There are extensive histological criteria to avoid this problem (table 1).¹⁻⁴ Each pathologist seems to feel as though he or she can use these parameters consistently, as shown by the good intraobserver variation (table 5). These comments imply that either the histological criteria for partial mole are not being consistently applied among pathologists or that they are less than ideal for diagnosis. Our collective experience shows that in a non-molar hydropic pregnancy vesicles are hardly ever seen macroscopically. In partial mole one can see quite large (not small) vesicles mixed with normal villi. There is only mild trophoblast hyperplasia in most cases of partial mole and it is quite incorrect to say that there is hyperplasia of syncytiotrophoblast; syncytiotrophoblast is post-mitotic terminally differentiated tissue, incapable of mitotic activity. Recent studies with proliferating cell

nuclear antigen support the low level of trophoblast hyperplasia in partial mole.⁸ Other histological features, such as scalloping of villi and the presence of small fibrosed villi, are also seen in surgical terminations of pregnancy and are, in our opinion, unhelpful in differential diagnosis. The important feature in the diagnosis of partial mole is the *atypical* pattern of trophoblastic hyperplasia with a circumferential or multifocal pattern rather than the polar growth seen in normal first trimester placenta.

There are other diagnostic modalities that may help. Ploidy has been shown to be diploid in complete mole and frequently triploid in partial mole.⁹⁻¹² Non-molar pregnancy, if anembryonic pregnancies are included, shows a wide variety of cytogenetic and ploidy abnormalities including tetraploidy, trisomy, and triploidy.^{13,14} Tetraploid and diploid partial mole however, have been described.^{15,16} Assessment of ploidy involves either flow cytometry or static image analysis cytometry, both techniques being mostly unavailable in district general hospitals. Nevertheless, in cases where there is a serious problem in differentiating partial mole from non-molar pregnancy with hydropic change, sending some wet tissue or a block for ploidy studies might be prudent.

What is the importance of an erroneous diagnosis of non-molar pregnancy being made when the "correct" diagnosis should be partial mole? There are very few documented cases of persistent trophoblastic disease after partial mole; the incidence has been reported to vary from 0 of 51 cases partial mole¹⁵ to eight of 81 partial mole.¹⁷ Even cases of choriocarcinoma consequent on partial mole have been described.¹⁸⁻²¹ The risk is real, therefore, if very small.

There are problems with the routine diagnosis of partial mole. This conclusion is not novel.²² It seems that histopathology alone cannot solve this diagnostic dilemma, but the situation may be helped by improving the diagnostic criteria for partial mole along the lines that we have suggested.

We thank Preston and Chorley Hospitals Research Fund for financial assistance.

Thanks also go to Ms M Jones, medical statistician, Northern Regional Health Authority, Newcastle upon Tyne, for expert statistical help.

- 1 Buckley CH, Fox H. Biopsy pathology of the endometrium. In: *Biopsy pathology series 14*. London: Chapman and Hall, 1989:248-62.
- 2 Elston CW. Gestational trophoblastic disease. In: Fox H, ed. *Haines and Taylor. Textbook of obstetrical and gynaecological pathology*. Edinburgh: Churchill Livingstone, 1987:1045-78.
- 3 Szulman AE, Phillippe E, Boue JG, Boue A. Human triploidy: association with partial hydatidiform moles and non-molar conceptuses. *Hum Pathol* 1981;12:1016-21.
- 4 Szulman AE. Trophoblastic disease: clinical pathology of hydatidiform moles. *Obstet Gynecol Clin North Am* 1988;15:443-56.
- 5 Berkowitz RS, Goldstein DP. Diagnosis and management of the primary hydatidiform mole. *Obstet Gynecol Clin North Am* 1988;15:491-503.
- 6 Womack CC, Elston CW. Hydatidiform mole in Nottingham: a 12 yr retrospective epidemiological and morphological study. *Placenta* 1985;6:93-106.
- 7 Brennan P, Silman A. Statistical methods for assessing observer variability in clinical measures. *BMJ* 1992;304:1491-4.
- 8 Suresh UR, Hale RJ, Fox H, Buckley CH. Proliferating cell nuclear antigen (PCNA) immunoreactivity as a means for distinguishing hydropic abortions from partial hydatidiform moles. *J Clin Pathol* 1993;46:48-50.
- 9 Benirschke K. Flow cytometry for all mole-like abortion specimens. *Hum Pathol* 1989;20:403-4.
- 10 Lage JM, Driscoll SG, Yavner DL, Olivier AP, Mark SD, Weinberg DS. Hydatidiform moles. Application of flow cytometry in diagnosis. *Am J Clin Pathol* 1988;89:596-600.
- 11 Bagshawe KD, Lawler SD. Unmasking moles. *Br J Obstet Gynaecol* 1982;89:255-7.
- 12 Hemming JD, Quirke P, Womack C, Wells M, Elston CW, Bird CC. Diagnosis of molar pregnancy and persistent trophoblastic disease by flow cytometry. *J Clin Pathol* 1987;40:615-20.
- 13 Guerneri S, Bettio D, Simoni G, Brambati B, Lanzani A, Fraccaro M. Prevalence and distribution of chromosome abnormalities in a sample of first trimester internal abortions. *Hum Reprod* 1987;2:735-9.
- 14 Procter SE, Watt JL, Gray ES. Cytogenetic analysis of 100 spontaneous abortions in North-West Scotland. *Clin Genet* 1986;29:101-3.
- 15 Lage JM, Weinberg DS, Yavner DL, Bieber FR. The biology of tetraploid hydatidiform moles: histopathology, cytogenetics and flow cytometry. *Hum Pathol* 1989;20:419-25.
- 16 Lawler SD, Fisher RA, Dent J. A prospective genetic study of complete and partial hydatidiform moles. *Am J Obstet Gynecol* 1991;164:1270-7.
- 17 Berkowitz RS, Goldstein DP, Bernstein MR. Natural history of partial molar pregnancy. *Obstet Gynecol* 1986;66:677-81.
- 18 Gardner HAR, Lage JM. Choriocarcinoma following a partial hydatidiform mole: a case report. *Hum Pathol* 1992;23:468-71.
- 19 Looi LM, Siganesaratnam V. Malignant evolutions with fatal outcome in a patient with partial hydatidiform mole. *Aust NZ J Obstet Gynaecol* 1981;21:51-2.
- 20 Heifetz SA, Czaja J. In situ choriocarcinoma arising in partial hydatidiform mole: implications for the risk of persistent trophoblastic disease. *Paediatr Pathol* 1992;12:601-11.
- 21 Bagshawe KD, Lawler SD, Paradinas FJ, Dent J, Brown P, Boxer GM. Gestational trophoblastic tumours following initial diagnosis of partial hydatidiform mole. *Lancet* 1990;335:1074-6.
- 22 Javey H, Borazjani G, Behmard S, Langley FA. Discrepancies in the histological diagnosis of hydatidiform mole. *Br J Obstet Gynaecol* 1979;86:480-3.