Short reports

Lymphocyte subset counts in skin puncture and venous blood compared

S E Cracknell, R F Hinchliffe, J S Lilleyman

Abstract

To determine whether skin puncture blood can be used reliably for CD4 lymphocyte counts, the numbers of the major subsets of lymphocytes were assessed in paired venous and skin puncture blood samples from 22 children and 10 adults. Paired values were highly correlated, with skin puncture values being about 7% higher than venous values for each cell type. Differences were of borderline statistical significance for total lymphocytes and for each subset except CD3+ CD8+ T lymphocytes. Nevertheless, the magnitude of the differences was small and unlikely to be of clinical importance, and it seems that skin puncture samples may be preferable for CD4 counts in children or adults with difficult venous access.

(J Clin Pathol 1995;48:1137-1138)

Keywords: Lymphocyte subsets, CD4, CD8.

The counting of lymphocyte subsets is widely used in the detection and monitoring of immunodeficiency states, and CD4 counts are frequently requested in patients with HIV infection. Flow cytometric analysis of the major subsets requires only a sample volume of <1 ml, and this study was undertaken to determine whether skin puncture (capillary) blood is a suitable alternative to venous blood. If so, such a technique might be useful in paediatric practice.

Methods

Paired venous and skin puncture samples were taken from children being treated for acute leukaemia. Skin puncture samples taken for blood counts when attending an outpatient clinic were used, venous samples being taken around two hours later, at the time of routine administration of intravenous drugs. No child had blood taken simply for the purpose of this study. Paired samples were also taken from healthy adult volunteers.

Paired samples were treated identically. Lymphocyte subset numbers were determined by flow cytometry (FACScan, Becton Dickinson, Oxford, UK) using instrument defined gates (Simulset software version 2.3.2) and

a panel of combined monoclonal antibodies (Simulset, Becton Dickinson). Instrument standardisation and methodology conformed to current guidelines.¹ Absolute numbers of lymphocytes and their subpopulations were derived from an automated total white cell count and the three part differential provided by the FACScan. Statistical analysis was by the Wilcoxon signed rank test.

CD3+ CD4+ T lymphocyte values in paired venous and skin puncture samples.

Section of Paediatric Haematology, University of Sheffield Department of Paediatrics, The Children's Hospital, Western Bank, Sheffield S10 2THS
S E Cracknell
R F Hinchliffe
J S Lilleyman

Correspondence to: Dr R F Hinchliffe.
Accepted for publication 13 March 1995

Number of white cells, lymphocytes and lymphocyte subpopulations (median and range, $\times 10^9/l$) in venous and skin puncture blood

Sample type	Total white cells	Lymphocytes	CD3+ T cells	CD3+ CD4+ T cells	CD3+ CD8+ T cells	CD19+ B cells	CD3+ CD16+ CD56+ natural killer cells
Venous blood	4.6 (1.3–13.0)	1.46 (0.47–4.45)	1.26 (0.38–3.69)	0.71 (0.23–2.31)	0.49 (0.12–1.32)	0.03 (0–0.36)	0.10 (0.01–0.42)
Skin puncture blood	4.6 (1.6–14.2)	1.51 (0.43–4.62)	1.27 (0.41–3.93)	0.74 (0.25–2.50)	0.47 (0.09–1.66)	0.03 (0–0.38)	0.14 (0.02–0.48)
p value	0.009	0.03	0.036	0.033	0.12	0.05	0.049

Results

Samples were obtained from 22 children aged from eight months to 16.4 years, median 5.5 years, and 10 adults aged between 26 and 44 years. Data from adults and children were combined for analysis. The numbers of the different cell types were higher in skin puncture blood (4.8–6.4% higher for subsets other than CD3+ CD16+ CD56+ natural killer cells, which were 10.7% higher), the difference being statistically significant for all cell types except CD3+ CD8+ T lymphocytes (table). Significance was lost for all pairings when total lymphocytes and subpopulations were expressed in proportional terms (data not shown). Paired values were highly correlated ($r=0.90$ for CD3+ CD16+ CD56+ natural killer cells, and 0.96–0.98 for total white cells, lymphocytes and other subsets, $p<0.0001$ for all pairings). The differences between paired CD3+ CD4+ T cell counts is shown in the figure.

Discussion

Collection of blood by skin puncture provides a rapid and simple alternative to venepuncture when small volumes of blood are required, and is useful when dealing with small children and subjects with difficult veins. In this study no undue difficulty was found in obtaining the required volume of skin puncture blood; should problems arise it may be possible to reduce test volumes by 25–50%, at least when the total lymphocyte count is within normal limits.

Previous studies have shown clinically acceptable comparability in white cell and differential counts between the two samples types,^{2,3} although in the earlier study neutrophil values

were significantly higher in skin puncture blood. Similar results were obtained in the present study, the usually slightly higher total white cell count of skin puncture samples being reflected proportionally throughout the range of lymphocyte subsets. Although of borderline statistical significance for most subpopulations, the magnitude of the difference was small and unlikely to be of clinical importance.

The precision of both replicate haemoglobin and platelet estimations from skin puncture samples has been reported to be poorer than that from venous blood,^{4,5} although the latter claim could not be verified using the same sample collection method as that of the present study.² While we did not count lymphocytes in sequential paired samples there appears to be no reason why their numbers should fluctuate to a greater degree in skin puncture blood, a contention supported by the highly significant correlations found for each lymphocyte subpopulation between the two sample types.

Our results indicate that skin puncture blood can be safely used as an alternative to venous blood for the enumeration of CD4 and other lymphocyte subsets. It would be prudent when monitoring cell numbers over time, however, to use one sample type alone in order to minimise the variables affecting the results obtained.

- 1 Calvelli T, Denny TN, Paxton H, Galman R, Kagan J. Guidelines for flow cytometric immunophenotyping: a report from the National Institute of Allergy and Infectious Diseases, Division of AIDS. *Cytometry* 1993;14:702–15.
- 2 Bellamy GJ, Hinchcliffe RF. Venous and skin puncture blood counts compared. *Clin Lab Haematol* 1988;10:329–34.
- 3 Dreyer R, Pillay GS, Jacobs P. Skin puncture blood counts [letter]. *Clin Lab Haematol* 1994;16:304–5.
- 4 Reeves JD, Vinchinsky E, Addiego J Jr, Lubin BH. Iron deficiency in health and disease. In: Barnes LA, ed. *Advances in pediatrics*. Chicago: Year Book Medical Publishers, 1983:281–319.
- 5 Brecher G, Schneiderman M, Cronkite EP. The reproducibility and constancy of the platelet count. *Am J Clin Pathol* 1953;23:15–26.

J Clin Pathol 1995;48:1138–1140

Detection of Ki67 antigen by a new sheep polyclonal antiserum

G M Reynolds, D C Rowlands, G P Mead

Department of Pathology,
The Medical School,
University of Birmingham,
Edgbaston,
Birmingham B15 2TT
G M Reynolds
D C Rowlands

IDRL Department of Immunology
G P Mead

Correspondence to:
Mr G M Reynolds.

Accepted for publication
1 June 1995

Abstract

This report describes the characterisation of a polyclonal sheep antiserum against the Ki67 antigen. On western blots, this antiserum recognises a pair of bands of high molecular weight identical with those

seen with another polyclonal Ki67 antiserum and the MIB 1 monoclonal antibody. The new antiserum showed nuclear staining of a proportion of cells in paraffin wax embedded tissue sections following antigen retrieval using a microwave oven