Membrane staining for hepatitis B surface antigen on hepatocytes: a sensitive and specific marker of active viral replication in hepatitis B

C-M Chu, Y-F Liaw

Abstract

Aims—To test the hypothesis that membranous staining of hepatitis B surface antigen (HBsAg) on the hepatocyte is a marker of active viral replication in chronic hepatitis B virus (HBV) infection.

Methods—Intrahepatic expression of HBsAg and hepatitis B core antigen (HBcAg) was studied by indirect immunofluorescence on frozen sections of liver specimens from 75 patients with chronic hepatitis B, and the results were correlated with serum levels of HBV-DNA assayed by spot hybridisation.

Results—Hepatocyte HBcAg was detected in all of 20 patients with serum levels of HBV-DNA >1000 pg/ml, 18 (75%) of 24 patients with levels of HBV-DNA ≤1000 pg/ml, and two (6.5%) of 31 patients without detectable serum HBV-DNA. The concordance between hepatocyte HBcAg and serum HBV-DNA was 89.3% (67/75). There were six patients (8%) who had detectable serum HBV-DNA but without hepatocyte HBcAg, and two patients (2.7%) who had detectable hepatocyte HBcAg but without serum HBV-DNA. Membranous staining of HBsAg associated with variable degrees of cytoplasmic HBsAg was found in all but one of 44 patients with serum HBV-DNA, irrespective of the levels, but in none of the 31 patients without serum HBV-DNA. Of the latter, HBsAg was distributed solely in the cytoplasm. In addition, there is an inverse correlation between serum levels of HBV-DNA and the degrees of cytoplasmic staining of HBsAg. The concordance between membranous staining of HBsAg and serum HBV-DNA was 98.7% (74/75), significantly higher than that between hepatocyte HBcAg and serum HBV-DNA.

Conclusions—Membranous staining of HBsAg on the hepatocyte correlated excellently with serum HBV-DNA and thus can be recognised as a sensitive and specific marker of active hepatitis B virus replication.

(*J Clin Pathol* 1995;48:470-473)

Keywords: Hepatitis B, HBV-DNA, HBsAg, HBcAg, immunostaining.

Previous immunohistochemical studies have suggested that hepatocyte expression of hepatitis B core antigen (HBcAg) had a significant

diagnostic and prognostic implication in patients with chronic hepatitis B virus (HBV) infection.¹⁻⁸ The demonstration of HBcAg in liver usually correlated with active viral replication and was associated with variable degrees of inflammatory activity in the liver, while failure to detect HBcAg in liver usually indicated low levels of viral replication and was associated with little or no inflammatory activity.¹⁻⁸ Assays of HBV-DNA in serum by molecular hybridisation techniques have long been recognised as a sensitive and specific method of evaluating viral replication in chronic hepatitis B virus infection.⁹⁻¹¹ However, these earlier data have shown that the discordance between serum HBV-DNA and hepatocyte expression of HBcAg ranged from 12% to 23%, including (1) 9-23% of patients who were positive for serum HBV-DNA without hepatocyte HBcAg and (2) 0-3% of patients who had hepatocyte expression of HBcAg without serum HBV-DNA.¹²⁻¹⁵ One possible reason for the former might be sampling error in the biopsy specimens because hepatocyte expression of HBcAg could be only focal in certain patients with chronic hepatitis B^{16,17}; in the case of the latter, false positive immunostaining of HBcAg cannot be excluded.¹⁸

It has been shown that there is membranous staining of HBsAg associated with variable degrees of cytoplasmic staining of HBsAg during the phase of active hepatitis B virus replication and, by the contrast, solely cytoplasmic staining of HBsAg during the non-replicative phase.^{12,19} Of note is that the distribution of membranous staining of HBsAg is usually diffuse and the staining intensity is rather strong during active viral replication.^{12,19} It is therefore postulated that immunostaining of HBsAg on the hepatocyte membrane might be more sensitive and specific than immunostaining of HBcAg as a marker of active hepatitis B virus replication. To test this hypothesis, serum levels of HBV-DNA were correlated with membranous staining of HBsAg on the hepatocyte versus hepatocyte staining of HBcAg in 75 patients with chronic hepatitis B.

Methods

Seventy five patients with chronic HBV infection were studied. All had been HBsAg positive for more than six months before a histological diagnosis of chronic hepatitis was made. Forty four patients were seropositive for HBeAg and the other 31 were seropositive for antibody against HBcAg (anti-HBe). All were

Liver Unit, Chang Gung Memorial Hospital, Chang Gung Medical College, 199 Tung Hwa North Road, Taipei, Taiwan
C-M Chu
Y-F Liaw

Correspondence to:
Dr Chia-Ming Chu.

Accepted for publication
26 September 1994

Table 1 Mean (SEM) clinical and laboratory data of patients studied

Category	No of cases	Age (years)	Sex (M:F)	AST (IU/l) (n <40)	ALT (IU/l) (N <40)
HBeAg positive CPH	22	24 (3)	18:4	48 (2)	60 (3)
HBeAg positive CAH	22	29 (2)	20:2	84 (8)	202 (18)
Anti-HBe positive CAH	5	33 (2)	5:0	80 (30)	180 (40)
Anti-HBe positive MHC	26	35 (1)	24:2	29 (3)	34 (3)

HBeAg=hepatitis B e antigen; anti-HBe=antibody against HBeAg; CPH=chronic persistent hepatitis; CAH=chronic active hepatitis; MHC=minimal histological changes.

negative for antibody against hepatitis δ virus (anti-HDV). None admitted intravenous drug abuse, nor had they ever received antiviral or immunosuppressive treatment. The histological diagnosis of chronic hepatitis was made according to standard criteria.²⁰ The clinical and laboratory data of the patients studied are listed in table 1.

Serum aspartate transferase (AST) and alanine transaminase (ALT) were measured by sequential multiple autoanalysers. Serum HBsAg, HBeAg, anti-HBe, and anti-HDV were assayed using commercially available radioimmunoassay kits (Ausria-II, HBeAg-RIA, and anti-delta, Abbot Laboratories).

Serum specimens for test of HBV-DNA were collected on the same day of liver biopsy and stored at -70°C until use. Serum HBV-DNA was assayed by spot hybridisation techniques using ^{32}P labelled cloned HBV-DNA, as reported previously.²¹

Liver specimens were obtained by percutaneous needle biopsy with a Menghini needle. Fragments of specimens were snap frozen in isopentane cooled with liquid nitrogen and stored at -70°C until use. Samples of the same biopsy specimens were also fixed in 10% formaldehyde and embedded in paraffin wax for routine histological diagnosis. Cryostat sections ($5\text{ }\mu\text{m}$) were dried overnight at room temperature and fixed in carbon tetrachloride at 4°C for 10 minutes, followed by an extensive wash with phosphate buffered saline (pH 7.2) before staining. Intrahepatic expression of

HBcAg was studied by indirect immunofluorescence using rabbit anti-HBc (Dako), followed by fluorescein isothiocyanate (FITC) labelled swine anti-rabbit immunoglobulin G. Intrahepatic expression of HBsAg was studied by indirect immunofluorescence using murine monoclonal anti-HBs (Chemicon), followed by FITC conjugated rabbit anti-mouse immunoglobulin G (Jackson Immuno Research Laboratories).

Statistical analyses were performed using χ^2 test with Yate's correction, Wilcoxon rank sum test, or Spearman's rank correlation where appropriate.

Results

Hepatocyte staining of HBcAg was detected in 40 patients, including 20 with predominantly nuclear staining and 20 with predominantly cytoplasmic staining. Serum HBV-DNA was detected in 44 patients, including 20 with levels $>1000\text{ pg/ml}$ and 24 with levels $\leq 1000\text{ pg/ml}$. As shown in table 2, the concordance between serum HBV-DNA and hepatocyte staining of HBcAg was 89.3% (67/75).

Figure 1 summarises the correlation between serum levels of HBV-DNA and hepatocyte staining of HBcAg. Of the 44 patients positive for serum HBV-DNA, 38 (86.4%) had HBcAg detectable in liver. All of the 20 patients with serum levels of HBV-DNA $>1000\text{ pg/ml}$ had detectable HBcAg in liver, with 70% (14/20) predominantly in the nucleus and 30% (6/20) predominantly in the cytoplasm, whereas only 75% (18/24) of those with serum levels of HBV-DNA $\leq 1000\text{ pg/ml}$ had detectable HBcAg in liver ($p<0.05$), with 67% (12/18) predominantly in the cytoplasm and 33% (6/18) predominantly in the nucleus ($p<0.05$). Serum levels of HBV-DNA were significantly higher in patients with predominantly nuclear staining of HBcAg than in those with predominantly cytoplasmic staining of HBcAg ($p<0.01$). Five of six patients who had HBV-DNA detectable in serum without hepatocyte staining of HBcAg had levels of HBV-DNA $\leq 500\text{ pg/ml}$.

Hepatocyte staining of HBsAg was detected in all of the 75 study patients, including six with purely membranous staining, 32 with purely cytoplasmic staining, and 37 with mixed membranous and cytoplasmic staining. Figure 2 summarises the correlation between serum levels of HBV-DNA and hepatocyte staining of HBsAg. All but one of 44 patients with serum HBV-DNA, including the six patients who had

Figure 1 Correlation between serum levels of HBV-DNA and hepatocyte staining of HBcAg in 75 patients with chronic hepatitis B. Serum levels of HBV-DNA were semiquantitatively scored on a 0 to 4+ scale, corresponding to undetectable, $\leq 500\text{ pg/ml}$, 501–1000 pg/ml, 1001–2000 pg/ml, and $>2000\text{ pg/ml}$. The empty circles indicate patients with membranous staining of HBsAg and the filled circles indicate patients with solely cytoplasmic staining of HBsAg. Spearman's rank correlation coefficient = -0.64 , $p<0.001$.

Table 2 Results of immunostaining of HBcAg on hepatocyte and serum HBV-DNA in 75 patients with chronic type B hepatitis

Category	Hepatocyte HBcAg/serum HBV-DNA			
	+/-	-/-	-/+	+/+
HBeAg positive CPH	20	1	1	0
HBeAg positive CAH	15	3	4	0
Anti-HBe positive CAH	3	1	1	0
Anti-HBe positive MHC	0	24	0	2
Total	38	29	6	2

HBcAg=hepatitis B core antigen; HBV=hepatitis B virus. For other abbreviations see table 1.

Instead it is suggested that the diffuse membranous staining of HBsAg during active HBV replication indicates absorption of circulating hepatitis B virus virions on the liver cell surface; the virions are surrounded by HBsAg which contains polymerised human albumin receptor and could thus be absorbed on the liver cell surface.^{24,25}

In conclusion, membranous staining of HBsAg on the hepatocyte correlated excellently with serum HBV-DNA and could be recognised as a marker of active hepatitis B virus replication. It was suggested that membranous staining of HBsAg possibly indicates absorption of circulating hepatitis B virus virions on the cell surface rather than expression of viral proteins on the cell membrane of infected cells.

The authors thank Ms Tsai MH and Leu HL for preparation of this manuscript.

- Gudat F, Bianchi L, Sonnabend W, Thiel G, Aenishaenslin W, Stalder GA. Pattern of core and surface expression in liver tissue reflects state of specific immune response in hepatitis B. *Lab Invest* 1975;32:1-9.
- Ray MB, Desmet VJ, Bradburne AF, Desmyter J, Fevery J, DeGroote J. Differential distribution of hepatitis B surface antigen and hepatitis B core antigen in liver of hepatitis B patients. *Gastroenterology* 1976;71:462-7.
- Yamada G, Nakane PK. Hepatitis B core and surface antigens in liver tissue. Light and electron microscopic localisation by the peroxidase-labeled antibody method. *Lab Invest* 1977;36:649-59.
- Huang SN, Neurath AR. Immunohistologic demonstration of hepatitis B viral antigens in liver with reference to its significance in liver injury. *Lab Invest* 1979;40:1-17.
- Chu CM, Liaw YF. Intrahepatic distribution of hepatitis B surface and core antigens in chronic hepatitis B virus infection: hepatocyte with cytoplasmic/membranous hepatitis B core antigen as a possible target for immune hepatocytolysis. *Gastroenterology* 1987;92:220-5.
- Hsu HC, Su IJ, Lai MY, Chen DS, Chang MH, Chuang SM, et al. Biologic and prognostic significance of hepatitis B core antigen expression in the natural course of chronic hepatitis B virus infection. *J Hepatol* 1987;5:45-50.
- Yoo JY, Howard R, Waggoner JG, Hoofnagle JH. Peroxidase-anti-peroxidase detection of hepatitis B surface and core antigen in liver biopsy specimens from patients with chronic type B hepatitis. *J Med Virol* 1987;23:273-81.
- Kakumu S, Arao M, Yoshioka K, Tsutsumi Y, Inoue M. Distribution of HBcAg in hepatitis B detected by immunoperoxidase staining with three different preparations of anti-HBc antibodies. *J Clin Pathol* 1989;42:284-8.
- Weller IVD, Fowler MJF, Monjardino J, Thomas HC. The detection of HBV-DNA in serum by molecular hybridization. *J Med Virol* 1982;9:273-80.
- Karayiannis P, Fowler MJF, Lok ASF, Greenfield G, Monjardino J, Thomas HC. Detection of serum HBV-DNA by molecular hybridisation: correlation with HBeAg/anti-HBe status, racial origin and liver histology. *J Hepatol* 1985;1:99-106.
- Berninger M, Hammer M, Hoyer B, Gerin JL. An assay for the detection of the DNA genome of hepatitis B virus in serum. *J Med Virol* 1982;9:57-68.
- Bonino F, Hoyer B, Nelson J, Engle R, Verme G, Gerin JL. Hepatitis B virus DNA in the sera of HBsAg carriers: a marker of active hepatitis B virus replication in the liver. *Hepatology* 1981;1:386-91.
- Hadziyannis SJ, Leiberhan HM, Karvountzis GB, Shafritz DA. Analysis of liver disease, nuclear HBcAg, viral replication, and hepatitis B virus DNA in liver and serum of HBeAg vs anti-HBe positive carriers of hepatitis B virus. *Hepatology* 1983;3:656-62.
- Freiman J, Eckstein R, McCaughan G, Parsons C, Davies JS, Dieguts P, et al. Significance of serum and hepatic markers of hepatitis B viral infection in HBsAg-positive and HBsAg-negative chronic active hepatitis. *Hepatology* 1985;5:50-3.
- Wu JC, Lee SD, Wang HY, Ting LP, Tsai YT, Tsay SH, et al. Correlation between hepatic hepatitis B core antigen and serum hepatitis B virus-DNA levels in patients with chronic hepatitis B virus infection in Taiwan. *Arch Pathol Lab Med* 1987;111:181-4.
- Chu CM, Liaw YF. Hepatocyte expression of HBcAg and serum HBeAg in hepatitis B. Comparison of polyclonal and monoclonal antibodies during a trial of interferon. *J Clin Pathol* 1991;44:21-4.
- Chu CM, Liaw YF. Immunohistological study of intrahepatic expression of hepatitis B core and e antigen in chronic type B hepatitis. *J Clin Pathol* 1992;45:791-5.
- Goodman ZO, Langloss JM, Bratthauer GL, Ishak KG. Immunohistochemical localization of hepatitis B surface antigen and hepatitis B core antigen in tissue sections. A source of false positive staining. *Am J Clin Pathol* 1988;89:533-7.
- Montano L, Miescher GC, Goodall AH, Wiedmann KH, Janossy G, Thomas HC. Hepatitis B virus and HLA antigen display in the liver during chronic hepatitis B virus infection. *Hepatology* 1982;2:557-61.
- International Group. Acute and chronic hepatitis revisited. *Lancet* 1977;iii:914-9.
- Chu CM, Karayiannis P, Fowler MJF, Monjardino J, Liaw YF, Thomas HC. Natural history of chronic hepatitis B virus infection in Taiwan: studies of hepatitis B virus DNA in serum. *Hepatology* 1985;5:431-4.
- Ou JH, Yeh CT, Yen TSB. Transport of hepatitis B virus precore protein into the nucleus after cleavage of its signal peptide. *J Virol* 1989;63:5238-43.
- Gowans EJ, Burrell CJ. Widespread presence of cytoplasmic HBcAg in hepatitis B infected liver detected by improved immunochemical methods. *J Clin Pathol* 1985;38:393-8.
- Imai M, Yanase Y, Mojri T. A receptor for polymerized human and chimpanzee albumins on hepatitis B virus particles co-occurring with HBeAg. *Gastroenterology* 1979;76:241-7.
- Thung SN, Gerber MA. Hepatitis B virus and polyalbumin binding sites. *Gastroenterology* 1983;83:466-8.