THE RELATION BETWEEN POLYAGGLUTINABILITY OF ERYTHROCYTES *IN VIVO* AND THE HÜBENER-THOMSEN-FRIEDENREICH PHENOMENON

BY

J. REEPMAKER

From the Central Laboratories of the Netherlands Red Cross Blood Transfusion Service, and the University Laboratory of Hygiene, Amsterdam

(RECEIVED FOR PUBLICATION DECEMBER 20, 1951)

Hübener (1925) and Thomsen (1927a and b, 1928) were the first to report that erythrocytes from old blood samples might sometimes be agglutinated by sera irrespective of their blood groups; in other words, they become panagglutinable. Friedenreich (1930) demonstrated that this phenomenon, which may be a cause of error in blood grouping, was due to contamination of the blood with certain enzyme-producing bacteria which cause a new agglutinin, "T," to appear at the red cell surface and it reacts with an agglutinin "T" present in all adult sera; it is apparently absent in newborn sera. The transformation process is generally known as the Hübener-Thomsen-Friedenreich phenomenon. The T antibody is a specific cold agglutinin which is adsorbed by "changed" cells only. It exists apart from the other known specific antibodies.

Friedenreich (1930) isolated two coryneform bacteria, "M" and "J," from old blood samples. Panagglutinability was induced either by direct inoculation of blood or with culture filtrates of these bacteria. Friedenreich also found that an enzyme with a similar action was produced by some strains of *Vibrio*, including *Vibrio cholerae*, and also by a strain of the *Coccobacillus pierantonii*. Other bacteria were added by later investigators, e.g., some strains of actinomycetes, especially among the acid-forming groups (von Magnus, 1936), a Gram-positive coccus (Terada, 1936), a Gram-negative coccobacillus (Kossovitch and Chabaud, 1938), a corynebacterium similar to the Orla-Jensen type, the *Corynebacterium H (Hektoenii)* (Davidson and Toharsky, 1940, 1942), and *Cl. welchii*, *Vibrio proteus*, pneumococci of all types, and some species of streptococci and staphylococci (Chu, 1948).

Viruses of the mumps-influenza group are also capable of forming a T antigen (Burnet, McCrea, and Stone, 1946; Stone, 1947; Chu and Coombs, 1947) and, finally, the same effect results from treatment with snake venom (Briody, unpublished

data), potassium periodate (Stewart, 1949; Moskowitz and Treffers, 1950), and also with trypsin under certain conditions (Rosenthal and Schwartz, 1951).

A similar phenomenon occurring *in vivo* has been less frequently observed. This is polyagglutinability of the red cells. Twelve well-documented cases have been described in the literature (Table I). In none of them was there convincing evidence that bacterial infection caused the polyagglutinability *in vivo*.

A further case, in which an organism was isolated from the urine of the patient which was able to render red blood cells panagglutinable *in vitro*, is described in this paper.

Case History

H.B., aged 25, was healthy until 1943. He then suffered from appendicitis and scarlatina. In 1946 he developed symptoms of pyelitis and was treated with sulphonamides. In the same year he was admitted to hospital at Djakarta because of amoebiasis and orchitis, probably following epidemic parotitis. By March, 1947, he had recovered. From April, 1947, onwards he has suffered from a relapsing infection of the urinary passages. This had been treated occasionally with sulphonamides, but without lasting effect. Pyelograms showed no abnormality, and cultures and guinea-pig inoculations were negative for tuberculosis.

A diagnosis of benign but persistent urinary infection, probably with prostatitis, was made. In June, 1951, the red blood cells were found to be polyagglutinable.

Serology and Urine Culture.—Till June 22, 1951, the patient's red blood cells (group A₁), containing a strong A antigen, had been regularly used as test erythrocytes for blood group determinations without giving abnormal results. On this date, however, it was found that they were agglutinated by nearly all sera of the same blood group and also by AB sera. An investigation of the exact frequency was not made at that time, but three days later the patient's corpuscles were titrated at four different temperatures against two normal A sera as well as his own serum (Table II).

TABLE I
ANALYSIS OF CASES OF POLYAGGLUTINABILITY OF RED CELLS

Investigators	Diagnosis	Sex and Age	Blood Group	Percentage of Agglutinating Sera	Optimum Temperature (°C.)	Auto-agglutination	Bacterial Infection
Levine and Katzin (1938) ..	Pneumococcal sepsis after measles	Child 4 years	O	15	20	Absent	Pneumococci type I and II
Gaffney and Sachs (1943) ..	Congenital syphilis	Male 11 years	O	47-89	4: 1/32 16: 1/16	"	<i>Treponema pallidum</i>
	Healthy	Female	O	73	4: 1/16 16: 1/8	"	Absent
Basil-Jones, Sanger, and Walsh (1946) ..	Puerperal sepsis and purulent parametritis after partus arte praematurus	Female	O	100 (45/45)	4 and 22	"	Gram-positive rods in parametric abscesses
Boorman, Loutit, and Steabben (1946) ..	Incomplete abortion	Female 29 years	O	±100 (38/39)	22	"	Infection present, bacteria not isolated
	Hysterectomy for parametritis	Female 37 years	O	±100 (142/143)	22	"	Idem
Engleson and Grubb (1949)	Parametritis	Female 28 years	B	100 (7/7)	22	Present at 22° C., not at 37° C.	"
	Sepsis with pyodermia	Child 9 years	A	100 (7/7)	4	Present at 22° C. and 37° C.	"
Henningsen (1949) ..	Healthy	Child 4 years	O	84 (adult sera) 18-4 (child's sera)	4	?	Absent
van Loghem, van der Hart, and Heier (1951) ..	Multiple tumour; metastases	Female 54 years	A	—	16	Absent	Blood culture negative. Gram-positive diplococci once isolated from sputum
Holländer (1951) ..	Pneumococcal meningitis	Child 9 months	O	95 infant sera negative	22	Present 4° C. 1/32 16° C. 1/16	Pneumococci type 19
Gasser and Holländer (1951)	Acute acquired haemolytic anaemia with thrombocytopenic purpura	Child 7 weeks	O	±100 (24/25)	22	Present at 37° C.	Absent
Reepmaker (195) ..	Urine infection	Male 25 years	A	—	16	Absent	α-Haemolytic streptococcus

TABLE II
AGGLUTINATION TITRES OF PATIENT'S RED CELLS WITH TWO NORMAL GROUP A SERA AND OWN SERUM AT FOUR DIFFERENT TEMPERATURES

Serum	4° C.	16° C.	22° C.	37° C.
Patient ..	—	—	—	—
A No 11139 ..	1:4 (+)	1:2 (+)	1:2 ±	—
A No. 11141 ..	1:8 (+)	1:8 ±	1:8 ±	1:1 (+)

The strongest reactions and the highest titres were obtained at 4° C. Auto-agglutination did not take place. By June 26, 1951, the polyagglutinability had totally disappeared.

In view of the history of chronic urinary infection the patient's urine was cultured to determine whether there was a relationship between the infection and the abnormal agglutinability of the red blood cells.

Two different micro-organisms were isolated and cultivated in broth. Filtrates made from these cultures were added to suspensions of thrice-washed group O erythrocytes which were then kept at room temperature for about an hour. The mixture was then centrifuged and the cells again washed three times. Two drops of a 4% suspension of treated cells were then added to two drops of AB serum on a slide and the reactions read after five minutes at room temperature. Only erythrocytes treated with a filtrate derived from a Gram-

positive streptococcus were agglutinated by the AB serum.

Further Investigations.—A filtrate of a four-day-old culture of the patient's streptococcus in serum broth proved to be the most useful for our experiments. The strain was dried *in vacuo* with phosphorus pentoxide in order to prevent variation through continuous re-inoculation. Dr. Skadhauge, from the Serum Institute at Copenhagen, kindly investigated the properties of this strain. It showed α-haemolysis on 5% blood agar and did not grow on 40% bile-blood agar. It did not split arginine with the formation of NH₃, nor sodium hippurate. It fermented lactose, sucrose, and trehalose, but not mannitol, salicin, aesculin, inulin, raffinose, glycerol, glycogen, or arabinose. According to Dr. Skadhauge the streptococcus belonged to a species, also isolated from urine, which was discovered by Gardborg and Borgen (1951). Dr. Friedenreich, in a personal communication, reported that this Oslo strain also caused panagglutinability of red blood cells.

Preparation of Panagglutinable T Erythrocytes.—Four drops of packed red cells were added to a mixture of equal volumes of streptococcus filtrate and normal saline and the suspension left at room temperature for half an hour. After centrifuging the red cells were washed three times in saline so as to stop the activity of the bacterial enzyme.

If they have been sufficiently transformed, they will be found to undergo agglutination within five to 10 minutes at room temperature in AB serum on a slide. If the cell filtrate mixture has been incubated for too long, the erythrocytes may undergo spontaneous agglutination in saline. It is noteworthy that Friedenreich (1930) also observed spontaneous agglutination as the result of the transformation process being carried on too far, but, contrary to our findings, this happened only when he used bacteria instead of filtrate.

Titration of T Agglutinin.—To determine the T titre, doubling dilutions of the serum were made in test-tubes (7×60 mm.). Two drops of a 4% suspension of transformed cells were added to two drops of the serum dilution. The reactions were read microscopically after one hour at 16° C. and recorded as ++, +, (+), ±, or —.

Absorption of T Agglutinin.—The T antibody was absorbed from one volume of a given serum with two volumes of transformed packed cells. After one hour at 16° C. the cells were rapidly centrifuged and the supernatant titrated against T erythrocytes, as described above.

Elution of T Agglutinin.—The T antibody was eluted from the T cells used for the absorption test. For this purpose the cells were washed twice in cold saline and centrifuged. The supernatant fluid was pipetted off and replaced by two volumes of fresh saline. The mixture was shaken and kept for ten minutes at 45° C. in a water bath. After rapid centrifugation the supernatant fluid was again titrated with T erythrocytes.

Results

As can be seen from Table III, the agglutination reactions at 16° C. were stronger than at 22° C. and 37° C. The T agglutinin can thus be classified

TABLE III
AGGLUTINATION TITRES OF STREPTOCOCCUS-TREATED GROUP O RH-NEGATIVE RED CELLS WITH GROUP AB SERUM AT FOUR DIFFERENT TEMPERATURES

Temperature	1:1	1:2	1:4	1:8	1:16	1:32
4° C.	++	++	+	+	(+)	—
16° C.	++	++	+	+	±	—
22° C.	++	+	+	(+)	±	—
37° C.	+	+	—	—	—	—

among the specific cold agglutinins. As a working temperature, 16° C. is preferable to 4° C., since non-specific cold antibodies, not identical with T agglutinins, are active at 4° C. The amount of T agglutinin varies in different sera, titres varying from 1 in 4 to 1 in 64 (Table IV).

An experiment was carried out to see if red blood cells treated with the patient's streptococcus filtrate developed the same T agglutininogen as cells treated with cholera filtrate. Erythrocytes from the same normal subject were treated with cholera filtrate in

TABLE IV
AGGLUTINATION TITRES OF STREPTOCOCCUS- AND CHOLERA-TREATED GROUP O RH-NEGATIVE RED CELLS WITH NORMAL SERA AT 16° C.

Type of Cells	Serum No.								
	5	6	7	9	14	15	17	19	26
Streptococcus-filtrate-treated cells	1/8 (+)	1/16 ±	1/16 ±	1/32 ±	1/64 ±	1/16 ±	1/16 ±	1/4 (+)	1/4 ±
Cholera-filtrate-treated cells	1/32 ±	1/16 (+)	1/16 ±	1/32 (+)	1/64 ±	1/16 ±	1/16 (+)	1/16 ±	1/8 ±

a manner similar to that described for the streptococcus filtrate. One volume of packed red cells was mixed with one and a half volumes of cholera filtrate and the mixture allowed to stand for one hour. Two parallel series of serum dilutions were made; streptococcal-filtrate-treated cells were added to the first and cholera-filtrate-treated cells to the second (Table IV). The two types of cells behaved in almost the same manner.

It was next shown that streptococcal-filtrate-treated cells specifically adsorbed the T antibody from AB serum, with the result that this serum failed to react not only with fresh streptococcal-filtrate-treated cells, but also with cholera-filtrate-treated cells. Adsorption with cholera-filtrate-treated cells gave the same result. It was also shown that an eluate of T antibody adsorbed by streptococcus-treated cells agglutinated both types of cells and the same was true of an eluate derived from cholera-filtrate-treated cells (Table V). It is concluded, therefore, that both T agglutinogens are serologically identical.

Finally, it was found that Rh-positive (D+) erythrocytes treated with streptococcus filtrate were

TABLE V
CROSS-ABSORPTION TEST

	Serum (No. 15) Adsorbed with Streptococcus-treated Red Cells	Serum (No. 15) Adsorbed with Cholera-treated Red Cells
Streptococcus-treated cells ..	—	—
Cholera-treated cells ..	—	—

CROSS-ELUTION TEST

	1:1	1:2	1:4	1:8
Dilutions of Eluate from Streptococcus-treated Cells:				
Streptococcus-treated cells	+	(+)	±	—
Cholera-treated cells ..	+	(+)	±	—
Dilutions of Eluate from Cholera-treated Cells:				
Streptococcus-treated cells	+	+	±	—
Cholera-treated cells ..	+	+	±	—

agglutinated in saline by a serum containing incomplete Rh antibodies (anti-D) from which the T antibodies had been previously absorbed with T-positive Rh-negative cells (Table VI).

TABLE VI
SERUM A.N. CONTAINING INCOMPLETE ANTI-D TITRATED
AGAINST STREPTOCOCC L- AND CHOLERA-TREATED GROUP O
D POSITIVE CELLS IN SALINE AT 37° C.

Serum Dilution:	1:1	1:2	1:4	1:8	1:16	1:32	1:64	1:128
	Agglutination							
Streptococcus-treated group O D+ cells ..	+	+	+	(+)	±	±	±	-
Cholera-treated group O D+ cells ..	++	+	+	+	(+)	-	-	-
Control cells: streptococcus-treated D negative cells ..	-							

The analogous action of cholera filtrate was discovered by Pickles (1946). Chu and Coombs (1947) showed that *Vibrio proteus*, *Cl. welchii*, pneumococci, and some virus strains behaved in the same manner, and Hummel (1951) described this behaviour with a bacillus resembling Friedenreich's "M" bacillus, which he had isolated from old stored blood.

Discussion

The observations now reported support the hypothesis that polyagglutinability of erythrocytes *in vivo* and panagglutinability *in vitro* may be induced by the same causal factor. In nearly all the published reports of polyagglutinability, except for the second case of Gaffney and Sachs (1943) and the case of Henningsen (1949), a bacterial infection was demonstrated as being the possible cause. The possibility that the abnormality of the red cells and the accompanying infection are related seems especially likely in the cases with pyogenic sepsis (Basil-Jones, Sanger, and Walsh, 1946; Engleson and Grubb, 1949) and in the two pneumococcal infections (Levine and Katzin, 1938; Holländer, 1951), because Chu (1948) demonstrated that these organisms might cause panagglutinability *in vitro*. However, no attempts seem to have been made to test for the ability of the isolated micro-organism to render red cells panagglutinable. Similarly, no clear relationship was demonstrated by van Loghem, van der Hart, and Heier (1951) and Gasser and Holländer (1951). In the case of van Loghem *et al.* (1951) it is possible that enzymes with activity similar to those derived from certain bacteria were released from neoplastic tissue (Rosenthal and

Schwartz, 1951). On the other hand, Gram-positive diplococci cultivated from the sputum could have been the cause of the abnormal agglutinability in this case.

Several authors showed that the erythrocytes of their patients behaved as did erythrocytes transformed *in vitro* (Gaffney and Sachs, 1943; Boorman, Loutit, and Steabben, 1946; van Loghem *et al.*, 1951; Holländer, 1951). Henningsen (1949), however, found the agglutinin adsorbed by the cells of his blood No. 38 to be not quite identical with that adsorbed by actinomyces-treated cells.

It has been pointed out that the transient nature of polyagglutinability *in vivo* compared with the stability of the T antigen and the regularity with which cells may be transformed *in vitro* is against the hypothesis that the two phenomena are identical. However, the apparent discrepancy between these two characteristics may be explained by the fact that body-temperature is unfavourable for the action of the enzyme on red cells. It is possible also that the presence of an enzyme inhibitor in human serum may weaken or prevent the reaction. It is worth noting in this connexion that Stack and Morgan (1949) found that the serum of rabbits immunized with enzymes from *Cl. welchii* developed anti-enzyme properties. Finally, the transient character of polyagglutinability can be explained by the disappearance of the responsible enzyme with the elimination of the local infection. This was confirmed in our case.

It has often been stressed that *in vivo* the phenomenon is usually that of polyagglutinability, whilst *in vitro* the cells become panagglutinable. This is, however, not true of all the published cases (Table I), and certainly not true of those described by Engleson and Grubb (1949), Holländer (1951), and Gasser and Holländer (1951), where auto-agglutination took place.

It has also been stressed that it is not possible to transmit the polyagglutinability to fresh normal blood, as can be done with blood artificially transformed with bacteria. These failures, however, do not argue against the Hübener-Thomsen-Friedenreich phenomenon manifesting itself *in vivo*, because previous experiments have shown that sterile culture filtrates are able to produce this change. The author concludes that polyagglutinability of red cells *in vivo* is in most cases probably caused by infection with certain enzyme-producing bacteria. These bacteria need not necessarily be detectable by blood culture. The occurrence of the phenomenon in apparently healthy individuals remains unexplained.

Summary

A case is described of a patient, aged 25, with a chronic, relapsing urinary infection, whose erythrocytes showed the transient phenomenon of polyagglutinability *in vivo*. A short review of the literature is given, together with a list of the micro-organisms known to cause the Hübener-Thomsen-Friedenreich phenomenon *in vitro*.

An α -haemolytic streptococcus was isolated from the urine of the patient. This organism was found to be capable of rendering red cells panagglutinable. It appeared to belong to the same species that Gardborg and Borgen (1951) isolated from urine and which had similar properties.

The T agglutininogen that arises at the surface of normal red cells after treatment with a filtrate of this streptococcus was shown to be identical with the one resulting from treatment with cholera filtrate. It was demonstrated that incomplete antibodies (anti-D) could be detected in saline with Rh-positive red cells previously treated with the bacterial filtrate.

The relationship between polyagglutinability of erythrocytes taking place *in vivo* and the Hübener-Thomsen-Friedenreich phenomenon is discussed. It is concluded that these phenomena are intimately related and most probably identical.

This paper has been prepared with the technical assistance of Miss J. Willems and Miss M. van der Hart, whom I wish to thank.

I am indebted to Dr. F. Dekking for the cultivation of the bacteria and to Dr. K. Skadhauge and Dr. V. Friedenreich for investigating the properties of the streptococcus.

All the clinical data were kindly provided by Dr. P. G. Hoorweg, of the Binnengasthuis, Amsterdam.

I wish to thank Dr. J. J. van Loghem, Jr., for his kind help and valuable criticism.

REFERENCES

- Basil-Jones, B., Sanger, R. A., and Walsh, R. J. (1946). *Nature, Lond.*, **157**, 802.
- Boorman, K. E., Loutit, J. F., and Steabben, D. B. (1946). *Ibid.*, **158**, 446.
- Briody, B. A. Unpublished data.
- Burnet, F. M., McCrea, J. F., and Stone, J. D. (1946). *Brit. J. exp. Path.*, **27**, 228.
- Chu, C. M. (1948). *Nature, Lond.*, **161**, 606.
- and Coombs, R. R. A. (1947). *Lancet*, **1**, 484.
- Davidson, I., and Toharsky, B. (1940). *J. infect. Dis.*, **67**, 25.
- (1942). *J. Immunol.*, **43**, 213.
- Engleson, G., and Grubb, R. (1949). *Amer. J. clin. Path.*, **19**, 782.
- Friedenreich, V. Personal communication to the author.
- (1930). *The Thomsen Hemagglutination Phenomenon*. Copenhagen.
- Gaffney, J. C., and Sachs, H. (1943). *J. Path. Bact.*, **55**, 489.
- Gardborg, O., and Borgen, L. O. (1951). *Acta path. microbiol. scand.*, **28**, 116.
- Gasser, C., and Holländer, L. (1951). *Rev. Hémat.*, **6**, 316.
- Henningsen, K. (1949). *Acta path. microbiol. scand.*, **26**, 339.
- Holländer, L. (1951). *Helv. paediat. Acta*, **6**, 149.
- Hübener, G. (1925). *Z. Immunforsch.*, **45**, 223.
- Hummel, K. (1951). *Ibid.*, **108**, 233.
- Kossovitch, N., and Chabaud, A. (1938). *C.R. Soc. Biol., Paris*, **128**, 851.
- Levine, P., and Katzin, E. M. (1938). *Proc. Soc. exp. Biol., N.Y.*, **39**, 167.
- Loghem, J. J. van, Hart, M. van der, and Heier, A. M. (1951). *Bull. centr. Lab. Bloedtr. Ned. Rode Kruis*, No. 2, 93.
- Magnus, R. von (1936). Thesis, *Undersøgelser over en Gruppe Actinomyceiter isolerede fra menneskets Svalg* (with English summary). Copenhagen.
- Moskowitz, M., and Treffers, H. P. (1950). *Science*, **111**, 717.
- Pickles, M. M. (1946). *Nature, Lond.*, **158**, 880.
- Rosenthal, M. C., and Schwartz, L. I. (1951). *Proc. Soc. exp. Biol., N.Y.*, **76**, 635.
- Stack, M. V., and Morgan, W. T. J. (1949). *Brit. J. exp. Path.*, **30**, 470.
- Stewart, F. S. (1949). *J. Path. Bact.*, **61**, 456.
- Stone, J. D. (1947). *Aust. J. exp. Biol. med. Sci.*, **25**, 137.
- Terada, S. (1936). Thesis, *J. med. Ass. Formosa*, **35**, 1285. Cited by Davidson, I., and Toharsky, B. (1940).
- Thomsen, O. (1927a). *Z. Immunforsch.*, **52**, 85.
- (1927b). *C.R. Soc. Biol., Paris*, **96**, 556.
- (1928). *Z. Immunforsch.*, **57**, 301.