Technical report

Use of alcohol fixed cytopspins protected by 10% polyethylene glycol in immunocytology external quality assurance

P Maxwell, A H Patterson, J Jamison, K Miller, N Anderson

Abstract

Aims—To provide cytopspins as a means of external quality assurance (EQA), while maintaining antigen expression integrity and achieving uniformity of material for all participating laboratories.

Methods—Cells were collected from two adenocarcinoma and one reactive pleural effusion specimens. Lymphoid cells were also collected through aspiration of a resected tonsil specimen. All cells were collected in Hanks balanced salt solution (HBSS); cytopspins were made and fixed in methanol or acetone alone or protected using a layer of 10% polyethylene glycol (PEG) in 50% methanol. Two laboratories participated (RGHT and UCL).

Results—Cytokeratin expression detected using either CAM5.2 or AE1/AE3 antibodies was sensitive to temperature. Without PEG, unacceptable or negative staining was seen within six weeks of preparation. LCA was not sensitive to temperature, with good staining scores being achieved after eight weeks following preparation.

Conclusions—It is possible to send alcohol fixed, air dried cytopspins to laboratories participating as part of an EQA scheme for immunocytology. Some antigens will require protection from temperature variations during transit. A layer of 10% PEG in 50% methanol, allowed to air dry, is suitable for this purpose. Participating laboratories will only have to remove this layer using methanol before their localisation technique for assessment.

(J Clin Pathol 1999;52:141-144)

Keywords: immunocytochemistry; cytopathology; external quality assurance

Immunocytology on cytopspins or smears has, to date, been used sparingly. Tissue sections made from cell blocks in a histopathology laboratory are often used in cytopathology for immunostaining. However, the increased use of cytological material for diagnostic purposes and the scarcity of cells in some specimens means that there is a need for this method of immunological localisation.¹⁻³ With this need

comes the requirement for the laboratory to participate in an external quality assurance scheme. In this study, we describe a means by which cytopspins may be used by such schemes to assess the localisation of antigens.

Methods

STORAGE AND TRANSPORT OF CYTOSPINS

The effect of storage at room temperature on antigen expression was tested by preparing cells collected into Hank's balanced salt solution (HBSS, Life Technologies). Cytopspins were prepared on a Shandon Cytospin 2 and fixed in either methanol or acetone for 20 minutes. Cells were stained immediately and at weekly intervals over a period of six to eight weeks after storing at room temperature.

The effect of protecting the antigen from degradation was assessed by comparison with cytopspins prepared at the same time as those described above and coated with 10% polyethylene glycol (PEG) in 50% methanol, air dried, and stained immediately and at weekly intervals for six weeks.

The remaining cells from each specimen were placed in Carboxifix (Shandon), a fixative/transport medium. Cytopspins were prepared at

Table 1 Assessment of cytokeratin localisation using CAM 5.2 antibody. Pleural effusion, first specimen of adenocarcinoma stained by both Royal Group of Hospitals Trust (RGHT) and University College, London (UCL)

	Week 1	Week 2	Week 3	Week 4	Week 5	Week 6
Acetone						
RGHT	6	7	8	ND	ND	ND
UCL	ND	ND	11	6	4	5
Methanol						
RGHT	7	4	6	ND	ND	ND
UCL	ND	ND	8	5	6	8
Carboxifix, acetone						
RGHT	17	9	4	9	5	7
UCL	ND	ND	11	12	13	12
Carboxifix, methanol						
RGHT	16	13	10	11	4	9
UCL	ND	ND	13	12	13	12
Acetone, PEG						
RGHT	15	17	16	17	17	16
UCL	ND	ND	ND	ND	16	16
Methanol, PEG						
RGHT	14	17	16	16	16	16
UCL	ND	ND	ND	ND	15	16

ND, not done.

Institute of Pathology,
Royal Group of
Hospitals Trust,
Grosvenor Road,
Belfast BT12 6BA, UK
P Maxwell
A H Patterson
N Anderson

Department of
Cytopathology, United
Hospitals Trust, Bush
Road, Antrim, UK
J Jamison

UKNEQAS for
Immunocytochemistry,
Department of
Histopathology,
University College
Medical School,
University Street,
London WC1E 6JJ, UK
K Miller

Correspondence to:
Dr P Maxwell.
email: p.maxwell@qub.ac.uk

Accepted for publication
11 August 1998

Table 2 Assessment of cytokeratin localisation using CAM 5.2 and AE1/AE3 antibodies. Pleural effusion, second specimen of adenocarcinoma, stained by the Royal Group of Hospitals Trust only

	Week 1	Week 2	Week 3	Week 4	Week 5	Week 6
Acetone						
CAM 5.2	15	16	15	12	15	14
AE1/AE3	15	16	10	8	12	12
Methanol						
CAM 5.2	11	17	16	15	14	14
AE1/AE3	18	13	12	12	11	12
Acetone, PEG						
CAM 5.2	19	19	15	15	16	17
AE1/AE3	16	16	12	12	12	9
Methanol, PEG						
CAM 5.2	13	15	15	19	19	20
AE1/AE3	19	19	19	9	16	16

Table 3 Assessment of cytokeratin localisation using CAM 5.2 and AE1/AE3 antibodies. Pleural effusion, diagnosed as reactive

	Week 1	Week 2	Week 3	Week 4	Week 5	Week 6
Acetone						
CAM 5.2	11	11	13	12	12	12
AE1/AE3	10	8	8	8	8	9
Methanol						
CAM 5.2	13	12	13	11	12	10
AE1/AE3	10	10	9	9	9	9
Acetone, PEG						
CAM 5.2	8	10	8	7	7	8
AE1/AE3	8	9	8	7	7	6
Methanol, PEG						
CAM 5.2	17	15	19	18	19	17
AE1/AE3	19	20	20	19	18	ND

weekly intervals, fixed in either acetone or methanol, and treated as for the protected cytopins before staining as described below.

IMMUNOCYTOCHEMISTRY AND ITS ASSESSMENT

Anti-leucocyte common antigen (LCA, 1:50, Dako) and anti-cytokeratin (CAM 5.2, 1:20, Becton-Dickinson; AE1/AE3, 1:50, Dako) were localised using a Duet kit, peroxidase streptavidin-biotin complex method (StABC, Dako). Diaminobenzidine acted as the chromogen and cells were counterstained in Harris's haematoxylin. Endogenous peroxidase was blocked using 3% alcoholic hydrogen peroxide. All cytopins either prepared from cells in Carboxif or protected by 10% PEG in 50% methanol, were incubated in two baths of 95% alcohol for five minutes before staining. Results were confirmed at a second centre (UCL, University College Medical School, London) using a standard alkaline phosphatase anti-alkaline phosphatase method (APAAP, Dako).

The staining was assessed by a panel of four assessors, one of whom was a consultant cytopathologist (NA). Each assessor independently scored the slides using a 5 point scale. Two marks were available for antigen demonstration. Once antigen demonstration was seen, two marks were available for clarity of background or lack of non-specific staining. One mark was available for the quality of the counterstain. Collectively, therefore, each slide was scored out of 20. Acceptable staining was recorded when scoring exceeded 12 points.

Figure 1 Pleural effusion, adenocarcinoma stained for low molecular weight cytokeratin (CAM 5.2). Methanol fixation after (A) one week and (B) six weeks. Methanol fixation with a layer of PEG after (C) one week and (D) six weeks.

Results

Both pleural specimens of adenocarcinoma showed an unacceptably low level of cytokeratin expression after six weeks when fixed in either methanol or acetone and compared with the results of staining immediately after

Figure 2 Aspirated tonsil cells stained for anti-leucocyte common antigen (LCA). Methanol fixation after (A) one week and (B) eight weeks.

Table 4 Assessment of LCA localisation. Cells collected from specimen of resected tonsil and stained by both Royal Group of Hospitals Trust (RGHT) and University College, London (UCL)

	Week 1	Week 2	Week 3	Week 4	Week 5	Week 6	Week 7	Week 8
Acetone								
RGHT	13	13	12	15	12	12	15	16
UCL	ND	ND	ND	ND	12	14	13	16
Methanol								
RGHT	16	16	13	15	16	16	15	16
UCL	ND	ND	ND	ND	13	15	13	12
Carboxif, acetone								
RGHT	16	15	16	15	17	16	13	ND
UCL	ND	ND	ND	ND	15	14	15	13
Carboxif, methanol								
RGHT	17	16	15	16	15	16	14	ND
UCL	ND	ND	ND	ND	15	14	15	14

preparation. When protected using PEG, however, cytokeratin expression was maintained and demonstration was scored as being excellent (tables 1–3). During week 1, higher scores for cytokeratin expression were noted in one case of adenocarcinoma and in reactive cells when placed in Carboxif or when PEG was used as a protective layer.

Overall, there was a similar reaction to all handling protocols when cytokeratin expression was demonstrated using either CAM5.2 (low molecular weight cytokeratin, fig 1) or a combined AE1/AE3 (pan-cytokeratin) antibody.

The demonstration of LCA in tonsil cells was unaffected by storage at room temperature after eight weeks (fig 2). There was also little difference in expression when fixed in acetone, methanol or when cells were placed in Carboxif (table 4).

The results were confirmed by the second laboratory (UCL) which received cytopsins made from the tonsil and one adenocarcinoma pleural specimen by post at week 3 of the study.

Discussion

Due to minimum invasiveness of the procedure and the potential speed of diagnosis, techniques such as fine needle aspiration (FNA) cytology are becoming increasingly popular in diagnostic pathology.¹ Provided adequate cells are available, it is possible to process the sample to paraffin wax to make a cell block which may be used for immunocytochemistry.^{2 3} Small numbers of cells, however, often mean that cytopsin preparations are all that are available for immunocytochemistry. The maximum amount of information therefore needs to be obtained from a minimum number of slides. Immunocytochemistry has become a standard procedure in most histopathology laboratories, but to date it has been used sparingly in cytopathology laboratories. The nature of the cytology material, however, does not exclude the use of immunocytochemistry and the slides are often passed on to the histopathology laboratory to stain cells obtained on sampling.

The need for a controlled immunocytochemical procedure on these samples is evident. Recently, 12 laboratories in The Netherlands participated in a limited external quality control study on material obtained from effusions.⁴ In this study, laboratories were sent cells to prepare their own cytopsins or unstained control specimens. In our experience, sending cells through the post to participating laboratories is restricted by Health and Safety practices and postal regulations.

The present study has shown that sending air dried cytopsins in the post requires fixation and a layer of protection in the form of PEG. The effect of degrading antigen expression was dependent upon the antigen. LCA appears to require no protection, whereas cytokeratin expression, irrespective of molecular weight, seems to be degraded at room temperature. Interestingly, methanol fixation gave consistently better results than acetone fixation, perhaps because of the removal of PEG by the methanol. This wash in methanol may also have a secondary fixation effect, contributing towards an enhanced staining reaction.

We have found, therefore, that it is possible to send alcohol fixed, air dried cytopsins in the post to other laboratories and expect the material to be robust enough to resist reasonable temperature changes. Other applications, such as using the procedure in storing cytopsins at room temperature for control material, can also be considered. For cytokeratin expression, the cytopsins require a further layer of protection. The application of 10% PEG in 50% methanol, appears to protect the cytokeratin from degradation.

This material should be suitable for use in EQA schemes in immunocytology on cytological material. Some laboratories already perform immunocytochemistry either on cytopsins or cell blocks.^{2 3} In the latter instance,

these laboratories participating in EQA schemes for immunohistochemistry still require adequate monitoring. For those cytopathology laboratories performing occasional immunocytochemistry, the proposed scheme is the first to offer cytopins which should be suitable for the production of uniform demonstration of antigen. This means that all participating laboratories will be able to be assessed equally.

- 1 Hanson CA. Fine-needle aspiration and immunophenotyping. A role in diagnostic hematopathology? *Am J Clin Pathol* 1994;**101**:555–6.
- 2 Zito FA, Gadaleta CD, Salvatore CT, et al. A modified cell block technique for fine needle aspiration cytology. *Acta Cytol* 1995;**39**:93–9.
- 3 Mayall F, Chang B, Darlington A. A review of 50 consecutive cytology block preparations in a large general hospital. *J Clin Pathol* 1997;**50**:985–90.
- 4 Kounen-Boumeester V, van Loenen P, de Bruijn EMCA, et al. Quality control of Immunocytochemical staining of effusions using a standardized method of cell processing. *Acta Cytol* 1996;**40**:475–9.

The Royal College of Pathologists “The Best” of the College Symposia

Academic activities, Continuing Professional Development, and Blackwell Healthcare Communications are publishing a CD-ROM containing highlights of past College Symposia.

This CD-ROM is approved for CPD credits and is of general interest to pathologists. It covers a wide range of topical issues, with lectures entitled:

- Allergy and asthma
- Peanut allergy
- A practical approach to unexpected infant death
- Vitamin K prophylaxis: can we ever reach a consensus?
- Epidemiology of adverse effects of “designer drugs”
- Multi-drug resistance modifiers: an alternative approach
- Idiopathic myelofibrosis: pathogenesis to treatment
- Adhesion molecules in pathology
- Chromosomal changes and cancer
- Interpretation of injury in road traffic accidents
- Reporting of cervical biopsies in the context of the cervical screening programmes
- The metastatic process: its biological basis
- Human papilloma virus infection
- The hepatitis virus
- Alcoholic liver disease

The CD-ROM will cost you £30 for a personal copy or £120 for an institutional copy with multiple user licence. To order a copy, call 0171 930 5862.