SHORT REPORT

Measurement of 5-hydroxy-2-aminovaleric acid as a specific marker of metal catalysed oxidation of proline and arginine residues of low density lipoprotein apolipoprotein B-100 in human atherosclerotic lesions

J Pietzsch, R Bergmann

J Clin Pathol 2003;56:622–623

γ -Glutamyl-semialdehyde (γ GSA) is a major product of the metal catalysed oxidation of apolipoprotein B-100 (apoB-100) proline and arginine residues. On reduction, γ GSA forms 5-hydroxy-2-aminovaleric acid (HAVA). This report describes the application of HAVA measurement to characterise the formation of γ GSA in low density lipoprotein (LDL) recovered from human atherosclerotic lesions. HAVA concentrations were greatly increased in LDL from early (mean, 10.25; SD, 3.49 mol/mol apoB-100; $p < 0.01$), intermediate (mean, 11.18; SD, 2.37 mol/mol apoB-100; $p < 0.01$), and advanced (mean, 9.91; SD, 2.15 mol/mol apoB-100; $p < 0.01$) lesions, when compared with LDL from normal aortic tissue (mean, 0.05; SD, 0.01 mol/mol apoB-100). These findings support the hypothesis that pathways involving metal catalysed oxidation of LDL apoB-100 are of pathological importance in atherogenesis.

Oxidative modification of low density lipoprotein (LDL) apolipoprotein B-100 (apoB-100) is regarded as a crucial event in atherogenesis.¹ Direct oxidation of apoB-100 amino acid residues is thought to result in the formation of new epitopes that are specifically recognised by scavenger receptors.¹ However, the nature of such new epitopes is still a matter of debate. One potential pathway for direct oxidation of apoB-100 involves myeloperoxidase, which finally generates specific oxidation products, such as 3-chlorotyrosine and 3-nitrotyrosine.² All of these compounds have been found in LDL apoB-100 recovered from human atherosclerotic vascular tissue.^{2,3} There is evidence supporting the hypothesis that metal catalysed processes are important mechanisms of protein oxidation.⁴ These processes involve binding of either free or, physiologically more relevant, complexed (porphyrin bound) redox active iron ($\text{Fe}^{2+}/\text{Fe}^{3+}$) to discrete binding sites of LDL and apoB-100, respectively, thus forming centres for redox cycling and repeated radical production.^{4,5}

"Oxidative modification of low density lipoprotein apolipoprotein B-100 is regarded as a crucial event in atherogenesis"

A major product of transition metal catalysed protein oxidation is γ -glutamyl-semialdehyde (γ GSA).⁶ Recently, oxidation of LDL apoB-100 proline and arginine residues to γ GSA, which by reduction forms 5-hydroxy-2-aminovaleric acid (HAVA), has been measured in vitro and in circulating LDL in vivo.^{7,8} In our present study, we used specific gas chromatography mass spectrometry (GC-MS) methodology to

demonstrate that concentrations of HAVA are greatly raised in LDL apoB-100 recovered from human aortic vascular lesions. These observations provide quantitative chemical evidence for metal catalysed oxidative processes in the human artery wall.

METHODS AND MATERIALS

Lesional LDL was isolated from the intima of normal and atherosclerotic specimens of human thoracic aortas obtained at necropsy within 10 hours of sudden death (20 male accident victims, aged 26 to 45 years; no signs of severe acute diseases). Normal and atherosclerotic tissue (fatty streaks, intermediate, and advanced plaques) was classified according to the criteria of the pathobiological determinants of atherosclerosis in youth study.⁹ Samples were processed in subdued light to prevent photooxidation. Aortic intima was dissected from the medium and homogenised for two hours at 4°C (30 mg wet tissue/ml phosphate buffered saline, pH 7.2). All buffers and solutions were degassed and stored under argon. Furthermore, all buffers and solutions were supplemented with 100 μM diethylenetriaminepentaacetic acid, 100 μM butylated hydroxyanisole, and 100 μM 3-amino-1,2,4-triazole. Lesional LDL (density, 1.006–1.063 g/ml) was then isolated by very fast ultracentrifugation at 120 000 revolutions/minute, corresponding to 625 000 $\times g$, with a run time of 100 minutes at 18°C.¹⁰ Delipidation of LDL, formation of HAVA by reduction of γ GSA with sodium borohydride, and enzymatic hydrolysis of apoB-100 were performed as described previously.⁷ The free amino acids from hydrolysates were derivatised to their *N*(*O*)-ethoxycarbonyl ethyl ester derivatives, and analysed by GC-MS, as described elsewhere.⁷ The HAVA content in all samples is expressed as mol/mol apoB-100 (intra-assay coefficient of variation (CV), < 4.5%; interassay CV, < 6.1%).

RESULTS AND DISCUSSION

Figure 1 shows the HAVA content of intima from various stages of lesion evolution as determined by sensitive and specific GC-MS analysis. The HAVA content was significantly higher in LDL recovered from all types of lesions when compared with normal aortic tissue (paired Student's *t* test). The overall yield of HAVA that has been found in lesional LDL (mean HAVA content, 10.45; SD, 0.66 mol/mol apoB-100; 328/10 000 proline plus arginine residues) is remarkably high, and indicates that proline and arginine residues are good targets

Abbreviations: apo, apolipoprotein; CV, coefficient of variation; GC-MS, gas chromatography mass spectrometry; γ GSA, γ -glutamyl-semialdehyde; HAVA, 5-hydroxy-2-aminovaleric acid; LDL, low density lipoprotein

Figure 1 Box plots showing low density lipoprotein apolipoprotein B-100 (apoB-100) 5-hydroxy-2-aminovaleric acid (HAVA) content in intima from various stages of lesion evolution in human thoracic aorta.

for metal catalysed oxidative attack in atherosclerotic tissue. Control experiments indicated that sample processing was unlikely to have contributed to HAVA formation as an artifact (data not shown in detail). The HAVA content of both atherosclerotic and normal aortic tissue exceeded the physiological concentrations of HAVA formerly found in native plasma LDL obtained from healthy normolipidaemic subjects (mean, 0.012; SD, 0.004 mol/mol apoB-100; 0.4/10 000 proline plus arginine residues).⁷ Furthermore, the findings are consistent with former data showing increased HAVA content (mean, 0.063; SD, 0.02 mol/mol apoB-100; 2.0/10 000 proline plus arginine residues) in circulating LDL recovered from hypercholesterolaemic subjects who were at high atherosclerotic risk.⁸ The experiments support the hypothesis that a pathway involving metal catalysed oxidation of LDL apoB-100 may be of pathological importance in atherogenesis. However, additional work is needed to understand the specific consequences of γ GSA formation for the metabolic fate of apoB containing lipoproteins in vivo.

ACKNOWLEDGEMENT

We thank the staff of the Institute of Legal Medicine, Medical Faculty, Technical University Dresden for expert technical assistance.

Take home messages

- 5-Hydroxy-2-aminovaleric acid concentrations were greatly increased in low density lipoprotein (LDL) from early, intermediate, and advanced atherosclerotic lesions, when compared with LDL from normal aortic tissue
- These findings support the hypothesis that pathways involving metal catalysed oxidation of LDL apolipoprotein B-100 are of pathological relevance for atherogenesis

Authors' affiliations

J Pietzsch, R Bergmann, Institute of Bioinorganic and Radiopharmaceutical Chemistry, Research Center Rossendorf, PO Box 510119, D-01314 Dresden, Germany

Correspondence to: Dr J Pietzsch, Institute of Bioinorganic and Radiopharmaceutical Chemistry, Research Center Rossendorf, PO Box 510119, D-01314 Dresden, Germany; j.pietzsch@fz-rossendorf.de

Accepted for publication 18 March 2003

REFERENCES

- 1 Ross R. Atherosclerosis—an inflammatory disease. *N Engl J Med* 1999;**340**:115–26.
- 2 Podrez EA, Abu-Soud HM, Hazen SL. Myeloperoxidase-generated oxidants and atherosclerosis. *Free Radic Biol Med* 2000;**28**:1717–25.
- 3 Hazen SL, Heinecke JW. 3-Chlorotyrosine, a specific marker of myeloperoxidase-catalyzed oxidation, is markedly elevated in low density lipoprotein isolated from human atherosclerotic intima. *J Clin Invest* 1997;**99**:2075–81.
- 4 Fu S, Davies MJ, Stocker R, *et al*. Evidence for roles of radicals in protein oxidation in advanced human atherosclerotic plaque. *Biochem J* 1998;**333**:519–25.
- 5 Miller YI, Shaklai N. Kinetics of hemin distribution in plasma reveals its role in lipoprotein oxidation. *Biochim Biophys Acta* 1999;**1454**:153–64.
- 6 Requena JR, Chao CC, Levine RL, *et al*. Glutamic and aminoaliphatic semialdehydes are the main carbonyl products of metal-catalyzed oxidation of proteins. *Proc Natl Acad Sci U S A* 2001;**98**:69–74.
- 7 Pietzsch J. Measurement of 5-hydroxy-2-aminovaleric acid as a specific marker of iron-mediated oxidation of proline and arginine side chain residues of low density lipoprotein apolipoprotein B-100. *Biochem Biophys Res Commun* 2000;**270**:852–57.
- 8 Pietzsch J, Latke P, Julius U. Oxidation of apolipoprotein B-100 in circulating low density lipoproteins is related to low density lipoprotein residence time: in vivo insights from stable isotope studies. *Arterioscler Thromb Vasc Biol* 2000;**20**:E63–7.
- 9 PDAY Research Group. Natural history of aortic and coronary atherosclerotic lesions in youth: findings from the PDAY study. *Arterioscler Thromb* 1993;**13**:1291–8.
- 10 Pietzsch J, Subat S, Nitzsche S, *et al*. Very fast ultracentrifugation of serum lipoproteins: influence on lipoprotein separation and composition. *Biochim Biophys Acta* 1995;**1254**:77–88.