REVIEW

The eicosanoid cascade: possible role in gliomas and meningiomas

N Nathoo, G H Barnett, M Golubic

J Clin Pathol: Mol Pathol 2004;**57**:6–13

Eicosanoids constitute a large family of biologically active lipid mediators that are produced by two enzyme classes, cyclooxygenases (COX-1 and COX-2) and lipoxygenases (5-LO, 12-LO, and 15-LO). Increasing evidence suggests that in addition to a variety of epithelial malignancies, the two most common types of human brain tumour, gliomas and meningiomas, aberrantly overexpress eicosanoid producing enzymes and release a spectrum of eicosanoids that may promote tumorigenesis and the development of peritumorous brain oedema. Glioma and meningioma cells are killed in vitro and in animal models when exposed to COX-2 and 5-LO inhibitors, and their effectiveness is under investigation in clinical trials for treatment of patients with malignant brain tumours. However, despite research into the role of the eicosanoid cascade in the tumorigenesis of human brain tumours, many important questions remain unanswered. Current and newer agents that specifically target key players of the eicosanoid cascade could change the approach to treating brain tumours, because their benefits may lie in their synergism with conventional cytotoxic treatments and/or with other novel agents targeted against other procarcinogenic pathways.

die from their disease in less than a year.⁴ In contrast, meningiomas are usually benign tumours that develop from the leptomeningeal cells of the arachnoid membrane that surrounds the brain and spinal cord.⁵ However, they often recur after seemingly complete surgical removal and occasionally progress to a fully malignant phenotype.⁶ Moreover, surgical cure is often not possible in those meningiomas involving the skull base or vital neurovascular structures.

In the past two decades, a large number of molecular and genetic pathways that are altered in brain tumour cells have been identified.⁴ However, the development of novel treatment strategies based on this molecular information has not yet been successfully translated to common clinical practice. A variety of promising drugs that selectively target precise molecular abnormalities, which create and drive the neoplastic phenotype of glioma and meningioma cells, are currently being investigated in clinical trials to improve the survival and quality of life of patients with brain tumours.

"In the past two decades, a large number of molecular and genetic pathways that are altered in brain tumour cells have been identified"

In this review, we will discuss the eicosanoid cascade of arachidonic acid (AA) and its relevance to brain tumours, the current evidence for the rationale of using inhibitors of eicosanoid producing enzymes for therapeutic purposes, and the limitations that hinder the full implementation of this form of treatment; in addition, we will suggest future directions for research and clinical practice.

INCREASED CONCENTRATIONS OF AA IN BRAIN TUMOURS

AA is a ω -6 polyunsaturated fatty acid (PUFA) that is converted into biologically active lipid compounds called eicosanoids. Eicosanoids constitute a large family of biologically active lipid mediators that are produced by two enzyme classes, cyclooxygenases (COX-1 and COX-2) and lipoxygenases (5-LO, 12-LO, and 15-LO).^{7–11}

Primary brain tumours are among the top 10 causes of cancer related death.¹ Approximately 13 000 people die from these tumours each year in the USA.¹ Gliomas and other neuroepithelial tumours make up 49% of primary brain tumours, and meningiomas are the second most common primary brain tumour (27%).²

The World Health Organisation (WHO) classifies gliomas as astrocytomas, oligodendrogliomas, and oligoastrocytomas on the basis of the phenotypical resemblance of tumour cells to normal glia (astrocytes or oligodendrocytes, or a combination of both). Astrocytomas are histologically graded as low grade (WHO I and II) and high grade tumours that are classified as anaplastic astrocytoma (WHO III) and glioblastoma multiforme (GBM) (WHO IV), respectively.³ The infiltrative growth pattern of high grade gliomas precludes them from being cured by neurosurgery and they respond poorly to other forms of treatment. The median survival of patients with a newly diagnosed GBM has improved only marginally during the past two decades, and today most patients with GBM still

See end of article for authors' affiliations

Correspondence to: Dr N Nathoo, Department of Neurosurgery, Cleveland Clinic Foundation, 9500 Euclid Avenue, Cleveland, 44195, Ohio, USA; nathoon@ccf.org

Accepted for publication 20 August 2003

These lipid compounds modulate diverse physiological and pathological processes including carcinogenesis.⁷⁻⁸ Several eicosanoids have been shown to promote cell survival,¹²⁻¹⁴ stimulate cell proliferation,¹⁵⁻¹⁶ and modulate cell adhesion and motility,¹⁴⁻¹⁶⁻¹⁷ angiogenesis,¹²⁻¹⁸ increased vascular permeability,¹⁹⁻²⁰ and inflammation,⁷⁻⁸ thus playing an important role in tumour growth and promotion (fig 1). In the past few years, numerous reports have indicated that not only a variety of epithelial human malignancies, but also glioma and meningioma cells, aberrantly overexpress eicosanoid producing enzymes, release a spectrum of eicosanoids, and are killed when exposed to specific eicosanoid inhibitors.

The importance of AA metabolism in human gliomas and meningiomas has been emphasised by reports demonstrating raised concentrations of ω -6 PUFAs, AA, and its precursor, linoleic acid (LA) in these tumours compared with normal white or grey matter.³⁴⁻³⁶ In contrast, concentrations of ω -3 PUFA docosahexaenoic acid were reduced in glioma samples compared with normal brain samples.³⁵⁻³⁶ These data suggested that the fatty acid composition of human brain tumours is different from that found in non-tumorous brain tissue, with ω -6 PUFAs, which may promote tumorigenesis in brain tumours, being expressed preferentially.

The causes and molecular mechanisms leading to the raised concentrations of LA and AA in gliomas and meningiomas are presently unknown. LA is an essential fatty acid that humans are unable to biosynthesise and is obtained exclusively through dietary intake.³⁷ Therefore, it is possible that raised concentrations of LA and AA in brain tumours may be related to high dietary consumption. Even though dietary LA (abundant in oils from various seeds) is the precursor of AA in tissues, dietary AA (exclusive to foods of animal origin) is perhaps the single most important nutritional determinant regulating AA values in humans.³⁸⁻³⁹ Furthermore, Nariai *et al* demonstrated in a rat tumour model using Walker 256 carcinosarcoma cells that nutritionally relevant fatty acids, including AA, were more readily incorporated into brain tumour tissue than normal brain when administered intravenously.⁴⁰

Epidemiological studies on dietary fat intake and the risk of brain tumours are sparse and not consistent.² Giles *et al* reported an increased risk of glioma in adult Australians with a high intake of oils, margarine, and salad dressing.⁴¹ Lubin *et al* reported that children with brain tumours had a higher consumption of vegetable fat than did controls.⁴² However, no significant association between the risk of glioma and intake of dietary fat was reported in a recent study from

Figure 1 Arachidonic acid (AA) metabolism. Brain tumour cells contain abnormally high concentrations of two dietary polyunsaturated fatty acids, AA and linoleic acid. AA is biosynthesised from dietary linoleic acid or is incorporated into membrane phospholipids directly from dietary AA. Activation of phospholipase A₂ (PLA₂) causes the release of free AA.²¹ Once AA is released from phospholipids, it is oxidised by one of three different groups of oxygenases into AA metabolites eicosanoids. The insertion of a single oxygen atom into AA by certain cytochrome P-450 enzymes produces epoxy-AA. Cyclooxygenases (COXs) introduce two molecules of oxygen into AA, and prostaglandins, thromboxanes, prostacyclin, and malondialdehyde are synthesised through a series of subsequent biochemical steps. COX-1 is a constitutively active enzyme present in most cell types, whereas COX-2 is almost undetectable in most unstimulated tissues.⁹⁻¹⁰ However, the expression and activity of COX-2 is inducible and regulated by a variety of ligands, including growth factors, Ras proteins, cytokines, and tumour promoters.¹⁰ Lipoxygenases (LOs) introduce one molecule of oxygen into AA to produce a series of isomeric hydroperoxy acids that are subsequently converted into leukotrienes and lipoxins. According to the positional specificity with respect to AA, mammalian lipoxygenases are categorised into three main groups, 5-LO, 12-LO, and 15-LO. 5-LO performs the first of two enzymatic reactions in the biosynthetic pathway for the leukotrienes.¹¹ COX and 5-LO derived eicosanoids stimulate cell proliferation, survival, adhesion, and motility, in addition to angiogenesis, increased vascular permeability, and inflammation, thus playing important roles in tumour growth and promotion.¹²⁻²⁰ They are also considered to be the main mediators of peritumorous brain oedema in patients with glioblastoma multiforme.¹⁹⁻²⁰ Tumorigenesis associated cellular functions of eicosanoids can be inhibited by the suppression of PLA₂ activity,²¹⁻²³ by cyclooxygenase inhibitors such as non-steroidal anti-inflammatory drugs (NSAIDs) and COX-2 specific inhibitors,¹⁰⁻²⁴⁻²⁵ by 5-LO inhibitors such as boswellic acids,²⁶⁻²⁹ and possibly by the reduced dietary intake of ω -6 linoleic and arachidonic acid and increased intake of ω -3 polyunsaturated fatty acids.³⁰⁻³³

Nebraska,⁴³ whereas increased consumption of total fat and cholesterol was found to be inversely associated with glioma risk in Israel.⁴⁴

Even if people who develop gliomas and meningiomas do not have a higher intake of AA and LA, it is possible that molecular changes in these tumours may lead to more efficient accumulation of these PUFAs in the tumour tissue in the presence of a normal dietary intake. For example, increased expression of fatty acid-CoA ligase 4, an enzyme that uses AA preferentially as a substrate for esterification into membrane phospholipids, was found in adenocarcinomas of the colon compared with normal tissue.⁴⁵ However, it remains to be determined whether similar molecular changes occur in brain tumours. Another reason why brain tumours contain raised concentrations of PUFAs may be that these tumours do not contain enzymes that would oxidise AA into eicosanoids. However, as described below, a large body of evidence does not support such an idea.

“Phospholipase A₂ is the key enzyme involved in the initiation of the entire eicosanoid cascade because it releases free arachidonic acid from the *sn*-2 position of membrane phospholipids”

Although ω -6 fatty acids are abundant in the membrane phospholipids of brain tumour cells, only free unesterified fatty acids that are released from the membrane phospholipids can be metabolised into eicosanoids. Intracellular concentrations of free AA are low, so that its formation limits the synthesis of eicosanoids. Phospholipase A₂ (PLA₂) is the key enzyme involved in the initiation of the entire eicosanoid cascade because it releases free AA from the *sn*-2 position of membrane phospholipids. Although mammalian cells normally contain more than 12 members of the PLA₂ group of enzymes,²¹ very limited information exists about PLA₂ enzymes in human brain tumours. The expression and enzymatic activity of the calcium sensitive cytosolic PLA₂ has been reported in normal human astrocytes, human astrocytoma cell lines 1321N1 and UC-11 MG, but not in U-373 MG cells.^{46, 47} Low amounts of the secreted form of PLA₂ and low enzymic activity were detected in normal rat astrocyte cell cultures and were dramatically induced by proinflammatory stimuli such as tumour necrosis factor α .⁴⁸ To our knowledge, the expression of other types of PLA₂ enzymes has not been reported in gliomas and meningiomas. Inhibition of PLA₂ by non-specific PLA₂ agents (4-bromophenacylbromide and quinacrine) suppressed the proliferation of primary meningioma cell cultures and the established human astrocytoma cell line, U373 MG, and these effects were partially reversed by the addition of exogenous AA.⁴⁹ These findings suggest that PLA₂ activity may be important for glioma and meningioma cell proliferation.

INCREASED EXPRESSION OF EICOSANOID PRODUCING ENZYMES IN GBM AND MENINGIOMA Cyclooxygenase

The crucial role of COX eicosanoids has been clearly demonstrated in human colon cancer, in which COX-2 is often overexpressed. COX inhibitors, such as non-steroidal anti-inflammatory drugs and COX-2 specific inhibitors, can reduce the incidence of colorectal cancer in humans and experimental animals, and can decrease the polyp number and size in patients with familial adenomatous polyposis.⁵⁰ In addition to colon cancer, COX-2 is overexpressed in a spectrum of human malignancies, such as cancer of the breast,^{51, 52} prostate,⁵³ pancreas,^{54, 55} oesophagus,⁵⁶ skin,^{57, 58} and lung.⁵⁹

Recently, gliomas and meningiomas were also found to overexpress both COX and lipoxygenase enzymes compared with normal brain tissue. Overexpression of both COX-1 and COX-2 enzymes was recently documented in human gliomas.^{60–65} In astrocytic gliomas, COX-1 was predominantly expressed in macrophages/microglial cells of the tumour mass, whereas COX-2 expression was mostly found in tumour cells and in single macrophages/microglial cells in perinecrotic areas.⁶⁴ COX-2 expression was shown to correlate with the pathological grade⁶¹ and proliferation rate of gliomas.⁶⁰ However, Matsuo *et al* reported that the staining intensity of GBM was relatively weak, but that astrocytomas (grades II and III) and meningiomas were strongly positive for COX-2.⁶⁵ Shono *et al* found that patients with glioma who had the highest expression of COX-2 had the shortest survival. COX-2 was identified as the strongest predictor of outcome, independent of all other variables examined.⁶³

A differential accumulation of COX isoforms during oligodendroglial tumour progression has also been recorded, with higher COX-1 expression in tumour cells of those patients whose tumours have “upgraded” to a more malignant phenotype.⁶⁴ Both survival and the time to tumour progression were longer in those patients with a low COX-1 labelling score compared with patients whose tumours displayed high COX-1 expression. Oligodendroglial tumours, in particular the anaplastic variant that recurred, expressed a higher number of COX-2 positive cells (mostly neurones and astrocytes) than did healthy control brains. Oligodendroglioma COX-2 protein expression was also examined in our laboratories by Castilla *et al*. In this study, the predominant COX-2 staining was not attributable to macrophages/microglial cells or reactive astrocytes but rather to tumour cells. In addition, COX-2 immunoreactivity was associated with decreased patient survival, along with retention of chromosome 1p and a high MIB-1 labelling index, whereas negative immunoreactivity for COX-2 and allelic loss of chromosome 1p correlated with better survival.^{65a}

5-Lipoxygenase

Although much attention has been focused on the role of COX derived metabolites in cancer development and progression, accumulating evidence suggests that 5-LO derived eicosanoids may play equally important roles. 5-LO mRNA and/or protein was found to be overexpressed in human lung,⁶⁶ prostate,⁶⁷ pancreatic,⁶⁸ and colon cancers.⁶⁹ Boado *et al* reported increased expression of 5-LO mRNA in three of three GBM and three of three meningioma surgical specimens examined. The increase in 5-LO mRNA was greater in the GBM samples than in the meningiomas.⁷⁰

Recent work from our laboratory suggested that surgical specimens of high grade astrocytomas and meningiomas overexpress 5-LO. Immunohistochemical analysis of surgical astrocytoma specimens shows that tumour cells and infiltrating macrophages/microglial cells express high amounts of 5-LO protein. In contrast, little if any 5-LO staining is detected in non-tumorous white matter control tissue. In addition, high amounts of both 5-LO mRNA and protein were expressed in some established GBM cell lines and a primary GBM cell culture. Staining for 5-LO was significantly more frequent in high grade than in low grade tumours (N Nathoo *et al*, unpublished data).

“Recent work from our laboratory suggested that surgical specimens of high grade astrocytomas and meningiomas overexpress 5-lipoxygenase”

We also detected high expression of 5-LO in primary meningioma cells and meningioma surgical specimens by

immunoblotting analysis.⁷¹ To date, it is unclear whether high 5-LO expression is present in meningioma precursor (leptomeningeal) cells or is a characteristic only of meningiomas. These data indicate that 5-LO is overexpressed in high grade astrocytomas and possibly in meningiomas and support the notion that, in addition to COX-2 derived eicosanoids, 5-LO eicosanoids may also play a role in tumorigenesis and progression of these brain tumours.

ENHANCED PRODUCTION OF EICOSANOIDS IN GLIOMAS AND MENINGIOMAS

The existence of high concentrations of COX and 5-LO enzymes and their substrate (AA) in glioma and meningioma cells creates a favourable environment for the efficient production of eicosanoids. Therefore, it is not surprising that cultured meningioma and glioma cells produce high amounts of prostaglandins, thromboxanes, and leukotrienes compared with normal brain tissue.^{72–76} AA metabolism capacity correlated well with proliferation patterns in neuroepithelial tumours,⁷⁷ and was shown to be related to steroid receptors in certain brain tumours; that is, some metabolites of AA were produced in different amounts in brain tumours, which either did or did not express steroid receptors.⁷⁸

In addition to *in vitro* production, eicosanoids can be detected systemically in patients with brain tumours, and their concentrations have been reported to decrease following neurosurgical resection or after treatment with specific inhibitors.^{73–79–80} In a recent clinical study, patients with malignant brain tumours (GBM, anaplastic astrocytoma, malignant meningioma, and medulloblastoma) had significantly higher concentrations of prostaglandin E₂ (PGE₂) in their plasma than did patients with benign brain tumours (meningiomas and pituitary adenomas), metastatic brain tumours, or non-tumorous control patients.⁷⁹ The high concentrations of PGE₂ decreased significantly (by a factor of three) three days after surgical removal of both malignant and benign brain tumours. Simmet *et al* detected high concentrations of leukotriene E₄ in the urine of eight patients with GBM.⁷³ The values dropped by 80% within seven days of surgical removal of the tumour. In contrast, five patients with meningiomas excreted less leukotriene E₄ and the values did not change significantly following meningioma removal. These data suggest that malignant brain tissue is the source of raised systemic COX and 5-LO derived eicosanoids.

“The use of cyclooxygenase 2 and 5-lipoxygenase inhibitors may even be an alternative to glucocorticoids for the treatment of eicosanoid induced peritumorous brain oedema”

Of particular importance is the observation that eicosanoids, produced by brain tumour cells, probably cause the development of tumour induced brain oedema.^{19–20} Peritumorous brain oedema is frequently found in both gliomas and meningiomas, and can contribute to the morbidity of patients with brain tumours. Glucocorticoids (such as dexamethasone) are used to treat peritumorous brain oedema. Although glucocorticoids often provide dramatic symptomatic relief for patients, they have many undesirable side effects.⁸¹

Of particular concern is the possibility that glucocorticoids may attenuate the cytotoxic effects of chemotherapy.^{82–85} For example, they activate the human O⁶-methylguanine DNA methyltransferase gene.⁸⁵ The product of this gene functions as a DNA repair protein that removes the mutagenic DNA adduct O⁶-alkylguanine, which is generated after exposure to alkylating antitumour drugs, such as those used for chemotherapy of GBM.

A substantial part of the beneficial effects of glucocorticoids in reducing oedema results from the suppression of PLA₂ activity and the reduction of free AA, in addition to inhibition of COX-2.²² However, there are reports suggesting that glucocorticoids upregulate the expression of 5-LO in rat neurones, human monocytes, and mast cells.^{86–88} Although the effects of glucocorticoids on 5-LO expression *in vivo* are unknown, these observations may have important therapeutic applications. These findings suggest that glucocorticoids and 5-LO inhibitors would have additive clinical benefits, and the use of COX-2 and 5-LO inhibitors may even be an alternative to glucocorticoids for the treatment of eicosanoid induced peritumorous brain oedema.

EICOSANOID INHIBITORS

An additional line of evidence supporting the concept that the COX-1, COX-2, and 5-LO proteins are involved in the tumorigenesis of gliomas and meningiomas comes from studies using COX and 5-LO inhibitors. Non-selective COX inhibitors, such as acetylsalicylic acid (aspirin), indomethacin, flurbiprofen, sulindac, and piroxicam, inhibited the growth of rat glioma cells both *in vitro* and *in vivo*,⁸⁹ and several human glioma cells *in vitro*.^{62–90–91} The COX-2 specific inhibitor NS-398 suppressed the proliferation of monolayer U-87 MG, U-251 MG human glioblastoma cell lines and the growth of three dimensional spheroids *in vitro*. In addition, NS-398 suppressed tumour cell migration and increased apoptotic cell death in spheroids.⁶¹ Another selective COX-2 inhibitor SC-236 was shown to be as effective as dexamethasone in extending the survival of rats implanted intracerebrally with 9L gliosarcoma cells,⁹² and to slow the tumour growth rate in nude mice with human U251 glioblastoma xenografts implanted in the hind leg.⁹³

These data suggest that gliomas and meningiomas should be added to the growing list of human malignancies that are considered suitable for treatment with COX-2 inhibitors.^{51–59–94} Malignant gliomas are resistant to radiotherapy and chemotherapy; therefore, selective COX-2 inhibitors are currently being examined in combination with these standard therapeutic modalities.⁹⁵ The rationale for these combination modality clinical trials comes from studies that reported the enhancing effect of a selective COX-2 inhibitor SC-236 on the tumour response to radiotherapy in a murine model using sarcoma and human U251 glioblastoma cell lines.^{93–96} Therefore, in the future the clinical use of COX-2 inhibitors may contribute to more efficient treatment for brain tumours.

Malignant gliomas are highly infiltrative tumours and this invasive nature represents a major obstacle to their effective treatment. Recent results obtained with inhibitors of thromboxane synthase suggest a novel anti-invasive approach for the treatment of gliomas.⁹⁷ Thromboxane synthase converts COX derived PGH₂ into thromboxane A₂ and it was found to be overexpressed in highly migratory subpopulations of glioma cells.⁹⁸ Specific thromboxane synthase inhibitors, such as furegrelate and dazmegrel, inhibited glioma cell migration and induced the apoptosis of these cells.⁶² When furegrelate was combined with camptothecin, it exerted a strong synergistic effect on the induction of apoptosis in glioma cells.⁶² Thus, further elucidation of the molecular mechanisms by which eicosanoids regulate glioma invasion will probably result in new ways to alter the migration and invasion of glioma cells and/or to increase their sensitivity to chemotherapeutic agents.

Boswellic acids, naturally occurring 5-LO inhibitors from the gum resin exudate of *Boswellia serrata* (frankincense), are known to kill meningioma and GBM cells *in vitro* by the induction of apoptosis.^{17–71–99–101} In a rodent glioma model, the gum resin exudate of *B. serrata* decreased glioma volume,

enhanced apoptosis, and more than doubled the survival time of treated rats compared with non-treated animals.¹⁰⁰ The effects of frankincense derived boswellic acids were examined in two other small studies of patients with gliomas. Boeker and Winking treated 29 patients with glioma using the H15 herbal preparation containing a gum resin exudate of *B serrata* for seven days.²⁶ There was no effect on tumour size, but a significant reduction of peritumorous brain oedema with a concurrent improvement of the clinical condition was recorded. Streffer *et al* treated 11 patients with glioma (including seven patients with GBM) and progressive peritumorous brain oedema with the H15 preparation, and saw a radiological and/or clinical response to treatment in three patients with GBM.²⁷

CURRENT LIMITATIONS AND FUTURE PERSPECTIVES

Investigations into the possible role of eicosanoids in the development and progression of brain tumours have progressed rapidly in the past few years. Today, specific inhibitors of eicosanoid pathways are being investigated in clinical trials of brain tumours. Nevertheless, many important questions remain unanswered.

Both benign meningiomas and GBMs contain raised amounts of AA and other ω -6 PUFAs, and express high amounts of COX-2 and 5-LO enzymes with active production of eicosanoids. If eicosanoids do indeed play an important role in the tumorigenesis of these tumours, it remains to be determined whether additional molecular changes in eicosanoid cascades or those unrelated to AA metabolism are responsible for such distinct biological behaviour. For example, it is not known which of the many PLA₂ enzymes are expressed and functionally active in brain tumour cells. Little is known about the expression of a variety of PG and leukotriene receptors and other eicosanoid enzymes, such as 12-LO, 15-LO, and cytochrome P450 enzymes in brain tumours. This is an important gap in our current knowledge because several specific prostaglandin and leukotriene receptor antagonists are presently available for non-tumour related clinical use.¹⁰² In addition, although the 12-LO pathway may be an important procarcinogenic cell regulator,^{103 104} 15-LO eicosanoids probably mediate the anticarcinogenic effects of PUFAs.^{103 105} The reasons for the existence of an unbalanced profile of fatty acids in brain tumour cells (accumulation of ω -6 and reduced amounts of ω -3 PUFAs) needs to be further elucidated. If dietary fatty acid intake is responsible, restoration of the equilibrium by dietary means in patients with brain tumours should be investigated.

Dietary fatty acids were shown to increase the expression of eicosanoid producing enzymes, such as COX-2,^{106 107} but they may modulate the growth and survival of brain tumour cells without their oxidation into eicosanoids. A family of steroid hormone receptors, peroxisome proliferator activated receptors, are activated not only by fatty acid metabolites but by dietary fatty acids themselves, and thus may provide a missing link between dietary fatty acids and cancer.^{108 109} Peroxisome proliferator activated receptor modulators are currently available for non-tumour related clinical use and are likely to stimulate investigations by brain tumour researchers in this rapidly growing field.^{108 110}

The tumour suppressive effects of COX and LO inhibitors may not only be related to the reduction in the production of eicosanoids, but they may also increase cellular concentrations of unesterified AA and induce the apoptosis of cancer cells.^{111 112} In other words, aberrantly raised expression of eicosanoid producing enzymes in tumour cells may promote tumorigenesis by increasing the intracellular utilisation of AA, thus suppressing unesterified AA mediated apoptosis. There is increasing evidence that polyunsaturated fatty acids,

such as γ linolenic, arachidonic, and eicosapentaenoic acids induce cell death and apoptosis of a variety of human cancer cell lines, including glioma cells, but not "normal" non-tumorigenic cells.¹¹³⁻¹¹⁶ However, the effects on cancer cells may depend on the concentration of exogenously added fatty acid. For example, γ linolenic acid impaired the growth of glioma spheroids only at concentrations higher than 100 μ M, whereas lower concentrations stimulated tumour cell growth and invasion.¹¹⁵ The cytotoxic action of PUFAs may be mediated through lipid peroxidation and free radical generation,¹¹⁷ in addition to AA stimulated conversion of sphingomyelin to the apoptosis inducer ceramide.¹¹² Intratumorous infusion of PUFAs in a rodent glioma model resulted in increased apoptosis and regression of tumour growth.¹¹⁸ Some improvements were documented in patients with malignant gliomas in whom γ linolenic acid was administered via a cerebral reservoir placed in the tumour bed or by direct intratumorous delivery.^{119 120} This treatment was found to be safe, but it remains to be determined whether higher concentrations of γ linolenic acid (or other PUFAs) and a longer duration of treatment alone or in combination with radiotherapy would result in better outcomes.

"Aberrantly raised expression of eicosanoid producing enzymes in tumour cells may promote tumorigenesis by increasing the intracellular utilisation of arachidonic acid (AA), thus suppressing unesterified AA mediated apoptosis"

Although the available evidence supports the idea that COX-2 and 5-LO proteins are aberrantly overexpressed and enzymatically active in gliomas and meningiomas, the data are less consistent with regard to the degree of staining and the nature of the cells that stain with specific antibodies. Previous investigators have used various antibodies, making it difficult to compare the results from different studies. Clearly, the use of specific antibodies directed against eicosanoid producing enzymes should be standardised so that the staining information can be used for clinical purposes, particularly those related to clinical trials with specific eicosanoid inhibitors. An additional concern when interpreting the COX-2 and 5-LO expression data is the possibility that upregulation of COX-2 and 5-LO in brain tumours may be partly caused by hypoxia introduced during and after tumour removal. Some studies suggest that hypoxia may induce both COX-2¹²¹ and 5-LO¹²² expression.

Although glucocorticoids, used to treat peritumorous brain oedema, may reduce the effectiveness of chemotherapy and radiotherapy, COX-2 inhibitors may in fact enhance these treatments, and combination trials with patients with brain tumours are currently in progress.⁹⁹ Further investigations are needed to establish whether COX-2 and 5-LO inhibitors can reduce peritumorous brain oedema in patients with brain tumours as effectively as glucocorticoids, but with less side effects. Selective COX-2 inhibitors were until recently considered to have fewer gastrointestinal adverse effects than non-selective, non-steroidal anti-inflammatory drugs. However, this premise is debatable because flaws in trial design and reporting of results cast doubts on the gastrointestinal, cardiovascular, and global safety profile of COX-2 specific inhibitors.^{24 123} Nevertheless, if effective for oedema reduction, these inhibitors will probably have fewer side effects than glucocorticoids. It remains to be determined whether the use of a combination of COX-2 and 5-LO inhibitors would be more effective than the use of either drug alone.

Take home messages

- The two most common types of human brain tumour, gliomas and meningiomas, overexpress eicosanoid producing enzymes, cyclooxygenases (COX-1 and COX-2) and lipoxygenases (5-LO, 12-LO, and 15-LO), and release a spectrum of eicosanoids that may promote tumorigenesis and the development of peritumorous brain oedema
- COX-2 and 5-LO inhibitors may be useful for the treatment of patients with malignant brain tumours, and they are currently under investigation in clinical trials
- Current and newer agents that specifically target key players of the eicosanoid cascade could change the approach to treating brain tumours, because their benefits may lie in their synergism with conventional cytotoxic treatments and/or with other novel agents targeted against other procarcinogenic pathways
- Further research is needed because the precise role of the eicosanoid cascade in the tumorigenesis of human brain tumours is still unclear

CONCLUSIONS

A new era of targeted treatments is rapidly evolving. The accrual of knowledge relating to the molecular pathways that govern the fundamental mechanisms of cell growth and survival, and how these pathways are altered in cancer, may be the key to improving patient survival and outcome from cancer in general, and particularly for patients with malignant gliomas. The downregulation or inhibition of tumour promoting eicosanoid cascades may represent one additional method of effectively and safely preventing and treating human brain tumours.

ACKNOWLEDGEMENTS

The authors would like to thank Ms M Tobin (Department of Neurosurgery) for helping to edit this manuscript.

Authors' affiliations

N Nathoo, G H Barnett, Brain Tumour Institute and Department of Neurosurgery, Cleveland Clinic Foundation, Cleveland, 44195 Ohio, USA
M Golubic, Brain Tumor Institute, Cleveland Clinic Foundation

REFERENCES

- Jemal A, Murray T, Samuels A, *et al.* Cancer statistics, 2003. *CA Cancer J Clin* 2003;53:5-26.
- Wrensch M, Minn Y, Chew T, *et al.* Epidemiology of primary brain tumors: current concepts and review of the literature. *Neuro-oncology* 2002;4:278-99.
- Kleihues P, Soylemezoglu F, Schauble B, *et al.* Histopathology, classification, and grading of gliomas. *Glia* 1995;15:211-21.
- Maher EA, Furnari FB, Bachoo RM, *et al.* Malignant glioma: genetics and biology of grave matter. *Genes Dev* 2001;15:1311-33.
- Russel DS, Rubinstein LJ. Tumours of the meninges and related tissues. In: Russel DS, Rubinstein LJ, eds. *Pathology of tumours of the nervous system*. Baltimore: Williams & Wilkins, 1989:449-532.
- Kujas M. Meningioma. *Curr Opin Neurol Neurosurg* 1993;6:882-7.
- Funk CD. Prostaglandins and leukotrienes: advances in eicosanoid biology. *Science* 2001;294:1871-5.
- Soberman RJ, Christman P. The organization and consequences of eicosanoid signaling. *J Clin Invest* 2003;111:1107-13.
- Smith WL, Garavito M, DeWitt DL. Prostaglandin endoperoxide H synthases (cyclooxygenases)-1 and -2. *J Biol Chem* 1996;271:33157-60.
- Turini ME, DuBois RN. Cyclooxygenase-2: a therapeutic target. *Annu Rev Med* 2002;53:35-57.
- Ford-Hutchinson AW. Regulation of leukotriene synthesis. *Cancer Metastasis Rev* 1994;13:257-67.
- Romano M, Catalano A, Nutini M, *et al.* 5-Lipoxygenase regulates malignant mesothelioma cell survival: involvement of vascular endothelial growth factor. *FASEB J* 2001;15:2326-36.
- Tong WG, Ding XZ, Witt RC, *et al.* Lipoxygenase inhibitors attenuate growth of human pancreatic cancer xenografts and induce apoptosis through the mitochondrial pathway. *Mol Cancer Ther* 2002;1:929-35.
- Tsuji M, DuBois RN. Alterations in cellular adhesion and apoptosis in epithelial cells overexpressing prostaglandin endoperoxide synthase 2. *Cell* 1995;83:493-501.
- Peppelenbosch MP, Tertoolen LG, Hage WJ, *et al.* Epidermal growth factor-induced actin remodeling is regulated by 5-lipoxygenase and cyclooxygenase products. *Cell* 1993;74:565-75.
- Capodici C, Pillinger MH, Han G, *et al.* Integrin-dependent homotypic adhesion of neutrophils. Arachidonic acid activates Raf-1/Mek/Erk via a 5-lipoxygenase-dependent pathway. *J Clin Invest* 1998;102:165-75.
- Park YS, Lee JH, Bondar J, *et al.* Frankincense-derived boswellic acids inhibit growth and motility of brain tumor cells [abstract]. *Complement Ther Med* 2002;10:117-18.
- Tsuji M, Kawano S, Tsuji S, *et al.* Cyclooxygenase regulates angiogenesis induced by colon cancer cells. *Cell* 1998;93:705-16.
- Constantini S, Tamir J, Gomori MJ, *et al.* Tumor prostaglandin levels correlate with edema around supratentorial meningiomas. *Neurosurgery* 1993;33:204-10.
- Black KL, Hoff JT, McGillicuddy JE, *et al.* Increased leukotriene C₄ and vasogenic edema surrounding brain tumors in humans. *Ann Neurol* 1986;19:592-5.
- Balsinde J, Balboa MA, Insel PA, *et al.* Regulation and inhibition of phospholipase A₂. *Annu Rev Pharmacol Toxicol* 1999;39:175-89.
- Goppelt-Strube M. Molecular mechanisms involved in the regulation of prostaglandin biosynthesis by glucocorticoids. *Biochem Pharmacol* 1997;53:1389-95.
- Burke JR, Witmer MR, Zusi C, *et al.* Competitive, reversible inhibition of cytosolic phospholipase A₂ at the lipid-water interface by choline derivatives that partially partition into the phospholipid bilayer. *J Biol Chem* 1999;274:18864-71.
- Wright JM. The double-edged sword of COX-2 selective NSAIDs. *Can Med Assoc J* 2002;167:1131-7.
- Hong SH, Avis I, Vos MD, *et al.* Relationship of arachidonic acid metabolizing enzyme expression in epithelial cancer cell lines to the growth effect of selective biochemical inhibitors. *Cancer Res* 1999;59:2223-8.
- Boeker DK, Winking M. Die Rolle von Boswellia-Sauren in der Therapie maligner Gliome. *Dtsch Arztebl* 1997;94:A1197-9.
- Streffler JR, Bitzer M, Schabet M, *et al.* Response of radiochemotherapy-associated cerebral edema to a phytotherapeutic agent, H15. *Neurology* 2001;56:1219-21.
- Steele VE, Holmes CA, Hawk ET, *et al.* Lipoxygenase inhibitors as potential cancer chemopreventives. *Cancer Epidemiol Biomarkers Prev* 1999;8:467-83.
- Safayhi H, Mack T, Sabieraj J, *et al.* Boswellic acids: novel, specific, nonredox inhibitors of 5-lipoxygenase. *J Pharmacol Exp Ther* 1992;261:1143-6.
- Lai LTY, German B. Modulation of fatty acid composition in murine brain by dietary unsaturated fats. *J Nutr Biochem* 1993;4:463-71.
- Clandinin MT, Cheema S, Field CJ, *et al.* Dietary fat: exogenous determination of membrane structure and cell function. *FASEB J* 1991;5:2761-9.
- Blair IA, Prakash C, Phillips MA, *et al.* Dietary modification of omega-6 fatty acid intake and its effect on urinary eicosanoid excretion. *Am J Clin Nutr* 1993;57:154-60.
- James MJ, Gibson RA, Cleland LG. Dietary polyunsaturated fatty acids and inflammatory mediator production. *Am J Clin Nutr* 2000;71(suppl):343S-8S.
- Ledwozyw A, Lutnicki K. Phospholipids and fatty acids in human brain tumors. *Acta Physiol Hung* 1992;79:381-7.
- Martin DD, Robbins MEC, Spector AA, *et al.* The fatty acid composition of human gliomas differs from that found in nonmalignant brain tissue. *Lipids* 1996;31:1283-8.
- Kokoglu E, Tuter Y, Yazici Z, *et al.* Profiles of the fatty acids in the plasma membrane of human brain tumors. *Cancer Biochem Biophys* 1998;16:301-12.
- Sinclair HM. Essential fatty acids—an historical perspective. *Biochem Soc Trans* 1990;18:756-61.
- German BJ, Dillard CJ, Whelan J. Introduction to symposium: biological effects of dietary arachidonic acid. *J Nutr* 1995;126:1076S-80S.
- Taber L, Chiu C-H, Whelan J. Assessment of the arachidonic acid content in foods commonly consumed in the American diet. *Lipids* 1998;33:1151-7.
- Narai T, Greig NH, DeGeorge JJ, *et al.* Intravenously injected radiolabelled fatty acids image brain tumor phospholipids in vivo: differential uptakes of palmitate, arachidonate and docosahexaenoate. *Clin Exp Metastasis* 1993;11:141-9.
- Giles GG, McNeil JJ, Donnan G, *et al.* Dietary factors and the risk of glioma in adults: results of a case-control study in Melbourne, Australia. *Int J Cancer* 1994;59:357-62.
- Lubin F, Farbstein H, Chetrit A, *et al.* The role of nutritional habits during gestation and child life in pediatric brain tumor etiology. *Int J Cancer* 2000;86:139-43.
- Chen H, Ward MH, Tucker KL, *et al.* Diet and risk of adult glioma in eastern Nebraska, United States. *Cancer Causes Control* 2002;13:647-55.
- Kaplan S, Novikov I, Modan B. Nutritional factors in the etiology of brain tumors: potential role of nitrosamines, fat, and cholesterol. *Am J Epidemiol* 1997;146:832-41.
- Cao Y, Dave KB, Doan TP, *et al.* Fatty acid CoA ligase 4 is up-regulated in colon adenocarcinoma. *Cancer Res* 2001;61:8429-34.

- 46 Stephenson DT, Manetta JV, White DL, et al. Calcium-sensitive cytosolic phospholipase A2 (cPLA2) is expressed in human brain astrocytes. *Brain Res* 1994;**637**:97-105.
- 47 Hernandez M, Burillo SL, Crespo MS, et al. Secretory phospholipase A2 activates the cascade of mitogen-activated protein kinases and cytosolic phospholipase A2 in the human astrocytoma cell line 1321N1. *J Biol Chem* 1998;**273**:606-12.
- 48 Oka S, Arita H. Inflammatory factors stimulate expression of group II phospholipase A2 in rat cultured astrocytes. Two distinct pathways of the gene expression. *J Biol Chem* 1991;**266**:9956-60.
- 49 Petr M, Origiano T, Wurster R. PLA₂ Activity regulates Ca⁺⁺ storage-dependent cellular proliferation. *Exp Cell Res* 1998;**244**:310-18.
- 50 Gupta RA, Dubois RN. Colorectal cancer prevention and treatment by inhibition of cyclooxygenase-2. *Nat Rev Cancer* 2001;**1**:11-21.
- 51 Ristimäki A, Sivula A, Lundin J, et al. Prognostic significance of elevated cyclooxygenase-2 expression in breast cancer. *Cancer Res* 2002;**62**:632-5.
- 52 Soslow RA, Dannenberg AJ, Rush D, et al. COX-2 is expressed in human pulmonary, colonic, and mammary tumors. *Cancer* 2000;**89**:2637-45.
- 53 Gupta S, Srivastava M, Ahmad N, et al. Over-expression of cyclooxygenase-2 in human prostate adenocarcinoma. *Prostate* 2000;**42**:73-8.
- 54 Tucker ON, Dannenberg AJ, Yang EK, et al. Cyclooxygenase-2 expression is up-regulated in human pancreatic cancer. *Cancer Res* 1999;**59**:987-90.
- 55 Okami J, Yamamoto H, Fujiwara Y, et al. Overexpression of cyclooxygenase-2 in carcinoma of the pancreas. *Clin Cancer Res* 1999;**5**:2018-24.
- 56 Zimmermann KC, Sarbia M, Weber AA, et al. Cyclooxygenase-2 expression in human esophageal carcinoma. *Cancer Res* 1999;**59**:198-204.
- 57 Buckman SY, Gresham A, Hale P, et al. COX-2 expression is induced by UVB exposure in human skin: implications for the development of skin cancer. *Carcinogenesis* 1998;**19**:723-9.
- 58 Higashi Y, Kanekura T, Kanzaki T. Enhanced expression of cyclooxygenase (COX)-2 in human skin epidermal cancer cells: evidence for growth suppression by inhibiting COX-2 expression. *Int J Cancer* 2000;**86**:667-71.
- 59 Ermet I, Dierkes C, Ermet M. Immunohistochemical expression of cyclooxygenase isoenzymes and downstream enzymes in human lung tumors. *Clin Cancer Res* 2003;**9**:1604-10.
- 60 Prayson RA, Castilla EA, Vogelbaum MA, et al. Cyclooxygenase-2 (COX-2) expression by immunohistochemistry in glioblastoma multiforme. *Ann Diagn Pathol* 2002;**6**:148-53.
- 61 Joki T, Heese O, Nikas DC, et al. Expression of cyclooxygenase 2 (COX-2) in human glioma and in vitro inhibition by a specific COX-2 inhibitor, NS-398. *Cancer Res* 2000;**60**:4926-31.
- 62 Kurzef F, Hagel Ch, Zapf S, et al. Cyclo-oxygenase inhibitors and thromboxane synthase inhibitors differentially regulate migration arrest, growth inhibition and apoptosis in human glioma cells. *Acta Neurochir (Wien)* 2002;**144**:71-87.
- 63 Shono T, Tofilon PJ, Bruner JM, et al. Cyclooxygenase-2 expression in human gliomas: prognostic significance and molecular correlations. *Cancer Res* 2001;**61**:4375-81.
- 64 Deininger MH, Meyermann R, Trautmann K, et al. Cyclooxygenase (COX)-1 expressing macrophages/microglial cells and COX-2 expressing astrocytes accumulate during oligodendroglioma progression. *Brain Res* 2000;**885**:111-16.
- 65 Matsuo M, Yonemitsu N, Zaitsum M, et al. Expression of prostaglandin H synthase-2 in human brain tumors. *Acta Neuropathol (Berl)* 2001;**102**:181-7.
- 65a Castilla EA, Prayson RA, Kanner AA, et al. Cyclooxygenase-2 in oligodendroglial neoplasms. *Cancer* 2003;**98**:1465-72.
- 66 Avis IM, Jett M, Boyle T, et al. Growth control of lung cancer by interruption of 5-lipoxygenase-mediated growth factor signaling. *Clin Invest* 1996;**97**:803-6.
- 67 Gupta S, Srivastava M, Ahmad N, et al. Lipoxygenase-5 is overexpressed in prostate adenocarcinoma. *Cancer* 2001;**91**:737-43.
- 68 Hennig R, Ding X, Tong W, et al. 5-Lipoxygenase and leukotriene B4 receptor are expressed in human pancreatic cancers but not in pancreatic ducts in normal tissue. *Am J Pathol* 2002;**161**:421-8.
- 69 Ohd JF, Nielsen CK, Campbell J, et al. Expression of the leukotriene D4 receptor CysLT1, COX-2, and other cell survival factors in colorectal adenocarcinomas. *Gastroenterology* 2003;**124**:57-70.
- 70 Boado RJ, Pardridge WM, Vinters HV, et al. Differential expression of arachidonate 5-lipoxygenase transcripts in human brain tumors: evidence for the expression of a multitranscript family. *Proc Natl Acad Sci U S A* 1992;**89**:9044-8.
- 71 Park YS, Lee JH, Harwalkar JA, et al. Acetyl-11-keto-beta-boswellic acid (AKBA) is cytotoxic for meningioma cells and inhibits phosphorylation of the extracellular-signal regulated kinase 1 and 2. *Adv Exp Med Biol* 2002;**507**:387-93.
- 72 Castelli MG, Chiabrando C, Fanelli R, et al. Prostaglandin and thromboxane synthesis by human intracranial tumors. *Cancer Res* 1989;**49**:1505-8.
- 73 Simmet T, Luck W, Winking M, et al. Identification and characterization of cysteinyl-leukotriene formation in tissue slices from human intracranial tumors: evidence for their biosynthesis under in vivo conditions. *J Neurochem* 1990;**54**:2091-9.
- 74 Inamura T, Abe M, Ikezaki K, et al. Leukotriene C₄ contents, synthase and catalase activity in human meningiomas. *Neural Res* 1992;**14**:405-10.
- 75 Cooper C, Jones HG, Weller RO, et al. Production of prostaglandins and thromboxane by isolated cells from intracranial tumors. *J Neural Neurosurg Psychiatry* 1984;**47**:579-84.
- 76 Kokoglu E, Tuter Y, Sandikci KS, et al. Prostaglandin E₂ levels in human brain tumor tissues and arachidonic acid levels in the plasma membranes of human brain tumors. *Cancer Lett* 1998;**132**:17-21.
- 77 Gaetani P, Butti G, Chiabrando C, et al. A study on the biological behavior of human brain tumors. Part I. Arachidonic acid metabolism and DNA content. *J Neurooncol* 1991;**10**:233-40.
- 78 Butti G, Gaetani P, Chiabrando C, et al. A study on the biological behavior of human brain tumors. Part II. Steroid receptors and arachidonic acid metabolism. *J Neurooncol* 1991;**10**:241-6.
- 79 Loh JK, Hwang SL, Lieu AS, et al. The alteration of prostaglandin E₂ levels in patients with brain tumors before and after tumor removal. *J Neurooncol* 2002;**57**:147-50.
- 80 Heldt RM, Winking M, Simmet T. Cysteinyl-leukotrienes as potential mediators of the peritumoral brain oedema in astrocytoma patients. Suppl to Naunyn-S [abstract]. *Arch Pharmacol* 1996;**353**:A538.
- 81 Frauman AG. An overview of the adverse reactions to adrenal corticosteroids. *Adverse Drug React Toxicol Rev* 1996;**15**:203-6.
- 82 Weller M, Schmidt C, Roth W, et al. Chemotherapy of human malignant glioma: prevention of efficacy by dexamethasone? *Neurology* 1997;**48**:1704-9.
- 83 Naumann U, Durka S, Weller M. Dexamethasone-mediated protection from drug cytotoxicity: association with p21^{WAF1/CIP1} protein accumulation? *Oncogene* 1998;**17**:1567-75.
- 84 Gorman AM, Hirt UA, Orrenius S, et al. Dexamethasone pre-treatment interferes with apoptotic death in glioma cells. *Neuroscience* 2000;**96**:417-25.
- 85 Biswas T, Ramana CV, Srinivasan G, et al. Activation of human O⁶-methylguanine-DNA methyltransferase gene by glucocorticoid hormone. *Oncogene* 1999;**18**:525-32.
- 86 Uz T, Swivedi Y, Savani PD, et al. Glucocorticoids stimulate inflammatory 5-lipoxygenase gene expression and protein translocation in the brain. *J Neurochem* 1999;**73**:693-9.
- 87 Colamorea T, Di Paola R, Macchia F, et al. 5-Lipoxygenase upregulation by dexamethasone in human mast cells. *Biochem Biophys Res Commun* 1999;**265**:617-24.
- 88 Riddick CA, Ring WL, Baker JR, et al. Dexamethasone increases expression of 5-lipoxygenase and its activating protein in human monocytes and THP-1 cells. *Eur J Biochem* 1997;**246**:112-18.
- 89 Aas AT, Tonnessen TI, Brun A, et al. Growth inhibition of rat glioma cells in vitro and in vivo by aspirin. *J Neurooncol* 1995;**24**:171-80.
- 90 Blomgren H, Kling-Andersson G. Growth inhibition of human malignant glioma cells in vitro by agents which interfere with the biosynthesis of eicosanoids. *Anticancer Res* 1992;**12**:981-6.
- 91 King JG Jr, Khalili K. Inhibition of human brain tumor cell growth by the anti-inflammatory drug, flurbiprofen. *Oncogene* 2001;**20**:6864-70.
- 92 Portnow J, Suleman S, Grossman SA, et al. A cyclooxygenase-2 (COX-2) inhibitor compared with dexamethasone in a survival study of rats with intracerebral 9L gliosarcomas. *Neurooncol* 2002;**4**:22-5.
- 93 Petersen C, Petersen S, Milas L, et al. Enhancement of intrinsic tumor cell radiosensitivity induced by a selective cyclooxygenase-2 inhibitor. *Clin Cancer Res* 2000;**6**:2513-20.
- 94 Smigel K. Arthritis drug approved for polyp prevention blazes trail for other prevention trials. *J Natl Cancer Inst* 2000;**92**:297-9.
- 95 Pannullo S, Balmaceda C, Serventi J, et al. Temozolomide (Temodar®) plus celecoxib (celebrex®) for treatment of malignant gliomas [abstract]. *Neuro-oncology* 2002;**4**:A239.
- 96 Milas L, Kishi K, Hunter N, et al. Enhancement of tumor response to gamma-radiation by an inhibitor of cyclooxygenase-2 enzyme. *J Natl Cancer Inst* 1999;**91**:1501-4.
- 97 Giese A, Bjerkvig R, Berens ME, et al. Cost of migration: invasion of malignant gliomas and implications for treatment. *J Clin Oncol* 2003;**21**:1624-36.
- 98 McDonough W, Tran N, Giese A, et al. Altered gene expression in human astrocytoma cells selected for migration: I. Thromboxane synthase. *J Neuropathol Exp Neurol* 1998;**57**:449-55.
- 99 Glaser T, Winter S, Groscurth P, et al. Boswellic acids and malignant glioma: induction of apoptosis but no modulation of drug sensitivity. *Br J Cancer* 1999;**80**:756-65.
- 100 Winking M, Sarikaya S, Rahmanian A, et al. Boswellic acids inhibit glioma growth: a new treatment option? *J Neurooncol* 2000;**46**:97-103.
- 101 Park YS, Lee JH, Bondar J, et al. Cytotoxic action of acetyl-11-keto-beta-boswellic acid (AKBA) on meningioma cells. *Planta Med* 2002;**68**:397-401.
- 102 Peters SP. Leukotriene receptor antagonists in asthma therapy. *J Allergy Clin Immunol* 2003;**111**(suppl 1):S62-70.
- 103 Shureiqi I, Lippman SM. Lipoxygenase modulation to reverse carcinogenesis. *Cancer Res* 2001;**61**:6307-12.
- 104 Pidgeon GP, Kandouz M, Meram A, et al. Mechanisms controlling cell cycle arrest and induction of apoptosis after 12-lipoxygenase inhibition in prostate cancer cells. *Cancer Res* 2002;**62**:2721-7.
- 105 Shappell SB, Gupta RA, Manning S, et al. 15S-Hydroxyeicosatetraenoic acid activates peroxisome proliferator-activated receptor gamma and inhibits proliferation in PC3 prostate carcinoma cells. *Cancer Res* 2001;**61**:497-503.
- 106 Badawi AF, El-Sohehy A, Stephen LL, et al. The effect of dietary n-3 and n-6 polyunsaturated fatty acids on the expression of cyclooxygenase 1 and 2 and levels of p21ras in rat mammary glands. *Carcinogenesis* 1998;**19**:905-10.
- 107 Rao CV, Hirose Y, Indranie C, et al. Modulation of experimental colon tumorigenesis by types and amounts of dietary fatty acids. *Cancer Res* 2001;**61**:1927-33.

- 108 **Kopelovich I**, Fay JR, Glazer RI, *et al.* Peroxisome proliferator-activated receptor modulators as potential chemopreventive agents. *Mol Cancer Ther* 2002;**1**:357–63.
- 109 **Vanden Heuvel JP**. Peroxisome proliferator-activated receptors: a critical link among fatty acids, gene expression and carcinogenesis. *J Nutr* 1999;**129**:575S–80S.
- 110 **Zander T**, Kraus JA, Grommes C, *et al.* Induction of apoptosis in human and rat glioma by agonists of the nuclear receptor PPARgamma. *J Neurochem* 2002;**81**:1052–60.
- 111 **Cao Y**, Pearman TA, Zimmerman GA, *et al.* Intracellular unesterified arachidonic acid signals apoptosis. *Proc Natl Acad Sci U S A* 2000;**97**:11280–5.
- 112 **Chan TA**, Morin PJ, Vogelstein B, *et al.* Mechanisms underlying nonsteroidal anti-inflammatory drug-mediated apoptosis. *Proc Natl Acad Sci U S A* 1998;**95**:681–6.
- 113 **Begin ME**, Sircar S, Weber JM. Differential sensitivity of tumorigenic and genetically related non-tumorigenic cells to cytotoxic polyunsaturated fatty acids. *Anticancer Res* 1989;**9**:1049–52.
- 114 **Williams JR**, Leaver HA, Ironside JW, *et al.* Apoptosis in human primary brain tumours: actions of arachidonic acid. *Prostaglandins Leukot Essent Fatty Acids* 1998;**58**:193–200.
- 115 **Bell HS**, Wharton SB, Leaver A, *et al.* Effects of N-6 essential fatty acids on glioma invasion and growth: experimental studies with glioma spheroids in collagen gels. *J Neurosurg* 1999;**91**:989–96.
- 116 **Vartak S**, McCaw R, Davis CS, *et al.* γ -Linolenic acid (GLA) is cytotoxic to 36B10 malignant rat astrocytoma cells but not to "normal" rat astrocytes. *Br J Cancer* 1998;**77**:1612–20.
- 117 **Preuss M**, Birnbaum GD, Darby CJ, *et al.* Role of antioxidant enzyme expression in the selective cytotoxic response of glioma cells to γ -linolenic acid supplementation. *Free Radic Biol Med* 2000;**28**:1143–56.
- 118 **Leaver HA**, Wharton SB, Bell HS, *et al.* Highly unsaturated fatty acid induced tumour regression in glioma pharmacodynamics and bioavailability of gamma linolenic acid in an implantation glioma model: effects on tumour biomass, apoptosis and neuronal tissue histology. *Prostaglandins Leukot Essent Fatty Acids* 2002;**67**:283–92.
- 119 **Das UN**, Prasad VVSK, Reddy DR. Local application of γ -linolenic acid in the treatment of human gliomas. *Cancer Lett* 1995;**94**:147–55.
- 120 **Bakshi H**, Mukherjee D, Backi A, *et al.* γ -Linolenic acid therapy of human gliomas. *Nutrition* 2003;**19**:305–9.
- 121 **Schmedtje JF Jr**, Ji YS, Liu WL, *et al.* Hypoxia induces cyclooxygenase-2 via the NF-kappaB p65 transcription factor in human vascular endothelial cells. *J Biol Chem* 1997;**272**:601–8.
- 122 **Tomimoto H**, Shibata M, Ihara M, *et al.* A comparative study on the expression of cyclooxygenase and 5-lipoxygenase during cerebral ischemia in humans. *Acta Neuropathol (Berl)* 2002;**104**:601–7.
- 123 **Gomez Cerezo J**, Lubomirov Hristov R, Carcas Sansuan AJ, *et al.* Outcome trials of COX-2 selective inhibitors: global safety evaluation does not promise benefits. *Eur J Clin Pharmacol* 2003;**59**:169–75.

ECHO

Tissue grading system is a reliable assay of response to treatment in gastric MALT lymphoma

Please visit the *Journal of Clinical Pathology* website (www.jclinpath.com) for a link to the full text of this article.

A histological grading system for gastric MALT lymphoma after treatment looks to be a promising tool for assessing residual disease in prospective drug trials. A pilot study has confirmed that the system has good internal reproducibility.

Good agreement was achieved among the scores of seven histopathologists who separately scored 45 sets of stained gastric biopsies for three criteria—lymphoid infiltrate, lymphoepithelial lesions, and stromal changes—from 10 patients treated for gastric MALT lymphoma. The results translated into four clinically meaningful scores: complete histological remission, probable minimal residual disease, responding residual disease, and no change.

The 10 patients comprised seven men and three women enrolled in one arm of a multicentre clinical trial of gastric MALT lymphoma and selected randomly. They had been treated to eradicate *Helicobacter pylori* and were followed up for a mean of 19 months, during which three to six sequential gastric biopsies were obtained. The histopathologists were blinded to the clinical results at follow up.

Gastric MALT lymphoma is closely associated with *H pylori* gastric infection. Eradicating the infection usually clears up the cancer, but ensuring patients are clinically clear of the disease entails prolonged follow up. Histological evaluation of residual disease is difficult but remains the only indication for the time being. The present standardised system was devised to help judge the extent of residual disease. Further testing of this scoring system on more patients is the next step, but in time it may be a useful complement to molecular studies.

▲ *Gut* 2003;**52**:1656.