ORIGINAL ARTICLE

Heterogeneity of the fibre sequence in subgenus C adenoviruses

A K Adhikary, U Banik, J Numaga, E Suzuki, T Inada, N Okabe

J Clin Pathol 2004;57:612–617. doi: 10.1136/jcp.2003.014944

See end of article for authors' affiliations

Correspondence to:
Dr A K Adhikary,
Infectious Disease
Surveillance Centre,
National Institute of
Infectious Diseases, 1-23-1
Toyama, Shinjuku-ku,
Tokyo 162-8640, Japan;
arun@nih.go.jp

Accepted for publication
17 December 2003

Aims: To determine the nucleotide sequences of adenovirus (Ad) types 1 and 6 fibre genes; to clarify the molecular basis of the distinct haemagglutination properties of subgenus C Ads and their phylogenetic relations.

Methods: Human Ad1 and Ad6 fibre genes were sequenced from genomic DNA by direct sequencing. Primer selection was based on alignment of the fibre gene of human Ad serotypes Ad2 and Ad5. Fibre based subgenus C specific polymerase chain reaction (PCR) was performed to check for deletions in field isolates of Ad6, as revealed by sequence analysis of the Ad6 prototype. A phylogenetic tree was constructed from the predicted amino acid (AA) sequences of the fibre gene of important Ads.

Results: Ad1 and Ad6 comprise 1746 and 1584 nucleotides, encoding 582 and 528 AA, respectively. Ad6 showed deletions in motifs 15–17 (51 AA) of the shaft when compared with Ad1, Ad2, and Ad5. Subgenus C specific PCR with both prototype and field isolates also showed deletions in Ad6. In the shaft and knob, AA homology was 58.82–72.91% and 68.89–74.59%, respectively. The tail was 100% conserved. Phylogenetically, Ad1 and Ad6, including Ad2 and Ad5, formed a subgenus specific cluster, like other serotypes.

Conclusions: The fibre gene (including the knob region) of subgenus C Ads is heterogeneous, providing the molecular basis for lack of crossreactivity in the haemagglutination inhibition test. This heterogeneity could be helpful in fibre based genotyping of subgenus C field isolates. Phylogeny might be useful for subgenus specific identification of important field strains.

In total, 51 adenovirus (Ad) serotypes have been identified as human pathogens, and on the basis of biochemical, immunological, and morphological criteria they are divided into six subgenera A–F.^{1–2} The members of subgenus C—Ad1, Ad2, Ad5, and Ad6—are the most frequently isolated human Ads, causing respiratory infection, gastroenteritis with prolonged excretion in children, ocular disease, and opportunistic infection among immunocompromised patients.^{3–5} Restriction endonuclease analysis of subgenus C field isolates revealed bewildering variations in their genome.^{6–9} Surprisingly, they have distinct antigenic sites for serum neutralisation (SN) and haemagglutination inhibition (HAI).^{10–11} The fibre protein, one of the major capsid proteins, is responsible for haemagglutination and the attachment of Ads to a specific cellular receptor at the early stage of infection through a receptor binding site present in the knob region.^{12–13} Analysis of the fibre protein of human Ad2 revealed that it possesses three distinct regions: an N-terminal tail of 43 amino acid (AA) residues; a central shaft of 357 residues, consisting of 22 pseudorepeats, usually 15 residues each (motif); and a C-terminal domain of 181 residues forming the knob.¹⁴ A later study by Chrobzek and Jarcot (1987)¹⁵ indicated that the tail comprises 44 AA residues. The tail and knob portions of the fibre are not significantly variable in length among the serotypes. However, the number of pseudorepeats in the shaft, which determines the length of the fibre, is variable. The tropism of Ads primarily depends on the presence of specific AA in the knob portion of the fibre.¹³ Investigation have shown that, as the length of fibre shaft increases, the efficiency of viral attachment to the cellular receptor also increases.^{16–17} Although subgenus C Ads account for 59% of all infections caused by Ads, Ad6 is responsible for only 4% of these infections, compared with 34.3% for Ad1, 42.8% for Ad2, and 18.6% for Ad5.⁴ Despite the fact that these subgenus C Ads

are important human pathogens, the fibre genes of Ad1 and Ad6 are yet to be studied.

“The fibre protein is responsible for haemagglutination and the attachment of adenoviruses to a specific cellular receptor at the early stage of infection through a receptor binding site present in the knob region”

In our study, the full length fibre genes of both Ad1 and Ad6 prototypes were sequenced for the first time with the following aims: (a) to find possible variations in the Ad1 and Ad6 fibre genes compared with the other serotypes of subgenus C, and to confirm any variations by fibre based polymerase chain reaction (PCR); (b) to correlate the variations of subgenus C Ads in the HAI test with heterogeneity of the fibre protein in the knob region; and (c) to clarify any evolutionary relations between Ad1 and Ad6 using a phylogenetic method.

MATERIALS AND METHODS

Viruses and DNA

Ad prototype strains Ad1 (adenoid 71), Ad2 (adenoid 6), Ad5 (adenoid 75), and Ad6 (tonsil 99) and serotypes of other subgenus C Ads were obtained from the American Type Culture Collection (ATCC; Rockville, Maryland, USA). Seventeen geographically and temporally diverse field isolates (Ad1:4, Ad2:4, Ad5:4, and Ad6:5) previously typed by the SN were used in our study.

Abbreviations: AA, amino acids; Ad, adenovirus; CAR, coxsackie virus and adenovirus receptor; HAI, haemagglutination inhibition; nt, nucleotides; PCR, polymerase chain reaction; SN, serum neutralisation

Extraction of viral DNA

The viruses were passaged in A549 cells and DNA was extracted from the infected cells by the method described previously.¹⁸ In short, when a greater than 75% cytopathic effect was seen, the cells were dislodged and pelleted by low speed centrifugation, washed three times with phosphate buffered saline, and resuspended in lysis buffer (10mM Tris/HCl, pH 7.4, 10mM EDTA, 1% sodium dodecyl sulfate). Next, the suspension was incubated with proteinase K (Sigma Chemical, St Louis, Missouri, USA). Cellular DNA was precipitated with 5M NaCl, and pelleted by centrifugation. After incubation with RNase A (Sigma Chemical) the supernatant was extracted twice in phenol/chloroform and precipitated in two volumes of absolute ethanol. After drying, DNA was suspended in 50 µl of double distilled water.

DNA sequencing

The genomic DNA was used as a template for DNA sequencing. The initial selection of primers was based on alignment of the Ad2 and Ad5 fibre genes (GenBank accession numbers j01907 (Ad2) and m18369 (Ad5)), and subsequently, because of the heterogeneous nucleotide sequences, the primer walking method was used.¹⁵ DNA sequencing was performed on the sense and antisense strands with overlapping primers. The cycle sequence reaction was carried out with an ABI prism big dye terminator cycle sequencing ready reaction kit (Applied Biosystems, Foster City, California, USA). The sequences were read by Genetic analyser model 310 (Applied Biosystems). The nucleotide sequences of the fibre gene were translated into AA sequences, and compared with the published AA sequences of Ad2 and Ad5 using the GENETYX-MAC program (Software Development Co, Tokyo, Japan).

Nucleotide sequence accession numbers

The nucleotide sequence data reported in this paper will appear in the DDBJ/EMBL/GenBank nucleotide sequence database with the accession numbers ab125750 (Ad1) and ab125751 (Ad6).

Ad subgenus C specific primers

Subgenus C specific PCR was performed with prototypes and field isolates to check the deletion in the shaft region of Ad6 field isolates noted by sequence analysis of the Ad6 prototypes. The design of the subgenus C specific primers (AdnCF, 5'-TGC TTG CGC THA AAA TGG GC-3' (654-674) and AdnCR, 5'-CGA TTC TTT ATT CYT GGG CAA TGT-3' (2228-2205)) was based on the alignment of the fibre gene sequences of Ad1 and Ad6 conducted in our laboratory (GenBank accession numbers Ad1 (ab125750), and Ad6 (ab125751)) and Ad2 and Ad5 obtained from GenBank (accession numbers Ad2 (j01917) and Ad5 (m18369)).¹⁵ The nucleotide positions of the primers refer to the fibre sequence of Ad2.

DNA amplification and detection

PCR amplification was carried out in a 50 µl reaction mixture containing 1 µl aliquots of DNA, 5 µl of 10× concentrated buffer, 0.5 µM of each of the primers, 200 µM of each deoxynucleoside triphosphate (dNTP), and 1.25 U of Taq polymerase (Boehringer-Mannheim, Mannheim, Germany). Sterile distilled water was used as a negative control. For field isolates, Ad2 DNA was used as a positive control. The assay was performed in a thermal cycler (model 9600-R; Perkin Elmer, Foster City, California, USA). Thermal cycling consisted of preliminary denaturation for three minutes at 94°C, followed by 35 cycles of denaturation at 94°C for one minute, annealing at 47°C for one minute, and extension at 72°C for two minutes, with a final extension at 72°C for seven

minutes. The reaction products (5 µl aliquots) were electrophoresed on a 1.2% horizontal agarose gel. After staining with ethidium bromide the gel was photographed with a CCD camera. The specificity of the test was evaluated with pooled DNA from the A, B, D, E, and F subgenera of Ad and non-adenoviral DNA.

Phylogenetic analysis

A phylogenetic tree was constructed according to the full length predicted AA sequences of the fibre genes obtained from the GenBank. Sequence alignment was performed with Clustal X (<http://www-igbmc.u-strasbg.fr/BioInfo/>) with parameters provided in Clustal X. Evolutionary distances were estimated by Kimura's two parameter method. A phylogenetic tree with 1000 bootstrap replicates was generated by the neighbour joining method with Clustal W and plotted by Tree View (<http://taxonomy.zoology.gla.ac.uk/rod/treeview.html>).

RESULTS

Sequence analysis

The Ad1 and Ad6 fibre genes were 1746 and 1584 nucleotides (nt) long (table 1), corresponding to 582 and 528 AA, respectively. The predicted AA sequences were consistent with the structural domains described for Ad2, which possesses three distinct regions: an N-terminal tail (44 AA); a central shaft of 357 residues consisting of 22 pseudorepeats, usually 15 residues; and a C-terminal knob of 181 residues. The fibre tail of Ad1 and Ad6 was 132 nt long, corresponding to 44 AA, and showed 100% conservation among the members of subgenus C. However, at the nucleotide level the homology varied from 89.39% to 93.18%. The two conserved sequences—2-KRλR (where λ is A in all the subgenus C serotypes) and FNPVYPYD—are present in subgenus C, similar to the other Ad subgenera (fig 1).¹⁹

The shaft region of Ad1 and Ad6 displayed a 15 AA basic structural unit (motif), containing hydrophobic glycine (G) or proline (P) residues at position 8 of the β sheet/β turn model (fig 1).¹⁴ Ad1 comprises 357 polypeptides and displayed 22 repeats of the 15 AA motifs, similar to Ad2 and Ad5. The shaft of Ad6 was relatively shorter, with 306 AA (19 motifs) as a result of the deletion of motifs 15 to 17 (fig 2). The third motifs were 19 AA long except for in Ad6, which showed deletion of a residue (T) in position 8. In the shaft region, AA homology among the serotypes varied from 58.82% to 72.91%. The Ad1 and Ad6 fibre knobs were 543 and 534 nt long, corresponding to 181 and 178 AA, respectively. Their AA homology varied between 68.89% and 74.59% (table 2). Ad1 and Ad6 showed an overall homology of 71.95% at the DNA level and 70.1% at the AA level. The overall AA homology among the members of subgenus C varied from 65.06% to 75.47%, and is considered to be heterogeneous.

Table 1 Length of the fibre genes (number of nucleotides) and proteins (number of amino acids) among the members of subgenus C adenoviruses

		Tail	Shaft	Knob	Overall
Ad1	DNA	132	1071	543	1746
	Protein	44	357	181	582
Ad2	DNA	132	1071	543	1746
	Protein	44	357	181	582
Ad5	DNA	132	1071	540	1743
	Protein	44	357	180	581
Ad6	DNA	132	918	534	1584
	Protein	44	306	178	528

Ad, adenovirus.

Tail									
Ad1	1	MKRARFSEDT FNPVYPYDTE	TGPPTVPFLLT	PPFVSPNGFQ	ESPP	44			
Ad2	1	MKRARFSEDT FNPVYPYDTE	TGPPTVPFLLT	PPFVSPNGFQ	ESPP	44			
Ad5	1	MKRARFSEDT FNPVYPYDTE	TGPPTVPFLLT	PPFVSPNGFQ	ESPP	44			
Ad6	1	MKRARFSEDT FNPVYPYDTE	TGPPTVPFLLT	PPFVSPNGFQ	ESPP	44			
Shaft									
Motif	45	GVLSRLRLSEPLVTSNGM	61	Motif	220	LTVGTGGQGVADNNA	234		
1	45	GVLSRLRVSEPLDTSNGM	61	12	220	LTVVTGPGVIVEQNS	234		
	45	GVLSRLRLSEPLVTSNGM	61		220	LTVATGPGVTINNTS	234		
	45	GVLSRLRLSEPLVTSNGM	61		219	LTLTGQGGVAVHNNL	233		
2	62	LALKMGSSGLALDKAGN	77		235	LHTKVTGAIGYDTSNG	250		
	62	LALKMGSSGLTLDKAGN	77	13	235	LRTKVAGAIYDSSNN	250		
	62	LALKMGSSGLSLDEAGN	77		235	LQTKVTGALGFDSQGN	250		
	62	LALKMGSSGLSLDQAGN	77		234	LHTKVTGAIGFDTSNG	249		
3	78	LTSQNTITVTEPLKTKSN	96		251	MELKTGGGVVRVDSVNR	267		
	78	LTSQNTITVTPQLKTKSN	96	14	251	MEIKTGGGMRINNNL	265		
	78	LTSQNTITVSPPLKTKSN	96		251	MQLNVAGGLRIDSQNR	267		
	78	LTSNTIT-VSQPLKTKSN	95		250	MELKTGDLVYDSAGPNQKIH	270		
4	97	ISLETSAFLITVTSGL	111		268	LILDVDYFFDAQSQR	283		
	97	ISLETSAFLITITSGA	111	15	266	LILDVDYFFDAQTKLR	281		
	97	INLETSAFLITVTSGL	111		268	LILDVSYFFDAQNQLN	283		
	96	INLETSAFLITVSSGA	110			-----			
5	112	LTVATAAPLVVAGNS	126		284	LKLGGGLYVNSSTHN	299		
	112	LTVATAAPLVITVSGA	126	16	282	LKLGGGLYINAS-HN	296		
	112	LTVAAAAPLMVAGNT	126		284	LRLGGGLFINS-A-HN	298		
	111	LTMATTSPLVSDNT	125			-----			
6	127	LTQSQAPLTVQDSK	141		300	LLDLNKNKGLHLFTTGN-SKKLE	320		
	127	LSVQSQAPLTVQDSK	141	17	297	LLDINYNRGLYLFNASNNTKKLE	318		
	127	LTMQSQAPLTVHDSK	141		299	LLDINYNKGLYLFNASNNTKKLE	320		
	126	LTMQSQAPLTVQDSK	140			-----			
7	142	LSIATKQPLTVSEGR	156		321	VNLKTKGLIFDTDA	335		
	142	LSIATKQPLTVSDGR	156	18	319	VSINKSSGLNFDNTA	333		
	142	LSIATQGLTVSEGR	156		321	VNLSTAKGLMFDATA	335		
	141	LSIATKQPLTVLDGR	155		271	INLNTTKGLAFDNTA	285		
8	157	LALQTSGLSATDNT	172		335	VAIHAAQGLEFGNDTS--TNTNP	356		
	157	LALQTSAPLSGSDSDT	172	19	334	IATNAGKGLFDTNTSESPIINP	356		
	157	LALQTSGLPLTIDSDT	172		335	IATNAGKGLFEGSPNA--PNTNP	356		
	156	LALQTSAPLSATDNTA	171		286	ITINAGKGLFETDSSNG---NP	305		
9	173	LEITTSPPFITINGS	187		357	LKTKLGLGLDYDSNGG	372		
	173	LTVTASPLITATGS	187	20	357	IKTKIGSGIDYNENGA	372		
	173	LTITASPLITATGS	187		357	LKTKIGHGLEFDNSKA	372		
	172	LTITASPLITANGS	186		306	IKTKIGSGIQNTINGA	321		
10	188	LGVNMENFLYNNNGK	202		373	MIPKLGTLGSLFDTTGA	388		
	188	LGINMEDPIYVNNNGK	202	21	373	MITKLGAFLSFDNSGA	388		
	188	LGLDLKEPIYTQNGK	202		373	MVPKLGTLGSLFDTTGA	388		
	187	LAVTMENFLYNNNGK	202		322	MVAKLGTLGSLFDSGGA	337		
11	203	LGLRVAGPLQVINDSHA	219		389	ITVGNKSDDKLTL	401		
	203	IGIKISGFLQVQNSDT	219	22	389	ITIGNKNDKDLTL	401		
	203	LGLKYGAPLHVTDLNT	219		389	ITVGNKNDKDLTL	401		
	202	LGLKIGGFLQVATDASHA	218		338	ITMGSINNDKDLTL	350		
Knob									
	402	WTPPDP SPNCQIHSEK	DAKLTVLTK	CGSQVLATVS	ALAVRGLAP	ISGTISSAHI	457		
	402	WTPPDP SPNCRIHSDN	CKFTVLTK	CGSQVLATVA	ALAVSGDLSS	MTGTVASVSI	457		
	402	WTPPAP SPNCRLNAEK	DAKLTVLTK	CGSQILATVS	VLA VKGLAP	ISGTQSAHL	457		
	351	WTPPDP SPNCRIASDK	CKLTILATK	CGSQILGTVS	ALAVSGNMA	ISGTQSAHL	406		
	458	ILRFNEHGVIL	MNHSLLDPQY	WNFRKGLTN	ATAYTNAVGF	MPNLKAYPKT	QSKTAKSNIV	517	
	458	FLRFQDQGVIL	MENSSLLKKHY	WNFRNGNSTN	ANPYTNAVGF	MPNLLAYPKT	QSKTAKSNIV	517	
	458	IIRFDENGVL	LNNSFLDPEY	WNFRNGDLTE	GTAYTNAVGF	MPNLSAYPKS	HGKTAKSNIV	517	
	407	VLRFDQDGVIL	MSNSLLDKQY	WNFRNGDSTN	GQPYTYAVGF	MPNLKAYPKT	QSKTAKSNIV	466	
	518	SQVYLNGEKE	KPMILTITLN	GTDENQTPA	-STYSISFSW	SWPSNQTYIG	QTFATNSYTF	576	
	518	SQVYLHGDKT	KPMILTITLN	GTSESTETSE	VSTYMSFTW	SWESGK-YTT	ETFATNSYTF	576	
	518	SQVYLNGDKT	KPVTLTITLN	GTQETGDTTP	-SAYSMSFSW	DWSG-HNYIN	EIFATNSYTF	575	
	467	SQVYLNGDKS	KPLHFTITLN	GTDE---TNQ	VSKYSISFSW	SWNSGQ-YTN	DKFATNSYTF	522	
	577	SYIAQE	582						
	577	SYIAQE	582						
	576	SYIAQE	581						
	523	SYIAQE	528						

Figure 1 Comparison of predicted amino acid (AA) sequences of adenovirus 1 (Ad1), Ad2, Ad5, and Ad6 fibre proteins. The nucleotide sequences of Ad1 and Ad6 have been deduced from their respective DNA sequences. Ad2 and Ad5 AA sequences were obtained from GenBank using the accession numbers j01917 and m18369, respectively. The conserved residues in the tail are underlined. The four sequences have been aligned to obtain maximal homology. The structural domains tail, shaft, and knob are marked. The shaft region has been divided into motifs. Bold letters indicate hydrophobic residues.

Polymerase chain reaction

The primer pair AdnCF/AdnCR amplified approximately 1575 bp, 1575 bp, and 1572 bp products from Ad1, Ad2, Ad5 prototypes, respectively, whereas the Ad6 prototype yielded a

1413 bp product (fig 2A). Seventeen field isolates (table 3) also yielded clearly visible products of the expected length (fig 2B). Ad6 could be distinguished from the other members of subgenus C by the length of the amplicon (1413 bp).

Figure 2 (A) Ethidium bromide stained agarose gel showing polymerase chain reaction (PCR) products of subgenus C prototype strains. The lane designations indicate the respective prototypes: lane N, negative control; lane M, DNA molecular size marker III (Boehringer Mannheim). (B) Ethidium bromide stained agarose gel showing PCR products of subgenus C field isolates. Lane i1 (66–315), Ad1; lane i2 (YC 66–283), Ad2; lane i5 (YC 77–82), Ad5; lane i6 (YC 92–516T), Ad6; lane N, negative control; lane P, positive control (Ad2p); lane M, DNA molecular size marker III.

PCR using DNA from other subgenera of adenovirus and non-adenoviral DNA yielded no amplified products, indicating a high specificity of the test.

Phylogenetic analysis

Ad1 and Ad6 were included in a cluster formed by subgenus C adenoviruses. Ad6 was found to be more closely related to Ad1 than either Ad2 or Ad5. Other serotypes included in phylogeny also formed subgenus specific clusters. However, the cluster formed by subgenus F was divided into two subclusters because of the presence of two different fibres (long fibre and short fibres; fig 3).

Table 2 Fibre gene nucleotide and amino acid homologies (%) among the members of the subgenus C adenoviruses						
Fibre		Ad1/2	Ad1/5	Ad1/6	Ad6/2	Ad6/5
Tail	DNA	90.15	90.91	93.18	90.91	89.39
	Protein	100.0	100.0	100.0	100.0	100.0
Shaft	DNA	72.71	74.91	69.16	64.52	63.90
	Protein	70.56	72.91	64.71	62.18	58.82
Knob	DNA	68.56	72.56	72.24	73.85	69.98
	Protein	70.88	74.59	73.48	74.03	68.89
Overall	DNA	72.73	75.39	71.95	69.42	67.72
	Protein	72.87	75.47	70.10	68.73	65.06

DNA and amino acid homologies were obtained using GENETYX-MAC software (Software Development Co, Tokyo, Japan).
Ad, adenovirus.

Figure 3 A phylogram showing adenovirus (Ad) subgenus specific clusters indicated by letters A–F. Ad1 and Ad6 are included in cluster C. The subgenus F is divided into short and long fibre subclusters marked by L and S with the respective prototypes. The phylogenetic tree was reconstructed from the predicted amino acid sequences of the full length fibre gene of all important human adenovirus serotypes under the GenBank accession for subgenus A: Ad12 (x73487) and Ad31 (x76548); subgenus B: Ad3 (m12411), Ad7p (m23696), Ad7h (z48954), Ad11p (l08231), Ad11a (l08232), Ad14 (ab065116), Ad16 (u06106), Ad21 (u06107), Ad34p (u10271, ab073168), Ad34a (u10272), and Ad35 (u10272); subgenus C: Ad1 (ab125750), Ad2 (j01917), Ad5 (m18369), and Ad6 (ab125751); subgenus D: Ad8 (x74660), Ad9 (x74659), Ad15 (x72934), Ad17 (nc002067), Ad19p (u69130), Ad19a (u69131), and Ad37 (u69132); subgenus E: Ad4 (x76547); and subgenus F: Ad40 (l19443 and m28822) and Ad41 (x16583, m60327).

DISCUSSION

Subgenus C Ads differ from other subgenera in that they have a highly variable genomic organisation, they are in widespread use as a vector for gene delivery, and they are responsible for approximately 60% of human infections caused by the different Ad serotypes. Our study revealed two important findings about the subgenus C Ad fibre protein. First, the structure of the knob is heterogeneous, which could explain the serological characteristics of subgenus C Ads and, second, the fibre shaft of Ad6 is relatively short, and this may be related to its lower rate of infection.

Subgenus C Ads are unique because of the lack of crossreactivity in both the SN and HAI tests.²⁰ The HAI test in a large number of geographically diverse isolates showed that the fibre epitopes are relatively conserved, although the exact location and size of these epitopes are yet to be determined.^{10 11 21} Serotypes such as Ad40 and Ad41 or Ad19

Table 3 Fibre based subgenus C specific polymerase chain reaction

Original identification		
Isolate	Serotype	Subgenus/amplicon (bp)
YC 66-113	Ad1	C/1572-1575
YC 66-283	Ad2	C/1572-1575
YC 66-284	Ad2	C/1572-1575
YC 66-315	Ad1	C/1572-1575
YC 72-141	Ad5	C/1572-1575
YC 72-143	Ad2	C/1572-1575
YC 77-82	Ad5	C/1572-1575
YC 84-31	Ad6	C/1413
YC 84-37	Ad5	C/1572-1575
YC 85-70	Ad6	C/1413
YC 85-75	Ad1	C/1572-1575
YC 85-77	Ad1	C/1572-1575
92-515T	Ad5	C/1572-1575
91-1141	Ad6	C/1413
92-516T	Ad6	C/1413
92-523T	Ad6	C/1413
92-566T	Ad2	C/1572-1575

Ad, adenovirus.

and Ad37 show 97.7% and 95.5% homology in the knob region, respectively, and show major crossreactivity in the HAI test. Therefore, higher homology (> 80%) in the knob region is thought to be the basis of serological crossreactivity as a result of epitope sharing.^{19, 20, 22} Although not an absolute rule, lower homology in the knob region may imply the presence of distinct epitopes, as demonstrated by the absence of major crossreactivity in HAI. Thus, subgenus C Ads have no major crossreactivity in HAI with other subgenera or among themselves probably because they have low homology in the fibre knob region, between 68.89% and 74.59.

“Adenovirus 1 (Ad1) is more closely related to Ad6 than to Ad2 or Ad5”

To date, studies with Ad2 and Ad5 have shown that the subgenus C fibre knob mediates binding to the coxsackie virus and Ad receptor (CAR), and is thus a major determinant of viral tropism.²³ Mounting evidence suggests that at least two factors determine viral infectivity and tropism, namely: (a) the fibre knob–primary receptor interaction and (b) the length and flexibility of the fibre shaft.^{16, 17} The length of the fibre shaft is variable because of deletion or insertion of certain numbers of repeats. The formation of a recombinant fibre shaft with decreasing length and increasing rigidity significantly reduced infectivity and attachment of Ad5.²⁴ The major differences between the Ad6 fibre shaft and that of the other subgenus C Ads are the deletion of the 15–17th motifs (51 residues) of the shaft and a single residue (T) deletion in the eighth position of motif 3. The deletion was also detected in PCR products of the field isolates of Ad6. It may be possible that deletion of 51 AA residues in the shaft decreases its flexibility and reduces the mobility of the knob.²² The third motif of subgenus C is 19 AA long except in Ad6. In all Ads, the third motif carries a hydrophilic residue in the second hydrophobic position of the β strand (position 15 in subgenus C). The fibre bends above the penton base at a position corresponding to this third motif of the shaft, which is also a flexing point to allow fibre movement. In Ad6, the third motif is 18 AA long (rather than 19 AA as in other subgenus C Ads), as a result of the deletion of a single residue (T) at position 8, and this may also slightly decrease the flexibility of the fibre shaft.²⁵ Both these factors present in Ad6 might act together to decrease its efficiency of binding to the cellular receptor. Further experiments with

Take home messages

- The fibre gene (including the knob region) of subgenus C adenoviruses (Ad) is heterogeneous, providing the molecular basis for their lack of crossreactivity in the haemagglutination inhibition test
- This heterogeneity might be useful in fibre based genotyping of subgenus C field isolates
- Phylogeny might be useful for subgenus specific identification of important field strains
- The fibre shaft of Ad6 is relatively short, and this may be related to its lower rate of infection

recombinant Ad6 fibre proteins might be useful to investigate the relation between the length of the Ad6 fibre shaft and the rate of infection. It is also important to investigate whether Ad6 uses CAR or another cellular receptor.

In the phylogram, Ad1 and Ad6, together with Ad2 and Ad5, formed a subgenus C specific cluster, indicating that these Ads have the same ancestral origin. Ad1 is more closely related to Ad6 than to Ad2 or Ad5. Subgenus specific clusters were also formed by other serotypes as a result of high homology (60–80%) among the members of each subgenus.²⁶ Subgenus specific clusters of the fibre gene that included important Ads could be useful in subgrouping and understanding the evolution of Ads.

Sequence heterogeneity in the fibre knob provides the molecular basis of the distinct serological properties of subgenus C Ads in the HAI test. The inclusion of serotypes Ad1, Ad2, Ad5, and Ad6 in a single cluster defines the evolutionary relations between the subgenus C Ads. Furthermore, the sequence data of Ad1 and Ad6 reported in our study will be useful in the identification of subgenus C field strains, either by fibre based type specific PCR or by comparison of sequences.

ACKNOWLEDGEMENTS

Supported by grant-in-aid for Scientific Research from the Japan Society for the Promotion of Science (JSPS), Japan.

Authors' affiliations

A K Adhikary, U Banik, T Inada, N Okabe, Infectious Disease Surveillance Centre, National Institute of Infectious Diseases, 1-23-1 Toyama, Shinjuku-ku, Tokyo 162-8640

J Numaga, Department of Ophthalmology, Graduate School of Medicine, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

E Suzuki, Department of Developmental Medical Sciences, Graduate School of Medicine, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

REFERENCES

1. **Shenk T**. Adenoviridae: the viruses and their replication. In: Fields BN, Knipe DM, Howley PM, eds. *Fields virology*, 5th ed. Philadelphia: Lippincott-Raven Publishers, 1996:2111–48.
2. **De Jong JC**, Wermenbol AG, Verweij-Uijterwall, et al. Adenovirus from human immunodeficiency virus-infected individuals, including two strains that represent new candidate serotype Ad50 and Ad51 of species B1 and D, respectively. *J Clin Microbiol* 1999;**37**:3940–45.
3. **Fox JP**, Hall CE, Cooney MK. The Seattle virus watch. VII. Observation of adenovirus infection. *Am J Epidemiol* 1977;**105**:362–68.
4. **Schmitz H**, Wigand R, Heinrich W. World wide epidemiology of adenovirus infections. *Am J Epidemiol* 1983;**117**:455–66.
5. **Hierholzer JC**. Adenoviruses in the immunocompromised host. *J Clin Microbiol* 1992;**5**:262–74.
6. **Adrian T**. Genome polymorphism of human adenovirus subgenus C. *Arch Virol* 1996;**141**:1021–31.
7. **Adrian T**, Wigand R, Wadell G. Serological and biochemical characteristics of intermediate adenovirus strains of subgenus D. *Arch Virol* 1987;**97**:347–57.

- 8 **Adrian T**, Wolf U. Adenovirus 6 genome types: mapping of restriction site alterations on the genome. *Res Virol* 1989;**140**:545–50.
- 9 **Fife KH**, Ashley R, Corey L. Isolation and characterization of six new genome types of human adenovirus types 1 and 2. *J Clin Microbiol* 1985;**21**:20–3.
- 10 **Adam E**, Nasz I, Lengyel A. Antigenic homogeneity among the members of the adenovirus hexon types of subgenus C. *Arch Virol* 1995;**140**:1297–301.
- 11 **Kaier B**, Wigand R. Antigenic homogeneity of adenovirus type 1, 2, 5, and 6. *J Med Virol* 1986;**18**:283–7.
- 12 **Norby E**. The structural and functional diversity of adenovirus capsid components. *J Gen Virol* 1969;**5**:221–36.
- 13 **Arnberg N**, Mei Y-F, Wadell G. Fiber genes of adenoviruses with tropism for the eye and the genital tract. *Virology* 1997;**227**:239–44.
- 14 **Green NM**, Wrigley NG, Russell WC. Evidence for a repeating cross- β sheet structure in the adenovirus fiber. *EMBO J* 1983;**2**:1357–65.
- 15 **Chrobozek J**, Jacrot B. The sequence of adenovirus fiber: similarities and differences between serotypes 2 and 5. *Virology* 1987;**161**:549–54.
- 16 **Shayakhmetov DM**, Lieber A. Dependence of adenovirus infectivity on length of the fiber shaft domain. *J Virol* 2003;**74**:10274–86.
- 17 **Nemerow GR**. Flexibility of the adenovirus fiber is required for efficient receptor interaction. *J Virol* 2000;**77**:7225–35.
- 18 **Adhikary AK**, Numaga J, Kaburaki T, *et al*. Genetic characterization of adenovirus type 8 isolated in Hiroshima city over a 15 year period. *J Clin Pathol* 2003;**56**:120–5.
- 19 **Pring-Akerblom P**, Heim A, Trijssenaar FEJ. Conserved sequences in the fibers of epidemic keratoconjunctivitis associated human adenovirus. *Arch Virol* 1997;**142**:205–11.
- 20 **Hierholzer JC**. Antigenic relationship among the 47 human adenoviruses determined in reference horse antisera. *Arch Virol* 1991;**121**:179–97.
- 21 **Adhikary AK**, Inada T, Suzuki E, *et al*. Characterization of hexon and fibre gene of a novel strain of adenovirus involved in epidemic keratoconjunctivitis. *J Clin Pathol* 2004;**57**:95–7.
- 22 **Kidd AH**, Erasmus MJ, Tiemessen CT. Fiber sequence heterogeneity in subgroup F adenoviruses. *Virology* 1990;**179**:139–50.
- 23 **Bergelson DM**, Cunningham JA, Droguett G, *et al*. Isolation of common receptor for coxsackie B viruses and adenoviruses 2 and 5. *Science* 1997;**275**:1320–3.
- 24 **Nakamura T**, Sato K, Hamada H. Reduction of natural adenovirus tropism to the liver by both ablation of fiber-coxsackie virus and adenovirus receptor interaction and use of replaceable short fiber. *J Virol* 2003;**77**:2512–21.
- 25 **Ruigrok RWH**, Barge A, Mittal SK, *et al*. The fiber of bovine adenovirus type 3 is very long but bent. *J Gen Virol* 1994;**75**:2069–73.
- 26 **Pring-Akerblom P**, Adrian T. Sequence characterization of the adenovirus 31 fiber and comparison with serotypes of subgenera A to F. *Res Virol* 1995;**146**:343–54.