CASE REPORT

Pineal yolk sac tumour with a solid pattern: a case report in a Chinese adult man with Down's syndrome

H W Tan, A Ty, S G N Goh, M C Wong, A Hong, K L Chuah

J Clin Pathol 2004;57:882–884. doi: 10.1136/jcp.2004.016659

Intracranial germ cell tumours are rare. The incidence of primary intracranial yolk sac tumour is even more uncommon, with only two reported cases being associated with Down's syndrome in the English literature. This report details the findings of yolk sac tumour in the pineal region affecting a 22 year old Chinese man with Down's syndrome. Histology revealed yolk sac tumour with only a solid pattern, potentially mimicking the more common germinoma in the pineal region. No other germ cell components were identified. This is the third report of intracranial yolk sac tumour manifesting in a patient with trisomy 21. The pathology of this tumour and its differential diagnoses are discussed.

In patients with Down's syndrome (DS), germinoma is the more common intracranial germ tumour,^{1–3} and two cases of intracranial yolk sac tumour (YST) have been documented in the English literature.^{1–4} In this report, we describe the appearance of YST with only a solid pattern identified on histology presenting as a pineal neoplasm associated with DS. Unlike the previously reported cases,^{1–4} our case lacked the reticulated pattern typically present in YST. Although DS related intracranial germ cell tumours tend to occur in atypical sites—for example, basal ganglia^{1–3}—the tumour in our patient was midline.

CASE REPORT

A 22 year old Chinese man with DS and Hashimoto's thyroiditis confirmed by positive serology for thyroid microsomal antibody, presented with headache of one month's duration associated with nausea, vomiting, and some weakness in the lower limbs when walking. Physical examination showed characteristic Down's syndrome, fair complexion, and coarse skin with increased buccal pigmentation. The testicles were not enlarged. Blood investigations

revealed hypopituitarism and diabetes insipidus. Computer tomography of the brain showed a large pituitary–pineal axis mass, with homogeneous enhancement and focal calcification, compressing the back of the third ventricle and the aqueduct and causing hydrocephalus. A ventriculoperitoneal shunt was inserted and a subsequent magnetic resonance imaging scan delineated two solid enhancing masses. The larger mass, measuring 4 × 3.7 × 3.6 cm, was centred in the pineal region and the smaller mass, measuring 1.8 cm in maximum diameter, was present in the pituitary infundibular/suprasellar region. Biochemistry of the cerebrospinal fluid revealed a raised α fetoprotein (α FP) concentration of up to 995 μ g/litre, whereas the human chorionic gonadotrophin (hCG) value was normal. A stereotactic biopsy was performed. He was scheduled for chemotherapy followed by radiotherapy; unfortunately, the patient developed fever of unknown origin postoperatively and died 1.5 months after the biopsy. Consent for necropsy could not be obtained from the next of kin.

PATHOLOGICAL FINDINGS

Grossly, the biopsy specimen received in small pieces was 1.3 cm in diameter when aggregated. On cytology, the crushed smears showed loose aggregates of medium sized hyperchromatic undifferentiated cells with ovoid nuclei, moderate anisonucleosis, discernible nucleoli, and scant to moderate amounts of cytoplasm (fig 1). Paraffin wax embedded sections showed a tumour composed of sheets of round to polygonal cells with apparently two populations of cells (fig 2). One population was composed of uniform cells resembling a germinoma, whereas another population was composed of larger cells disclosing eosinophilic polygonal cytoplasm and pale nuclei with prominent nucleoli. Occasional, more pleomorphic cells were noted but syncytiotrophoblastic cells were absent in the biopsy. Mitotic activity was readily apparent. Importantly, scattered periodic acid Schiff (PAS)-diastase positive hyaline globules were seen (fig 3). However, a typical reticulated pattern with Schiller-Duval bodies of YST was not identified and no fibrovascular septae associated with a lymphocytic infiltrate were present. No other germ cell components were featured in the biopsy. Immunohistochemistry showed the tumour cells to be positive for α FP (RB365A1; 1/60 dilution; Neomarkers, Fremont, California, USA) (fig 4) and α 1 antitrypsin or AAT (A012; 1/1200 dilution; Dako, Glostrup, Denmark). Staining for cytokeratin AE 1/3 (M3515; 1/200 dilution; Dako) was positive in the larger cells, sparing the areas resembling germinoma (fig 5). Staining for placenta-like alkaline phosphatase or PLAP (MS734-S1; 1/20 dilution; NeoMarkers), S100 protein (Z0311; 1/2000 dilution; Dako), leucocyte common antigen (250-2136; prediluted; Ventana,

Figure 1 Smear cytology revealing single and clusters of medium sized cells, which appeared to be undifferentiated. Note the presence of nucleoli (Papanicolaou stain; original magnification, $\times 600$).

Abbreviations: α FP, α fetoprotein; AAT, α 1 antitrypsin; β hCG, β human chorionic gonadotrophin; DS, Down's syndrome; PLAP, placenta-like alkaline phosphatase; YST, yolk sac tumour

Figure 2 Histology revealing a cellular tumour composed of solid sheets of cells with clear to eosinophilic cytoplasm lacking fibrous septae, with accompanying lymphocytes. No Schiller-Duval bodies were noted (haematoxylin and eosin; original magnification, $\times 200$).

Tucson, Arizona, USA), hCG (RB059-A1; 1/300 dilution; NeoMarkers), glial fibrillary acidic protein (Z0334; 1/1000 dilution; Dako), and CD30 (M0751; 1/60 dilution; Dako) was negative. MIB-1 (M7240; 1/70 dilution; Dako) staining was 90%. The light microscopy and immunohistochemical findings agreed with a diagnosis of YST with a solid pattern.

DISCUSSION

The histological classification of germ cell tumours arising in the brain follows those arising from the gonads and extragonadal sites.⁵ YST is known to exhibit a variety of microscopic patterns.⁶ Although the epithelial element may proliferate in solid sheets,⁵ the solid component often coexists with more typical patterns, having at least a focally reticular pattern with the characteristic Schiller-Duval bodies,⁶ thereby aiding the diagnosis. Although it is possible that the tumour in our report may contain other morphological patterns typical of YST, which were not biopsied, the presence of only the solid component of YST on biopsy makes the diagnosis challenging. Although the tumour in our case can be classified as pure using the criteria of Matsutani *et al*,⁷ the possibility of coexisting germ cell components apart from YST, which were not identified on biopsy, has to be entertained.

“The possibility of a purely coincidental manifestation of yolk sac tumour in the background of Down’s syndrome in our case cannot be excluded”

Although raised serum and cerebrospinal fluid α FP concentrations are helpful in the diagnosis and follow up treatment of intracranial YST,¹⁻⁸ they are insufficient for a definitive diagnosis of YST in lieu of proper histological diagnosis because embryonal carcinomas and immature teratomas without YST can be associated with raised α FP values. Histologically, YST with a solid pattern in our case mimics germinoma or even embryonal carcinoma,^{3,9} by exhibiting non-overlapping, relatively uniform nuclei, clear cytoplasm, and well defined cell borders. However, fibrous septae with lymphocytic infiltrates characteristic of germinomas are absent. In contradistinction to YST, embryonal carcinoma displays a greater degree of nuclear pleomorphism. PAS-diastase resistant globules of YST are not featured in germinoma or embryonal carcinoma, underscoring the need of careful light microscopic examination with the PAS stain to identify this feature. In problematic situations, immunohistochemical studies are useful because YST is positive for cytokeratin, α FP, and AAT, whereas germinoma

Figure 3 Careful examination of tumour disclosing occasional eosinophilic globules present within the tumour (periodic acid Schiff with diastase pretreatment; original magnification, $\times 600$).

Figure 4 Antibodies to α fetoprotein decorating the tumour cells, confirming the diagnosis of yolk sac tumour (avidin-biotin complex method; original magnification, $\times 600$).

is PLAP positive. Cytokeratin expression alone cannot be used to differentiate YST from germinoma because germinomas can be cytokeratin positive.^{9,10} Embryonal carcinoma, although positive for cytokeratin and CD30, is negative for α FP. Interestingly, in our case, cytokeratin staining in some of the solid areas, especially those resembling germinoma, was negative, as seen in the solid variant of testicular YST.⁹ However, immunohistochemical staining for α FP should not be used alone in the diagnosis of YST because gastrointestinal epithelium and primitive neuroepithelial elements of immature teratoma may be positive for α FP.¹¹ Rather, a constellation of light microscopy features in conjunction with a broad

Figure 5 Note the presence of tumour cells resembling germinoma surrounded by cytokeratin (AE1/3) positive cells (avidin-biotin complex method; original magnification, $\times 200$).

Take home messages

- We report a yolk sac tumour in the pineal region affecting a 22 year old Chinese man with Down's syndrome—this is the third report of intracranial yolk sac tumour manifesting in a patient with trisomy 21
- Histology revealed yolk sac tumour with only a solid pattern, potentially mimicking the more common germinoma in the pineal region
- In cases of diagnostic difficulty a constellation of light microscopy features in conjunction with a broad panel of immunohistochemical stains and clinical findings is essential

panel of immunohistochemical stains and clinical correlation was pivotal in achieving the right diagnosis in our case.

Although an association between DS and an increased occurrence of intracranial germ cell tumours has been noted,^{1–3 12} where trisomy 21 is proposed to cause tumour suppressor gene inactivation or direct activation of an oncogene,³ the possibility of a purely coincidental manifestation of YST in the background of DS in our case cannot be excluded. Whether trisomy 21 plays a significant role in the pathogenesis of YST or not, it is nonetheless apparent that intracranial YST has an extremely poor prognosis, with survival of more than three years being exceptional, regardless of therapeutic modalities.^{8 13}

Authors' affiliations

H W Tan, S G N Goh, K L Chuah, Department of Pathology, Singapore General Hospital, Outram Road, Singapore 169608, Singapore
A Ty, M C Wong, Department of Neurology, National Neuroscience Institute, Singapore General Hospital Campus
A Hong, Department of Neurosurgery, National Neuroscience Institute

Correspondence to: Dr K Leong Chuah, Department of Pathology, Singapore General Hospital, Outram Road, Singapore 169608, Singapore; gzzckl@singnet.com.sg

Accepted for publication 25 February 2004

REFERENCES

- 1 **Chik KW**, Li CK, Shing MMK, *et al*. Intracranial germ cell tumours in children with and without Down syndrome. *J Pediatr Hematol Oncol* 1999;**21**:149–51.
- 2 **Nakashima T**, Nishimura Y, Sakai M, *et al*. Germinoma in cerebral hemisphere associated with Down syndrome. *Childs Nerv Syst* 1997;**13**:563–6.
- 3 **Tanabe M**, Mizushima M, Anno Y, *et al*. Intracranial germinoma with Down's syndrome: a case report and review of the literature. *Surg Neurol* 1997;**47**:28–31.
- 4 **Rueda-Pedraza ME**, Heifetz SA, Sesterhenn IA, *et al*. Primary intracranial germ cell tumors in the first two decades of life. A clinical, light-microscopic, and immunohistochemical analysis of 54 cases. *Perspect Pediatr Pathol* 1987;**10**:160–207.
- 5 **Rosenblum MK**, Matsutani M, Van Meir EG. CNS germ cell tumours. In: Kleihues P, Cavenee WK, eds. *World Health Organisation classification of tumours. Pathology and genetics of tumours of the nervous system*. Lyon: IARC Press, 2000:208–14.
- 6 **Scully RE**, Young RH, Clement PB. Germ cell tumors: general features and primitive forms. In: *Tumors of the ovary, maldeveloped gonads, fallopian tube and broad ligament*, 3rd series. Washington DC: AFIP, 1998:245–55.
- 7 **Matsutani M**, Sano K, Takakura K, *et al*. Primary intracranial germ cell tumours: a clinical analysis of 153 histologically verified cases. *J Neurosurg* 1997;**86**:446–55.
- 8 **Kirikae M**, Arai H, Hidaka T, *et al*. Pineal yolk sac tumour in a 65-year-old man. *Surg Neurol* 1994;**42**:253–8.
- 9 **Eglen DE**, Ulbright TM. The differential diagnosis of yolk sac tumour and seminoma. Usefulness of cytokeratin, alpha-fetoprotein, and alpha-1-antitrypsin immunoperoxidase reactions. *Am J Clin Pathol* 1987;**88**:328–32.
- 10 **Nagakawa Y**, Perentes E, Ross GW, *et al*. Immunohistochemical differences between intracranial germinomas and their gonadal equivalents. An immunoperoxidase study of germ cell tumors with epithelial membrane antigen, cytokeratin, and vimentin. *J Pathol* 1988;**156**:67–72.
- 11 **Yamashita M**, Kancai H, Kamiya K, *et al*. Immature teratoma producing alpha-fetoprotein without components of yolk sac tumour in the pineal region. *Childs Nerv Syst* 1997;**13**:225–8.
- 12 **Hori A**, Walter GF, Haas J, *et al*. Down syndrome complicated by brain tumours: case report and review of the literature. *Brain Dev* 1992;**14**:396–400.
- 13 **Ushio Y**, Kachi M, Kuratsu J, *et al*. Preliminary observations for a new treatment in children with primary intracranial yolk sac tumour or embryonal carcinoma. Report of five cases. *J Neurosurg* 1999;**90**:133–7.