ty in using the chair. The patient's arm is at a height which allows the operator the maximum comfort at what can become, under adverse conditions, a back-breaking task.

We are grateful to Dr E. G. Rees, consultant pathologist, for his helpful comments, and to Mr J. D. Hamlett, chief technician, Morbid Anatomy Department, for the photographs

Letter to the Editor

Survey of Hepatitis in Laboratories

Members of the ACP who helped with this survey by submitting data on proformas may be interested in a preliminary summary of the findings for the three years of the survey (1970-72). Brief reports concerning the first two years of the survey were made to the Pathology Section of the Royal Society of Medicine and to the 90th General Meeting of the ACP (Grist, 1973 a and b).

Hepatitis cases were reported by 12 (5%) of 244 laboratories in 1970, 15 (7%) of 215 in 1971, and eight (2%) of 337 in 1972. Of the laboratories reporting cases, in each of the three years more than average tested specimens from haemodialysis, transplant, and haemophilia units (see table), and to a lesser extent tested for hepatitis B antigen (HBAg). Different organizational arrangements for HBAg testing did not appear to be associated with differences in hepatitis risk.

The attack rate per 100 000 in staff of all grades was 116 (13/11 253) in 1970, 174 (18/10 380) in 1971, and 60 (9/15 103) in 1972, ie, 109 per 100 000 per annum for the three years combined. Considered by grade and discipline, the highest attack rates in 1970 were in science micro-

biologists (1/126 = 0.79%) and biochemical technicians (7/1438 = 0.49%); in 1971 the highest rates were in science biochemists (3/315 = 0.95%), medical haematologists (2/315 = 0.63%), and medical morbid anatomists (2/390 = 0.51%); the highest rates in 1972 were in science biochemists (1/469 = 0.21%) and haematology technicians (4/2775 = 0.14%). The highest attack rates per 100 000 per annum for the three years combined were 316 (1/317) for science microbiologists, 266 (3/1120) for science biochemists, 210 (3/1429) for medical anatomists, and 183 (9/4900) for biochemical technicians.

Tests for HBAg had been carried out in 47 cases. Of the 16 positive reactors, five were admitted to hospital as were three of the 31 negatives. No deaths were reported. More than half the patients were under 30 years old, both sexes were involved, and cases were scattered through the country over the three-year period without significant clusters.

Although no remarkable or alarming trends emerged from the survey, some of the differences in hepatitis attack rate between different areas of work underline the modest but significant risk associated with laboratory work and the need to take sensible, but not exaggerated, precautions.

These findings will be reported more fully after further analysis.

I thank the ACP for assistance with the expenses of this survey and all those members whose cooperation made it possible. I am also grateful for the help given by my secretary, Miss E. H. Simpson, and by Mrs M. E. Martin, of Glasgow Northern Hospitals Medical Records Department.

NORMAN R. GRIST
University of Glasgow,
Department of Infectious Diseases,
Ruchill Hospital,
Glasgow, G20 9NB.

References

Grist, N. R. (1973a). Hazards in the clinical pathology laboratory. *Proc. roy. Soc. Med.*, 66, 795-796.
Grist, N. R. (1973b). A survey of hepatitis in laboratories. (Abstr.) *J. clin. Path.*, 26, 388.

Feature	Percentages of Laboratories in Each Year with Indicated Features ¹					
	1970		1971		1972	
	T	H	T	H	T	H
Testing haemodialysis unit specimens	28	50	29	60	25	50
Testing transplant unit specimens	16	33	18	40	17	38
Testing haemophilia unit specimens	18	33	19	27	19	38
Testing drug addiction unit specimens	9	8	13	27	11	0
Performing HB Ag. tests	15	25	16	27	17	25

Table Work of laboratories in relation to hepatitis

¹T = total reporting laboratories
H = laboratories reporting hepatitis

J Clin Pathol: first published as 10.1136/jcp.27.1.84 on 1 April 1974. Downloaded from <http://jcp.bmj.com/> on April 26, 2023 by guest. Protected by copyright.