



In UTI
first-time success
in 9 out of 10 patients*

— plus the safety
of a penicillin

Uticillin (carfecillin sodium) is indicated for the treatment of acute and chronic urinary tract infections caused by sensitive organisms including *E. coli*, *Proteus* spp., *Strep. faecalis*, and particularly *Pseudomonas*. Following oral administration, Uticillin is rapidly absorbed and hydrolysed to carbenicillin, achieving high concentrations in the urine.

**Brit. J. Clin. Pract.* (1974), 28, 349.

Uticillin

Full prescribing information available on request

Beecham Research Laboratories, Welwyn, Herts. SG8 2PE

BRL 025



Even more signs of quality in Radioassays



Oestriol
New 1976



Digoxin
New 1976



Folate
New 1976



FSH
New 1976



cGMP
New 1976



Thyopac* - 3



Thyopac* - 4



Thyopac* - 5



T₃RIA



T₄RIA



Cortipac



ACTH



Insulin



HPL



Anti-DNA



cAMP

Every time you see our symbol on a radioassay kit you know you can depend on its performance. Because we spend a lot of time discovering the needs of radioassay users and on the production and quality control of our kits, we can guarantee they are precise, reliable and simple to use.

In 1976 we added five more kits to our range, making sixteen in all, and there will be more to come.



**The Radiochemical Centre
Amersham**

Full information is available on request.
The Radiochemical Centre Limited, Amersham, England. Telephone: 024-04-4444.
In the Americas: Amersham/Searle Corp., Illinois 60005. Telephone: 312-593-6300.
In W. Germany: Amersham Buchler GmbH & Co., KG., Braunschweig. Telephone: 05307-4693-97.

*TradeMark 0952