


\*trademark

## four new kits join our radioassay line-up

Our range of radioassay kits for diagnosis or research keeps growing to meet your needs. What stays the same is the quality. Our four latest kits are designed for simplicity, reliability and convenience, and maintain the high standards of manufacture and quality control our customers have come to expect.



A unique kit for the RIA of  $\beta$ -thromboglobulin in platelet research.

Fast, convenient assay for T3 using PEG separation.

New 400 test pack especially suitable for automated use.

Sensitive and specific assay for unconjugated oestriol in maternal serum.

### The sign of quality



The Radiochemical Centre  
Amersham


Full information on all our radioassay kits is available on request.  
The Radiochemical Centre Limited, Amersham, England. Telephone: 024-04-4444  
In the Americas: Amersham Corporation, Illinois 60005. Telephone: 312-593-6300  
In W. Germany: Amersham Buchler GmbH & Co KG, Braunschweig. Telephone: 05307-4693-97

# Septtrin success *in vivo* begins *in vitro*

Septtrin can be relied on to perform as well in the patient as it does in the Petri dish.

Many published reports show that a wide range of bacterial isolates are sensitive *in vitro* to Septtrin, with sensitivities of up to 99% achieved for urinary tract pathogens<sup>1,2</sup>. And clinical success rates are correspondingly high so you can recommend Septtrin with confidence.

1. Robertson, M.H. (1976). *Update*, 12, 11.  
2. McAllister, T.A. (1974). *Int. med. Res.*, 2, 400.




Wellcome

Septtrin contains trimethoprim and sulphamethoxazole.  
Full prescribing information is available on request.  
Wellcome Medical Division, The Wellcome Foundation Ltd,  
Berkhamsted, Herts

RESEARCH IS OUR ONLY SHAREHOLDER

# SEPTRIN



In UTI  
first-time success  
in 9 out of 10 patients\*

— plus the safety  
of a penicillin

Uticillin (carfecillin sodium) is indicated for the treatment of acute and chronic urinary tract infections caused by sensitive organisms including *E. coli*, *Proteus* spp., *Strep. faecalis*, and particularly *Pseudomonas*. Following oral administration, Uticillin is rapidly absorbed and hydrolysed to carbenicillin, achieving high concentrations in the urine.

\**Brit. J. Clin. Pract.* (1974), 28, 349.

# Uticillin


For further information, contact your local representative or write to:  
Beecham Research Laboratories, 9100 Rockledge Drive, Broomfield, Colorado 80020, U.S.A.  
Beecham Research Laboratories, 5080 Kingsway, London W8 5AH, U.K.

BRL 025



**Beecham Research Laboratories,**

# MYOCARDIAL INFARCTION?



**ELECTROCARDIOGRAM** and **CARDIO-RELATED ENZYMES**  
and CK-MB\*\*

**TOGETHER**  
**Your diagnosis is only minutes away**

\*CK - Creatine kinase \*\*CK-MB=CK-MB Isoenzyme

By performing CK, CK-MB and ECG you can now rapidly differentiate myocardial infarction from other causes of chest pain.

CK and CK-MB can easily detect re-infarctions

Save your time and coronary bed time with these simple tests.

For further information on how Cardio-related Enzyme assays can help you, please contact:

## **BCL**

The Boehringer Corporation (London) Ltd.,  
Bell Lane, Lewes, East Sussex BN7 1LG Telephone: (07916) 71611