Significance of bacterial and white cell counts in midstream urines

PJ LITTLE, BARBARA A PEDDIE, AND ADRIENNE R SINCOCK

From the Department of Renal Medicine, Christchurch Hospital, Christchurch, New Zealand

SUMMARY The results of cultures carried out on urine collected by a midstream technique (MSU) have been compared with the results of cultures of urine collected by suprapubic bladder aspiration (SPA) in the same 903 patients. Comparisons indicate that when there are more than $100 \times 10^6/l$ ($100\,000/ml$) Gram-negative bacteria in the midstream urine this finding is confirmed by suprapubic aspiration in 92% of patients, and in 70% when the urine contained this number of Gram-positive bacteria. When the culture contained more than one species (mixed organisms), the presence of infection was confirmed in only 11%. When there were $10\text{--}100 \times 10^6/l$ bacteria in the MSU the finding of Gram-negative bacilli still indicated urinary infection in 74% of patients, but the presence of Gram-positive organisms was confirmed in only 30% and mixed organisms in 2%. In asymptomatic patients, the presence of white cells in the urine was not helpful in confirming the diagnosis of infection. The nature of the organism found in the MSU is almost as important as the number of bacteria in assessing the validity of the result.

The results of urine cultures performed on urine collected by a midstream technique are often misleading in that in some patients heavy contamination may occur. Kass¹ provided statistical analyses of multiple midstream urines (MSUs) to establish the criteria separating contamination from true infection. He found the presence of a concentration of organisms greater than $100 \times 10^6/l$ ($> 100\,000/ml$) could be regarded as indicating urinary infection, but when the counts were less than $100 \times 10^6/l$ repeat specimens were rarely alike.² He showed that the finding of greater than 100×10^6 organisms per litre ($> 100\,000$ per ml) of urine is reproducible on 80% of occasions and that it may therefore be assumed that on 20% of occasions it does not represent true infection of the urinary tract.³ Although the technique of suprapubic bladder aspiration (SPA) gives a more accurate result, this method is often inconvenient and difficult to those not familiar with its use. In this paper an attempt is made to correlate the results of urine examinations using the two techniques so that a more accurate estimate can be made of the significance of the result of examination of an MSU. The results were assessed with

particular reference to the identity of the organisms present in the urine.

Method

In this study SPA was carried out in 903 women, and the result was compared with a preceding MSU examination. The patients were 840 asymptomatic women examined at the first attendance at an antenatal clinic, 53 asymptomatic student nurses, and 10 women presenting with symptomatic urinary tract infection (UTI). All the patients had an initial MSU containing $\geq 10 \times 10^6/l$ organisms ($\geq 10\,000/ml$).

The MSU was collected by a nurse employed for this purpose. Perineal cleansing with sterile water preceded collection. SPA was carried out within seven days of the MSU by a method previously described.⁴ All urine cultures were carried out on the day of urine collection. Bacterial counts were made by pour plate dilution technique.⁵

Urinary white cell counts (WCC) were made on 462 of the urine samples collected by MSU and on 37 of those collected by SPA.^{6,7} An upper limit of $10 \text{ cells} \times 10^6/l$ ($10/cmm$) in uncentrifuged urine was used in this study.⁸ No limits of normality of

Received for publication 4 April 1979

Table 1 Result of suprapubic aspiration (SPA) after a midstream urine (MSU) containing $10\text{--}100 \times 10^6/\text{l}$ bacteria ($10\ 000\text{--}100\ 000/\text{ml}$)

Organism	MSU (No. tested)	SPA	
		No. infected	%
<i>E. coli</i>	28	21	75
<i>Klebsiella</i> and <i>Enterobacter</i> spp	2	2	
<i>Proteus mirabilis</i>	5	3	
<i>Staphylococcus</i> spp	36	9	25
<i>Streptococcus</i> spp	11	4	36
Mixed	100	2	2
Total	182	41	23

WCC in urine collected by SPA are available. The technique used requires the administration of a diuretic and forced fluid intake so that considerable dilution of urine occurs.

Results

The results of SPAs done on patients with $10\text{--}100 \times 10^6/\text{l}$ ($10\ 000\text{--}100\ 000/\text{ml}$) bacteria in the initial MSU are shown in Table 1. Infection was confirmed in 26 of 35 patients with $10\text{--}100 \times 10^6/\text{l}$ Gram-negative bacilli in the MSU. Where the MSU had $10\text{--}100 \times 10^6/\text{l}$ Gram-positive organisms, 13 out of 47 bladder urines were found to be infected. Only two of 100 urines with mixed cultures of $10\text{--}100 \times 10^6/\text{l}$ bacteria in the MSU had infected SPAs.

Table 2 shows the results of the SPA when the MSU showed $> 100 \times 10^6/\text{l}$ ($> 100\ 000/\text{ml}$) bacteria. This demonstrates that where the initial MSU had $100 \times 10^6/\text{l}$ of a pure growth of *Escherichia coli*, *Proteus* spp, *Klebsiella* spp, or *Enterobacter* spp, 92% of the SPAs were infected. With $> 100 \times 10^6/\text{l}$ streptococci in the MSU 81% of SPAs were infected, and with $> 100 \times 10^6/\text{l}$ staphylococci 67% were infected. Where the initial MSU had a mixed culture of $> 100 \times 10^6/\text{l}$ bacteria only 11% of the bladder

Table 2 Result of suprapubic aspiration (SPA) after a midstream urine (MSU) containing $> 100 \times 10^6/\text{l}$ bacteria ($> 100\ 000/\text{ml}$)

Organism	MSU (No. tested)	SPA	
		No. infected	%
<i>E. coli</i>	399	370	93
<i>Klebsiella</i> and <i>Enterobacter</i> spp	12	11	92
<i>Proteus mirabilis</i>	26	24	92
<i>Staphylococcus</i> spp	75	50	67
<i>Streptococcus</i> spp	31	25	81
Mixed	146	16*	11
Total	689	496	72

*One had true mixed infection with two bacterial species present in SPA; 15 had only one species present in SPA.

Table 3 Result of white cell concentration (WCC) on initial midstream urine (MSU) compared with bacterial count of MSU and suprapubic aspiration (SPA)

MSU bacterial count ($\times 10^6/\text{l}$)	Infected WCC $\times 10^6/\text{l}$		Not infected WCC $\times 10^6/\text{l}$	
	≤ 10	> 10	≤ 10	> 10
10-100	13	11	8	13
100	241	130	27	20

Note: Bacterial counts $100 \times 10^6/\text{l}$ (SI units) = $100\ 000/\text{ml}$
WCC $10 \times 10^6/\text{l}$ (SI units) = $10/\text{cmm}$

urines were infected. In only one specimen was a true mixed infection found.

Table 3 shows the WCC of the initial MSU together with the result of culture of the SPA. When the initial MSU had $> 100 \times 10^6/\text{l}$ organisms, the WCC was raised in 65% of those in whom infection was confirmed and in 57% of those in whom the SPA was sterile. There were also 130 patients (35%) with infected SPAs but no raised WCC. In a further group of 37 patients, a sterile initial MSU was accompanied by a raised WCC. This finding was confirmed by SPA in 13, while 24 had a normal white cell concentration.

Discussion

Kass⁹ introduced the concept of quantitative urine cultures to enable the diagnosis of urinary tract infection in symptomatic and asymptomatic patients to be made with more accuracy. Since then, a concentration of organisms exceeding $100 \times 10^6/\text{l}$ in an MSU has been taken to indicate infection. In asymptomatic pregnant women, a count of $> 100 \times 10^6/\text{l}$ was reproducible on 80% of occasions.³ Kass¹ also showed that if urine contained less than $10 \times 10^6/\text{l}$ ($10\ 000/\text{ml}$) bacteria then this represented contamination. The results presented here indicate that knowledge of the identity of organisms in the urine may help in assessing the probability of the presence of a true infection. Infection was confirmed by SPA for 72% of all patients with MSU $> 100 \times 10^6/\text{l}$ bacteria. When MSUs with $> 100 \times 10^6/\text{l}$ mixed organisms were excluded SPAs confirmed infection in 88% of cases.

The technique of SPA is the simplest way to establish with certainty whether or not bacteria are present within the urinary tract.⁴ However, it is not always possible or desirable to take a bladder aspirate before starting treatment for symptomatic UTI. Only one-half of patients presenting with the symptoms generally suggestive of UTI will be found to have bacterial infection, and it is probable that those without infection should be dealt with differently

from those with infection.^{10,11} The interpretation of the initial culture of urine collected by the MSU technique is therefore important. Moreover, when an MSU is part of a routine examination, it is desirable to know the significance of an equivocal result in an asymptomatic patient.

When the MSU has $10\text{--}100 \times 10^6$ organisms/l, the results of SPAs are very different with different species of bacteria. If a mixed growth is obtained from an MSU at counts below 100×10^6 organisms/l the probability that SPA will confirm the presence of organisms is very low. When a pure culture of Gram-positive cocci at this concentration is obtained in an MSU one-third can be confirmed by SPA. However, when a pure growth of Gram-negative bacilli is obtained this may be confirmed by SPA even at counts as low as 10×10^6 /l. When the number of organisms in an MSU is above 100×10^6 /l infection is likely to be confirmed except when there is a mixed growth. It is of interest that in only one of the 903 SPAs was a true mixed infection found. The probability that a culture result indicates true infection should be taken into account when carrying out sensitivity testing of isolated organisms and also when initiating or assessing the results of treatment. It has been assumed throughout that if the MSU contains less than 10×10^6 /l bacteria then infection is very unlikely and SPA is unnecessary.

The WCC was not a useful diagnostic aid to confirm the presence or absence of UTI. It should be emphasised that most of these patients were asymptomatic and therefore not clinically comparable with patients with symptomatic UTI.

From this study it is concluded that the nature of the organism found in the urine is almost as important as the concentration of bacteria in assessing whether or not true urinary infection is present.

References

- ¹Kass E H. Asymptomatic infections of the urinary tract. *Trans Assoc Am Physicians* 1956; **69**: 56-64.
- ²Kass E H. Bacteriuria and the diagnosis of infections of the urinary tract. *Arch Intern Med* 1957; **100**: 709-714.
- ³Kass H E. Pyelonephritis and bacteriuria. A major problem in preventive medicine. *Ann Intern Med* 1962; **56**: 46-53.
- ⁴Bailey R R, Little P J. Suprapubic bladder aspiration. In: Kincaid-Smith P, Fairley K F, eds. *Renal Infection and Renal Scarring: Proceedings of an International Symposium on Pyelonephritis, Vesico-ureteric Reflux and Renal Papillary Necrosis*. Melbourne, Australia: Mercedes. 1971: 81-84.
- ⁵Mackie T J, McCartney J E. *Handbook of Bacteriology*. 10th ed. Cruickshank R, ed. Edinburgh: Livingstone. 1960: 303.
- ⁶Little P J. Urinary white-cell excretion. *Lancet* 1962; **1**: 1149-1151.
- ⁷Little P J. Diagnostic criteria of pyelonephritis. *J Clin Pathol* 1965; **18**: 556-558.
- ⁸Little P J. A comparison of the urinary white cell concentration and the white cell excretion rate. *Br J Urol* 1964; **36**: 360-363.
- ⁹Kass E H. Chemotherapeutic and antibiotic drugs in the management of infections of the urinary tract. *Am J Med* 1955; **18**: 764-781.
- ¹⁰Tapsall J W, Bell S M, Taylor P C, Smith D D. Relevance of 'significant bacteriuria' to aetiology and diagnosis of urinary-tract infection. *Lancet* 1975; **2**: 637-639.
- ¹¹Peddie B A, Little P J, Sincock A R. Urinary tract infection in general practice. *NZ med J* 1978; **88**: 317-320.

Requests for reprints to: Dr PJ Little, Department of Renal Medicine, Riyadh Military Hospital, PO Box 7897, Riyadh, Saudi Arabia.