

- Parallel tubular arrays in severe combined immunodeficiency disease: an ultrastructural study of peripheral blood lymphocytes. *Blood* 1977;**50**:55-64.
- ⁴ Payne CM, Nagle RB. Complement receptors on normal human lymphocytes containing parallel tubular arrays. *Amer J Pathol* 1980, in press.
 - ⁵ McKenna RW, Parkin J, Gajl-Peczalska KJ, Kersey JH, Brunning RC. Ultrastructural, cytochemical, and membrane surface marker characteristics of the atypical lymphocytes in infectious mononucleosis. *Blood* 1977;**50**:505-15.
 - ⁶ Lo Buglio AF, Cotran RS, Jandl JH. Red cells coated with immunoglobulin G: Binding and sphering by mononuclear cells in man. *Science* 1967;**158**:1582-5.
 - ⁷ Douglas SD, Huber H. Electron microscopic studies of human monocyte and lymphocyte interaction with immunoglobulin- and complement-coated erythrocytes. *Exp Cell Res* 1972;**70**:161-72.
 - ⁸ Elson CJ, Bradley J, Howells RE. The mechanism of rosette formation between Rh(D)-positive erythrocytes and peripheral blood lymphocytes from Rh isoimmunized individuals. The role of surface micro-projections. *Immunology* 1972;**22**:1075-86.
 - ⁹ Chen LT, Eden A, Nussenzweig V, Weiss L. Electron microscopic study of the lymphocytes capable of binding antigen-antibody-complement complexes. *Cell Immunol* 1972;**4**:279-88.
 - ¹⁰ Bentwich Z, Douglas SD, Siegal FP, Kunkel HG. Human lymphocyte-sheep erythrocyte rosette formation: some characteristics of the interaction. *Clin Immunol Immunopath* 1973;**1**:511-22.
 - ¹¹ Galey FR, Prchal JT, Amromin GD, Jhurani Y. "Hairy" B cells and "smooth" T cells. (Letter.) *N Engl J Med* 1974;**290**:690.
 - ¹² Marchalonis JJ, Bucana C, Hoyer L, Warr GW, Hanna MG, Jr. Visualization of a guinea pig T lymphocyte surface component cross-reactive with immunoglobulin. *Science* 1978;**199**:433-5.
 - ¹³ Spurr AR. A low viscosity epoxy resin embedding medium for electron microscopy. *J Ultrastruct Res* 1969;**26**:31-43.
 - ¹⁴ Kataoka K, Minowada J, Pressman D. Electron microscope study on human lymphocyte-sheep erythrocyte rosettes. *J Natl Can Inst* 1975;**55**:1323-5.
 - ¹⁵ Reyes F, Le Go A, Delrieu F, Bach JF. Ultrastructure of cells binding immunoglobulin-coated erythrocytes in rheumatoid arthritis. *Clin Exp Immunol* 1974;**17**:533-46.
 - ¹⁶ Lay WH, Nussenzweig V. Receptors for complement on leukocytes. *J Exp Med* 1968;**128**:991-1009.
 - ¹⁷ Bell GH, Davidson JN, Scarborough H. Textbook of Physiology and Biochemistry, 5th ed. Edinburgh: Livingstone. Baltimore: The Williams and Wilkins Company, 1961.
 - ¹⁸ Abramson N, Lo Buglio AF, Jandl JH, Cotran RS. The interaction between human monocytes and red cells. Binding characteristics. *J Exp Med* 1970;**132**:1191-206.
 - ¹⁹ Anderson N, Doane FW. Epoxy embedding of thin-layer material in commercially available vinyl cups for light and electron microscopy. *Stain Tech* 1967;**42**:169-73.

Requests for reprints to: Dr Claire M Payne, Department of Pathology, University of Arizona Health Sciences Center, Tucson, Arizona 85724, USA.

Letters to the Editor

Use of Intralactam for detection of β -lactamase production by *Neisseria gonorrhoeae*

In your Journal of July 1979 (p 738), DB Wheldon and Mary PE Slack reported an evaluation of Intralactam for the detection of β -lactamase production by *Haemophilus influenzae*. At the end of their paper they commented that it might also be used to detect gonococcal β -lactamase.

We have used this acidometric method to detect β -lactamase production in primary isolates of *Neisseria gonorrhoeae* for the past year. Initially we prepared paper strips impregnated with penicillin and indicator, as described by Slack *et al.*,¹ but then began using the commercially available Intralactam.

All confirmed isolates of *N. gonorrhoeae* have minimum inhibitory concentrations (MIC) determined to penicillin, spectinomycin, and tetracycline. At first, testing for β -lactamase production was limited to isolates having an MIC to penicillin of > 2.0 mg/l. (Exact quantification of the MIC value above 2 mg/l was not performed.) Of 2357 *N. gonorrhoeae* isolated in this laboratory from clinical specimens in 1978-79, 38 organisms (1.6%) had an MIC to penicillin of > 2.0 mg/l. These 38 organisms had a β -lactamase determination performed, the majority using Intralactam strips, but a small number using our own prepared strips with penicillin and neutral red incorporated, as described by Slack *et al.*¹ Of these 38 organisms, 36 were identified as

β -lactamase producers. However, as MIC results and identification of β -lactamase production were not usually available to the clinicians until four days after primary isolation, we decided to perform β -lactamase determination on all isolates using Intralactam strips at the same time as a presumptive diagnosis of *N. gonorrhoeae* was made.

We now provide clinicians with a presumptive diagnosis of *N. gonorrhoeae* if oxidase-positive, Gram-negative diplococci are present on primary isolation plates after 18-24 hours' incubation, and at the same time report the presence or absence of β -lactamase production. The primary isolation medium used is modified New York City Medium.²

We have used Intralactam strips to

Letters to the Editor

determine β -lactamase production in 267 presumptive gonococcal isolates. Seven organisms had MICs to penicillin of > 2.0 mg/l, the remainder having MICs to penicillin of 0.03-1.0 mg/l. The Intralactam test identified the seven organisms with MICs of > 2.0 mg/l as β -lactamase producers, and none of the other 260 isolates gave a positive result with the test.

Due to Australia's proximity to south-east Asia, where β -lactamase producing *N. gonorrhoeae* strains are frequently encountered, we occasionally see such organisms in patients returning from these countries. There have also been some β -lactamase producing strains causing infections contracted locally. The standard treatment at the Venereal Diseases Clinic in Sydney is amoxycillin, 3 g orally, preceded by probenecid. This is highly effective except in the case of β -lactamase producing strains. For this reason we feel that rapid detection of β -lactamase production is desirable so that antibiotic treatment can be modified accordingly. In our experience, Intralactam strips provide a rapid, simple method for detecting β -lactamase production in *N. gonorrhoeae*.

R MUNRO
R MALLON
D DALEY

Department of Bacteriology,
Institute of Clinical Pathology
and Medical Research,
Sydney,
New South Wales,
Australia

References

- Slack MPE, Wheldon DB, Turk DC. A rapid test for β -lactamase production by *Haemophilus influenzae*. *Lancet* 1977; 2:906.
- Young H. Cultural diagnosis of gonorrhoea with Modified New York City (MNYC) medium. *Br J Ven Dis* 1978;54:36-40.

A human strain of *Campylobacter fetus* subsp. *intestinalis* grown at 42°C

A recent report of an infection with *Campylobacter fetus* subsp. *intestinalis* in a patient with multiple myeloma¹ leads us to mention several unusual features of the case. The strain was first isolated from a presumably traumatic joint. A blood culture was requested following that isolation. Tryptic Soy Broth and Thio-glycollate Medium (w/o Indicator) bottles (Pfizer Inc), inoculated with 5 ml blood each, remained clear (both contained

12 v/v % CO₂) but the organism grew in a one-day subculture to chocolate agar at 37°C in 8% CO₂. Most remarkable, however, was the fact that the *C. fetus* isolate grew at both 25 and 42°C. Since it proved to be resistant to nalidixic acid (no zone around the 30 μ g disc) and grew in 1% glycine broth the identification of *C. fetus* subsp. *intestinalis* was established.²

Such strains have previously been reported from ovine sources³ but not from man. Conventional wisdom has it that the subspecies *intestinalis* and *fetus* (so far never isolated from man) grow at 25°C but not at 42°C while the subspecies *jejuni* grows at 42°C but not at 25°C. Our finding prompts us to re-emphasise that growth at 25°C is a more reliable characteristic in the differential diagnosis of *C. fetus* than growth at 42°C.² For the differentiation of clinical strains, we therefore recommend growth at 25°C and susceptibility to nalidixic acid rather than growth at both 25°C and 42°C.

We also want to emphasise the importance of 'blind' subcultures of blood culture bottles to suitable media, such as chocolate agar in a CO₂ atmosphere. It is rare to see turbidity due to *C. fetus* in blood culture media, presumably because of an inadequate growth atmosphere in commercial bottles.²

ROBERT M SMIBERT
ALEXANDER VON GRAEVENITZ
Anaerobe Laboratory, Virginia
Polytechnic Institute and State University,
Blacksburg, Virginia 24061; and
Clinical Microbiology Laboratories,
Yale-New Haven Hospital, New Haven,
Connecticut 06504, USA

References

- Fick R, Isturiz R, Cadman E. *Campylobacter fetus* septic arthritis. Report of a case. *Yale J Biol Med* 1979;52:339-44.
- Smibert RM. The genus *Campylobacter*. *Annu Rev Microbiol* 1978;32:673-709.
- Firehammer BD. The use of temperature tolerance in the identification of *Vibrio fetus*. *Amer J Vet Res* 1965;26:995-7.

Is there a place for the GammaCoat (125I) Gentamicin Radioimmunoassay Kit in a routine clinical pathology laboratory?

Waterworth¹ has pointed out that a delay in obtaining results detracts from the value of a gentamicin assay service. In an attempt to minimise this delay a variety

of rapid methods have been devised. All of these have their protagonists, but none has yet achieved general acceptance. Included among these rapid methods is radioimmunoassay in a number of forms.²⁻⁵ Radioimmunoassay techniques are simple, sensitive, and specific.² Thus, when a kit, the GammaCoat (125I) Gentamicin Radioimmunoassay Kit (Clinical Assays Cat. No. CA-540), recently became available in this country, we made a preliminary assessment of it.

Paired (pre- and post-dose) sera from patients receiving gentamicin therapy, either alone or in combination with other antibiotics, were assayed by our current overnight plate diffusion method and then, together with standards used to calibrate this method and controls of varying values made up in the laboratory, reassayed blind using the GammaCoat (RIA) kit. The principle of the GammaCoat technique and the recommended procedure for use of the kit are detailed in the instruction booklet. We followed the recommendations except that, for reasons of economy in the trial, only one tube per standard, control, or sample was used.

Altogether 109 samples were analysed by both methods. These comprised 15 standards, 12 controls, and 41 pairs of sera. The correlation between the results obtained by RIA and plate diffusion are shown in Figures 1 and 2. The linear correlation coefficient (r) for the standard/control results was +0.963 and for the patients' sera was +0.961. A slightly better correlation was obtained using a quadratic model (parabola) in which y (value by RIA method) = $-1.22 + 1.49x - 0.038x^2$ (where x = value by plate diffusion method) for the standard/control results and $y = -0.23 + 1.21x - 0.0257x^2$ for the patients' sera. Discrepancies seemed to be most marked at higher levels. Mahon *et al.*³ commented on discrepancies between the results by the two methods in their study and suggested that it might be due to the effects of multiple antibiotic therapy. This did not seem to be so in our case, other antibiotics present being inactive against our plate diffusion assay indicator organism, and a similar correlation being found for both patients' samples and standard/control samples. The RIA method may have been reading low at the higher levels, but we cannot explain why at present. An assessment of each pair of sera was made retrospectively by one of us (CDR), and it was found the variations