Diagnosis of hairy-cell leukaemia by tartrate-resistant acid phosphatase activity in paraffin-embedded tissue sections

V GROULS

From the Institute of Pathology of the University of Bonn, Germany

SUMMARY Tartrate-resistant acid phosphatase activity may be demonstrated in paraffin-embedded liver and spleen specimens as well as in decalcified bone marrow biopsies after fixation in a mixed glutaraldehyde-formaldehyde-calcium acetate solution. This technique may routinely be applied to haematologically relevant material and aids in the differential diagnosis of hairy-cell leukaemia.

Jamshidi needle biopsies of the bone marrow are used extensively in the initial evaluation of patients with non-Hodgkin's lymphoma. Hairy-cell leukaemia has to be considered in the differential diagnosis of lymphoid myelofibrosis. A positive tartrate-resistant acid phosphatase activity of the cells usually establishes the diagnosis. In the following short

communication the possibility of employing this marker enzyme also on paraffin-embedded decalcified bone marrow sections and other tissues is discussed.

Methods

Needle biopsy specimens of bone marrow from the iliac crest as well as liver and spleen tissues were

Received for publication 3 December 1979

Fig. 1 Positive tartrate-resistant acid phosphatase activity in lymphoid cells. Decalcified, paraffin-embedded bone marrow; Jamshidi needle biopsy, no nuclear counterstaining $\times 1300$.

Fig. 2 Infiltration of a portal tract in hairy-cell leukaemia. Paraffin-embedded wedge biopsy of liver; positive cytoplasmic staining with tartrate-resistant acid phosphatase; nuclear counterstaining $\times 400$.

fixed in a solution containing 20 ml 25% glutaraldehyde, 40 ml 35% formaldehyde, and 15.8 g calcium-acetate, made up to 1000 ml with distilled water.¹ Osseous structures were decalcified in a solution of 100 g EDTA and 33 g Tris buffer with distilled water added to 1000 ml at room temperature for two to three days.²

The tartrate-resistant acid phosphatase reaction was carried out using naphthol-AS-BI-phosphate as a substrate and hexazonium pararosaniline as a coupler.³

Results and comment

Three cases of suspected hairy-cell leukaemia were investigated. In the first case, a Jamshidi bone marrow biopsy (Fig. 1), in the second a wedge excision of the liver (Fig. 2) as well as the removed spleen, and in the third the spleen only were examined. In every instance the typical diffuse bright red enzyme activity in the cytoplasm of the hairy cells is apparent, thus confirming the tricholeucocytic nature of these malignant lymphomas. An improved staining contrast will sometimes be obtained, especially in bone marrow biopsies without nuclear counterstaining. Furthermore, it is shown in Fig. 2 that the lymphoid infiltrate is not only located in the liver sinusoids but also to a great extent in the

portal tracts. In the bone marrow, macrophages and osteoclasts exhibit a distinct positive reaction, but this may not pose any diagnostic problems as far as hairy-cell leukaemia is concerned.

From the practical point of view it may be stated that, besides AS-D-chloroacetate-esterase⁴ and peroxidase,⁵ acid phosphatase is the third cellular enzyme which may easily be applied in the examination of haematologically relevant paraffin-embedded tissue material, and thus be used successfully in routine work. In suitable cases with a high enzyme content, it surely will be helpful in making the final diagnosis of hairy-cell leukaemia. But it should also be mentioned that both the fixation and decalcification of material in the solutions originally recommended by Schaefer^{1,2} are necessary preconditions for an optimal staining result.

References

- ¹ Schaefer HE. 1978, personal communication.
- ² Schaefer HE, Hellriegel KP, Fischer R. Vorkommen von Tartrat-resistenter saurer Phosphatase in verschiedenen Zelltypen des lympho-retikulären und hämatopoetischen Systems. *Blut* 1977;**34**:393-7.
- ³ Leder LD, Stutte HJ. Seminar für hämatologisch-zytochemische Techniken. *Verh dtsch Ges Pathol* 1975;**59**:503-9.

- ⁴ Leder LD. Der Nachweis von Naphthol-AS-D-Chloracetat Esterase und seine Bedeutung für die histologische Diagnostik. *Verh dtsch Ges Pathol* 1964;**48**:317-20.
- ⁵ Schaefer HE, Fischer R. Der Peroxydasenachweis an Ausstrich-präparaten sowie an Gewebsschnitten nach Entkalkung und Paraffineinbettung. *Klin Wschr* 1968;**46**:1228-30.
- Requests for reprints to: Dr V Grouls, Pathological Institute, University of Bonn, 53 Bonn 1, Germany.