Table 1 Isolation rate of group B streptococcus from faeces

Source	No positive/No examined (%)		
	Total specimens	Fresh specimens	Frozen specimens
Males	11/108 (10.2)	8/48 (16.7)	3/60 (5)
Females	6/67 (9)	5/33 (15.2)	1/34 (2.9)
Total	17/175 (9.7)	13/81 (16)	4/49 (4.3)

Table 2 Distribution of group B streptococcal serotypes in faecal isolates

No of faecal carriers	Serotype distributions							
	Ia	Ib	Ic-protein	II	III/Ic-protein	III	X	NT
Adult female	5	1	1	0	1	0	1	0
Female <16 yr	1	1	0	0	0	0	0	0
Adult male	10	0	0	4	0	1	3	1
Male <16 yr	1	1	0	0	0	0	0	0
Total	17	3	1	4	1	1	4	1

report of GBS serotypes in faecal samples is that of Islam and Thomas¹ who detected four type II strains in their series and suggested that type II may be the only resident gut flora in adults. Type II is an uncommon cause of serious neonatal infection, but serotypes I and III are much more frequently implicated (unpublished, PHLS Commun Dis Rep).

The isolation rate of GBS from both rectal and perianal swabs is much higher than from faeces.³ It may be that carriage is mainly limited to the perianal skin, which could contaminate rectal swabs,¹ or alternatively, the rectum and anal canal may also be colonised.³ If the latter is true, it is surprising that faeces are not more often contaminated. In either case, it is impossible to be certain that GBS are gut commensals without examining stool specimens collected in such a way as to avoid contamination—for example, by sigmoidoscopy. Further studies are required.

CJ NOBLE
Maidstone Public Health Laboratory,
Preston Hall Hospital,
Maidstone, Kent ME20 7NH

References

¹ Islam AKMS, Thomas E. Faecal carriage of group B streptococci. *J Clin Pathol* 1980;33:1006-8.
² Islam AKMS. Rapid recognition of group B streptococci. *Lancet* 1977;i:256-7.
³ Easmon CSF, Tanna A, Munday P, Dawson S. Group B streptococci—gastrointestinal organisms? *J Clin Pathol* 1981;34:921-3.

An evaluation of API-20 STREP and isolation of *Streptococcus milleri*

The API-20 STREP system has been under

evaluation in this laboratory for several months, and our results largely agree with those reported by Tillotson.¹ The system is quick and reliable.

My particular interest has been infections due to *Streptococcus milleri*, and I can confirm that the API-20 STREP system enables the rapid identification of this organism. The Table shows the 15 isolations made from clinically significant lesions in a four-month period, in Glasgow Royal Infirmary. In 14 cases, *Strep milleri* was grown in pure culture or as the predominant organism from a purulent lesion, and the remaining isolate was from a blood culture, though not associated with endocarditis. Peritoneal and pelvic sepsis accounted for most of our isolates. There were no cases of brain abscess because such patients are not treated in Glasgow Royal Infirmary.

Rapid diagnosis is important, and, since the organism responds readily to penicillin, prompt adequate antibiotic treatment together with surgical drainage (where appropriate) can achieve an excellent clinical response. Untreated, the organism is responsible for considerable morbidity associated with failure of healing of surgical wounds.

Isolations from significant lesions

Site/lesion	No of isolates	Comments
Peritonitis	5	4 post-op bowel surgery 1 appendicitis
Pelvic sepsis	4	3 post-hysterectomy 1 pelvic inflammatory disease
Pancreatic pseudocyst	1	drainage fluid
Head and neck	2	1 post-op skin flap-reconstruction 1 spontaneous abscess
Empyema of pleura	1	
Skin ulcer	1	
Blood culture	1	bacteraemia after bowel surgery

In my hands, the API-20 STREP system is of real value in the busy routine diagnostic laboratory as a rapid and reliable method of identifying streptococcal species; it is particularly useful with the difficult "indifferent" streptococci, such as *Strep milleri*.

MORAG I BROWN
Department of Bacteriology,
Royal Infirmary,
Glasgow G4 0SF

Reference

¹ Tillotson GS. An evaluation of the API-20 STREP system. *J Clin Pathol* 1982;35:468-72.

Quality control terminology and some practical implications

The International Federation of Clinical Chemistry's (IFCC) approved recommendations on general principles and terminology on quality control¹ has resolved several outstanding issues and significant among these appears to be the affirmation that the negative forms of the terms *accuracy* and *precision* should be used to describe the quantification of these parameters.

Though the measurement of *imprecision* never posed a problem, the generally accepted definition of accuracy—that of agreement of results with the *true value*—and the premise that the true value was an unknowable entity,^{2,3} have resisted efforts at quantifying it directly. Recovery experiments hence became the generally acknowledged means of assessing *inaccuracy* and the IFCC has accepted this.

Whitehead⁴ circumvented the rigidity of the aforementioned definition of accuracy by introducing a broader one—namely, the relationship of a set of results to the *correct value*, where the correct value is the best estimate of a quantity for a particular material using a defined analytical technique. As values can vary widely depending on the method used, this definition runs