Inhibitory effects of peripheral blood cells on in vitro colony formation by autologous bone marrow in aplastic anaemia: relation with response to immunosuppressive therapy

A BLASETTI, A FAILLE, N BALITRAND, E GLUCKMAN, A DEVERGIE, C DRESCH

From the Service Central de Médecine Nucleaire, Hôpital Saint-Louis, 2 Place du Docteur Fournier, 75475 Paris Cedex 10, France

SUMMARY The inhibitory activity of peripheral blood lymphocytes on autologous bone marrow was studied in 27 patients with aplastic anaemia after treatment with androgens. Inhibitory activity was hard to assess in 10 patients studied during the first year of treatment. The colony count was too low to be certain of differences between the samples incubated with or without lymphocytes. Among the 17 patients who had more than 10 colonies per 2×10^5 mononuclear bone marrow cells, nine showed inhibitory activity by peripheral blood lymphocytes. After 12 months of androgen therapy each of these patients showing inhibitory activity of bone marrow colony forming cells by peripheral lymphocytes responded to antithymocyte globulin. None of nine patients with few colony forming cells or no inhibitory activity of lymphocytes responded to immunosuppression.

Aplastic anaemia is not a single entity but a heterogeneous group of pancytopenias associated with bone marrow hypoplasia. The physiopathological basis of the condition remains unclear so therapeutic approaches are pragmatic. Spontaneous bone marrow recovery in some cases indicates that a residual population of stem cells may be able to repopulate the bone marrow.¹ The fact that some autologous reconstitution may follow androgen therapy and also immunosuppressive therapy may indicate the existence of a population of immune cells involved in the pathogenesis of aplastic anaemia.²⁻⁴ In vitro bone marrow cultures have shown that pre-incubation with antithymocyte globulin (ATG) may increase granulomonocyte colony forming cells (GM-CFC) and suggests the involvement of lymphocytes in the pathogenesis.⁵⁻⁷ Decreased colony formation by normal GM-CFC has been detected in the presence of peripheral blood cells from patients with aplastic anaemia.⁷⁻⁹ However these studies were performed first in an allogenic situation and the possibility of an HLA sensitisation due to multiple transfusion was suggested.^{10 11} More recent studies have been conducted in autologous situations.¹² However due to the scarcity of remaining stem cells in the early phase

of aplastic anaemia, these studies have to be performed after treatment for some time. We present 27 coculture studies of autologous bone marrow and lymphocytes performed two months to 10 yr after the beginning of androgen therapy. Inhibitory activity of lymphocytes could be assessed only after one year of treatment. There was a good correlation between this activity and the response to immunosuppression.

Patients and methods

PATIENTS

Clinical laboratory data are summarised in Tables 1 and 2. Seventeen patients met the criteria for severe aplastic anaemia at the onset of the disease³ (platelets $< 20 \times 10^9/l$; reticulocytes $< 20 \times 10^9/l$; polymorphonuclears $< 2 \times 10^9/l$) with an "empty" bone marrow smear. The 10 remaining patients had pancytopenia with a hypocellular but not empty bone marrow smear.

Ten of the patients with severe aplasia (Table 1) were studied between two and 12 months after the onset of disease. Seventeen patients were studied one to ten years after the onset of the disease (Table 2): seven with severe aplasia (cases 11 to 17) and 10 with pancytopenia (cases 18 to 27).

Table 1 *BM-GM-CFC and % variations with autologous peripheral blood mononuclear leucocytes (PBML) in short term follow-up patients: relation to response to immunosuppressive therapy*

Case No/ sex/age (yr)	Duration of disease before study (months) and therapy		BM-GM-CFC / 2×10^5 cells	Colonies in coculture/colonies in control culture			Response to immunosuppressive therapy
				+PBML	+E rosette + cells	+E rosette – cells	
1 F 18	2	Androgens	3	1	0.67	1.3	NT
2 F 22	3	Androgens	0	0	0	0	NR (ATG)
3 F 16	3	Androgens	2	1	3	1	NT
4 M 14	3	Corticoids	3	1.5	—	—	NR (corticoids)
5 F 15	4	Androgens	0	0	0	0	NT
6 M 18	4	Androgens	0	0	0	0	NR (corticoids)
7 M 41	6	Androgens	9	1.6	—	—	NR (ATG)
8 M 21	8	Androgens	3	1	1	1.5	NR (ATG)
9 M 15	8	Androgens	7	1.40	—	—	NT
10 M 12	11	Androgens	5	0.80	1.2	1	NR (ATG)

BM-GM-CFC = bone marrow granulomonocyte colony forming cells.

NT = not tested.

NR = non-responders.

ATG = antithymocyte globulin.

METHODS

Rosetting procedures

Peripheral blood mononuclear leucocytes (PBML) were obtained by Ficoll-Hypaque (Lymphoprep, Nyegaard D = 1.077 g/cm^3) separation of heparinised whole blood diluted twice with Hanks' solution (HBSS). Cells were then incubated with sheep red blood cells sensitised by 2-aminoethylisothiuronium bromide hydrobromide (AET) (Sigma) to produce rosette formation.¹³ The cell suspension was then centrifuged over Ficoll-Hypaque. E rosette-negative cells were taken up from the interphase,

resuspended in HBSS and washed twice in HBSS. The pellet of E rosette-positive cells was processed with a hypotonic solution to disrupt the rosettes and harvest E rosette-positive cells. The E rosette-positive cells were then washed twice.

Coculture studies

Bone marrow cells were separated in the same way as blood cells on a Ficoll Hypaque density gradient and the supernatant cells were harvested, resuspended and washed in McCoy's 5 A medium. Aliquots of 6×10^5 nucleated cells in 0.1 ml were pipetted into sterile

Table 2 *BM-GM-CFC and % variations with autologous PBML in long term follow-up patients: relation to response to immunosuppressive therapy*

Case No/ sex/age (yr)	Duration of disease before study (months) and therapy		BM-GM-CFC /2 × 10 ⁵ cells	Colonies in coculture/colonies in control culture			Response to immunosuppressive therapy
				+ PBML	+ E rosette + cells	+ E rosette – cells	
Severe aplasias							
11 M 50	12	Androgens	21	0.43	0.77	0.14	R (ATG)
12 M 15	16	Androgens	13	1.3	1	0.67	NT
13 F 35	30	Androgens	14	0.86	1.43	0.86	NR (ATG)
14 M 27	36	Androgens	13	0.40	0.50	0.50	R (ATG)
15 F 54	37	Androgens	34	0.40	—	—	R (ATG)
16 M 31	36	Androgens	48	0.67	0.45	0.37	R (ATG)
17 F 18	36	Androgens	10	1.45	—	—	NT
Hypoplasias							
18 M 32	12	Androgens	12	0.36	0.58	0.75	R (ATG)
19 M 23	12	—	71	1	0.61	0.29	R (ATG)
20 M 21	15	Androgens	45	1.04	0.98	0.87	NT
21 F 17	20	Androgens	18	0.50	0.50	0.11	R (ATG)
22 M 15	20	Androgens	23	2	1.7	4	NT
23 F 39	22	Androgens	28	0.82	0.68	0.57	R (ATG)
24 M 33	22	Androgens	10	0.50	0.60	0.60	R (ATG)
25 F 39	24	Androgens	10	0.67	1.17	1	NT
26 F 51	25	Androgens	13	1	0.92	0.85	NR (ATG)
27 F 25	120	Androgens	16	0.68	0.81	1.22	NR (corticoids)

8 mm plastic tubes with or without 6×10^5 PBML or 6×10^5 peripheral E rosette-positive cells or 6×10^5 E rosette-negative cells. The cells were incubated at 37° C for 2 h in 5% CO₂. The cells were then centrifuged and the whole cellular sediment was resuspended in 0.3% agar and medium containing 15% fetal calf serum and plated in triplicate 1 ml volume with 0.1 ml of placenta conditioned medium.¹⁴ After 10 days of incubation at 37° C in a fully humidified atmosphere containing 5% CO₂, the colonies (groups containing 50 cells or more) were counted.

Results

When tested in 14 normal subjects, normal PBML peripheral E rosette-positive or E rosette-negative cells were not inhibitory for autologous bone marrow GM-CFC. Mean values were as follows: bone marrow + PBML = 1.02 ± 0.22 ($\pm$ SD); bone marrow + E rosette-positive cells = 1.05 ± 0.29 ($\pm$ SD); bone marrow + E rosette-negative cells = 1.02 ± 0.30 ($\pm$ SD) compared to bone marrow cells incubated without PBML. Lymphocytes were considered inhibitory when colonies in cocultures were less than 0.6 of control culture.

Results for aplastic patients are summarised in Tables 1, 2, and 3.

Ten patients studied between 2 and 11 months after the onset of disease had received androgen therapy for two months or more (except case 4 who received only corticosteroids for two months). Their granulocyte count was still below $1 \times 10^9/l$ and platelet count below $35 \times 10^9/l$; bone marrow cellularity was much reduced in all of them and all needed blood support. In every case GM-CFC values were too low to allow any valid study; in the seven cases with few colonies no inhibitory activity of PBML could be seen. There was no response to immunosuppressive therapy in the six cases where this therapy was given (two cases with no colony growth and four cases with rare colonies).

All the 17 patients studied more than one year after the onset of the disease had received androgen therapy, except case 19 who had post-hepatic

aplastic anaemia. All had more than 10 colonies 2×10^5 bone marrow mononuclear cells; cases 16, 19, 20 were in the normal range. No correlation was found between the number of colonies and the severity of the disease at onset or the clinical data at the time of the study. Four of seven aplasias and 5/10 hypoplasias had suppressor cells. Inhibitory activity was found in both E rosette-positive and E rosette-negative cells; however in three cases E rosette-negative cells were much more inhibitory than E rosette-positive cells (cases 11, 19, 21).

Immunosuppressive therapy was given in 11 cases using ATG and in one case using corticosteroids. All nine patients who displayed inhibitory activity responded well (cases 11, 14, 15, 16, 18, 19, 21, 23, 24) with an increase in granulocytes $> 1 \times 10^9/l$ and no further requirement for blood support within a month of immunosuppression. Among the eight patients without inhibitory activity at the time of the study, three received immunosuppressive therapy and failed to respond (case 13, 26, 27).

Discussion

Reports of bone marrow recovery after unsuccessful allogeneic bone marrow graft or after treatment with high doses of cyclophosphamide or ATG^{15 16} suggest that an immunological mechanism is involved in some cases of aplastic anaemia. Therefore it is important to try to identify patients who might benefit from immunosuppressive therapy. Coculture studies in an autologous system are impossible to perform at the onset of the disease due to the scarcity of the remaining bone marrow cells. Coculture studies in an allogeneic system have been criticised on the grounds that inhibition could be related to transfusion.^{10 11} Our results show that it is useless to try, by autologous coculture, to identify patients who might benefit from immunosuppression at an early stage of the disease. On the one hand results of coculture are unreliable when the number of colonies is too low; on the other hand none of the six patients treated in the first year of the study responded to immunosuppression.

If one supposes that stem cell failure may be

Table 3 Coculture results: relation to response to immunosuppressive therapy

Patients	Invitro inhibition	Immunosuppressive therapy	
		Patients tested	Response
Short term follow up patients = 10	No growth 3	2	2 non-responders
	No inhibition 7	4	4 non-responders
Long term follow up patients = 17	Inhibition 9	9	9 responders
	No inhibition 8	3	3 non-responders
		5 (not tested)	

accompanied in some cases by a dysfunction of the immune system, evidence of this dysfunction could only be obtained after several months of androgen-therapy, when this treatment has allowed multiplication of the remaining stem cells. Our results show that a treatment of at least 12 months is necessary to increase the CFC concentration to levels permitting valuable coculture studies.

After one year or more of androgen-therapy autologous coculture showed a residual autoimmune process in half the patients studied. As in other studies¹² we did not find in most cases a predominance of inhibiting activity in the E rosette-positive lymphocyte population. This could be due to the fact that the suppressor activity is shared by subpopulations of cells present in both E-positive and E-negative cells or by monocyte-macrophages, which would be present in the E-negative cells, as suggested by Suda.¹⁷ Recent studies of normal subjects seem to indicate that a subset of null lymphocytes may normally be inhibitory against bone marrow GM-CFC.¹⁸ Bacigalupo and coworkers¹⁹ found that E rosette-positive bone marrow cells are not normally inhibitory, but may become so if previously activated by pokeweed mitogen. PHA and Con-A have also been shown to be capable of inducing suppressor activity in T lymphocytes.²⁰ Thus the term "auto-immune inhibition" in aplastic anaemia may be incorrect and it would be more appropriate to speak of "dysfunction" of the lymphocyte and monocyte populations that normally control haematopoiesis.

It appears that lymphomonocyte dysfunction persists in about half of the patients with prolonged aplasia. Coculture inhibition of GM-CFC in these cases correlated with their responsiveness to immunosuppressive therapy. Further study of the links between immune cells and haematopoietic stem cells in pancytopenias would therefore seem to be worthwhile in defining new approaches to treatment.

References

- Speck B, Cornu P, Jeannet M, *et al.* Autologous bone marrow recovery following allogeneic bone marrow transplantation in a patient with severe aplastic anaemia. *Exp Hematol* 1976;4:131-7.
- Jeannet M, Speck B, Rubinstein A, Pelet B, Wyss M, Kummer H. Autologous marrow reconstitution in severe aplastic anaemia after ALG pretreatment and HLA semicompatible bone marrow cell transfusion. *Acta Hematol* 1976;55:129-39.
- Gluckmann E, Devergie A, Faille A, *et al.* Treatment of severe aplastic anaemia with antilymphocyte globulin. *Exp Hematol* 1978;6:679-87.
- Baran DT, Griner PF, Klemperer MR. Recovery from aplastic anaemia after treatment with cyclophosphamide. *N Engl J Med* 1976;295:1522-9.
- Faille A, Barrett AJ, Balitrand N, Ketels F, Gluckmann E, Najean Y. Effect of antilymphocyte globulin on granulocyte precursors in aplastic anaemia. *Br J Haematol* 1979;42:371-80.
- Ascensao JA, Pahwa R, Kagan W, Hansen JA, Moore MAS, Good RA. Aplastic anaemia: evidence for an immunological mechanism. *Lancet* 1976;i:669-71.
- Kagan WA, Ascensao JA, Pahwar N, *et al.* Aplastic anaemia: presence in human bone marrow of cells that suppress myelopoiesis. *Proc Natl Acad Sci USA* 1976;73:2890-4.
- Singer JW, Brown JE, James MC, *et al.* Effect of peripheral blood lymphocytes from patients with aplastic anaemia on granulocyte colony growth from HLA matched and mismatched marrows: effect of transfusion sensitisation. *Blood* 1978;52:37-46.
- Hoffman R, Zanjani ED, Lutton JD, Zalusky R, Wasserman LR. Suppression of erythroid colony formation by lymphocytes from patients with anemia. *N Engl J Med* 1977;296:19-27.
- Singer JW, Doney KC, Thomas ED. Coculture studies of 16 untransfused patients with aplastic anaemia. *Blood* 1979;54:180-5.
- Török-Storb BJ, Sieff C, Storb R, Adamson J, Thomas ED. In vitro tests for distinguishing possible immune mediated aplastic anaemia from transfusion induced sensitization. *Blood* 1980;55:211-5.
- Nissen C, Cornu P, Gratwohl A, Speck B. Peripheral blood cells from patients with aplastic anaemia in partial remission suppress growth of their own bone marrow precursors in culture. *Br J Haematol* 1980;45:233-43.
- Hockland P, Hockland M, Heron I. An improved technique for obtaining E rosettes with human lymphocytes and its use for B cell purification. *J Immunol Methods* 1976;13:175-80.
- Pike B, Robinson WA. Human bone-marrow growth in agar gel. *J Cell Phys* 1970;76:78-80.
- Territo MC. For the UCLA bone-marrow transplantation team. Autologous bone-marrow repopulation following high dose cyclophosphamide and allogeneic marrow transplantation in aplastic anaemia. *Br J Haematol* 1977;36:305-12.
- Thomas ED, Storb R, Giblett ER, *et al.* Recovery from aplastic anaemia following attempted bone-marrow transplantation. *Exp Hematol* 1976;4:97-102.
- Suda T, Misoguchi H, Miura Y, Kubota K, Takaku F. Enhancement of granulocytic colony formation by deletion of phagocytic cells in the bone-marrow of patients with idiopathic aplastic anaemia. *Exp Hematol* 1980;8:659-64.
- Morris TCM, Vincent PC, Sutherland R, Hersey P. Inhibition of normal human granulopoiesis in vitro by non B non T lymphocytes. *Br J Haematol* 1980;45:541-50.
- Bacigalupo A, Podesta M, Mingari MC, *et al.* Generation of CFU-C suppressor T cells in vitro. An experimental model for immune mediated marrow failure. *Blood* 1981;57:491-6.
- Banisadre, Ascensao JL, Kay NE, Ash RC, Zanjani ED. PHA induced suppression of human blood BFU-E is mediated by T cells. *Exp Hematol* 1980;8 (suppl 7):44 (Abstract).

Requests for reprints to: Dr A Faille, Service du Professeur Y Najean INSERM U 204, Centre Hayem, Hôpital Saint-Louis, 75475 Paris Cedex 10, France.