

We thank Mr J Reid, Mr J McDermott, and Mr W Kirkwood for the histological staining and S Campbell for typing the paper. RAJ Spence was in receipt of an Eastern Health and Social Services Board research fellowship during this study.

#### References

- <sup>1</sup> Tompsett DH. *Anatomical techniques*. 2nd ed. London: Churchill Livingstone, 1970:9-27.
- <sup>2</sup> Butler H. The veins of the oesophagus. *Thorax* 1951;**6**:276-96.
- <sup>3</sup> Hales MR, Carrington CB. A pigmented gelatin mass for vascular injection. *Yale J Biol Med* 1971;**43**:257-70.
- <sup>4</sup> Reynolds DG, Swan KG. Intestinal microvasculature architecture in endotoxic shock. *Gastroenterology* 1972;**63**:601-20.
- <sup>5</sup> Drury RAB, Wallington EA. *Carleton's histological technique*. 4th ed. Oxford: Oxford University Press, 1967:167.
- <sup>6</sup> Murakami T. Application of the scanning electron microscope to the study of the fine distribution of the blood vessels. *Arch Histol Jap* 1971;**32**:445-54.
- <sup>7</sup> Northover JMA, Williams EDF, Terblanche J. The investigation of small vessel anatomy by scanning electron microscopy of resin casts. *J Anat* 1980;**130**:43-54.
- <sup>8</sup> Dollinger RK, Armstrong PB. Scanning electron microscopy of injection replicas of the chick embryo circulatory system. *J Microscopy* 1974;**102**:179-86.
- <sup>9</sup> Page RB, Munger BL, Bergland RM. Scanning microscopy of pituitary vascular casts. *Am J Anat* 1976;**146**:273-302.
- <sup>10</sup> Manenti F, Williams R. Injection studies of the splenic vasculature in portal hypertension. *Gut* 1966;**7**:175-80.
- <sup>11</sup> Greenway CV, Murthy VS. Effects of vasopressin and isoprenaline infusions on the distribution of blood flow in the intestine: criteria for the validity of microsphere studies. *Br J Pharmacol* 1972;**46**:177-88.
- <sup>12</sup> Micflikier AB, Bond JH, Sircar B, Levitt MD. Intestinal villus blood flow measured with carbon monoxide and microspheres. *Am J Physiol* 1976;**230**:916-9.
- <sup>13</sup> Grandison AS, Yates J, Shields R. Capillary blood flow in the canine colon and other organs at normal and raised portal pressure. *Gut* 1981;**22**:223-7.
- <sup>14</sup> Anderson JM. Histometry. In: Bancroft JD, Stevens A, eds. *Theory and practice of histological techniques*. 2nd ed. Edinburgh: Churchill Livingstone, 1982:548-63.
- <sup>15</sup> Aberne WA, Dunnhill MS. *Morphometry*. 1st ed. London: Edward Arnold, 1982:46-118.
- <sup>16</sup> Bradbury S. Commercial image analysers and the characterisation of microscopical images. *J Microscopy* 1983;**131**:203-10.
- <sup>17</sup> Laursen H. Cerebral vessels and glial cells in liver disease: a morphometric and electron microscope investigation. *Acta Neurol Scand* 1982;**65**:381-412.
- <sup>18</sup> Wilkinson JH, Hopewell JW, Reinhold HS. A quantitative study of age-related changes in the vascular architecture of the rat cerebral cortex. *Neuropathology and applied neurobiology* 1981;**7**:451-62.
- <sup>19</sup> White FH, Al-Azzawi B. Vascularity in experimental oral neoplasia: a stereological approach. *Acta Stereol* 1983;**2**:119-26.

Requests for reprints to: Mr RAJ Spence, Senior Surgical Registrar, The Queen's University of Belfast, Department of Surgery, Institute of Clinical Science, Grosvenor Road, Belfast, BT12 6BJ, Northern Ireland.

## Letters to the Editor

### Necrotising granulomata in prostatic resection specimens—a sequel to previous operation

We read with interest the article by Dr Lee and Dr Shepherd.<sup>1</sup> We have observed similar appearances in two patients, aged 61 and 62 years, who had second transurethral prostatectomies 22 and 25 days respectively after their first operations. There was no clinical evidence of preceding urinary infection or tuberculosis.

In both cases there were granulomata in the second specimen, whereas they were absent from the first, which showed only benign nodular hyperplasia. The giant cells, most of which were of the foreign body type and occasionally contained pigmented particles or fibrinoid material, were situated close to necrotic areas, and there was also an infiltrate of lymphocytes, plasma cells, neutrophils, and eosinophils; the last were numerous in one patient. There was frequent squamous metaplasia of prostatic ducts. We did not find distinctive linear and non-linear changes, as did Dr Lee and Dr Shepherd.

We agree that these lesions are induced by operative trauma. The squamous metaplasia may be the result of ischaemia, but the latter is probably not directly related to the granulomatous reaction since granuloma is not a feature of spontaneous prostatic infarction.

LINDA SINGH  
FE DISCHE

Department of Morbid Anatomy,  
King's College School of Medicine and  
Dentistry,  
Denmark Hill, London SE5 8RX

#### Reference

- <sup>1</sup> Lee G, Shepherd N. Necrotising granulomata in prostatic resection specimens—a sequel to previous operation. *J Clin Pathol* 1983;**36**:1067-70.

### Anti-DNA immunoassay

Concern recently expressed<sup>1</sup> about the quality of anti-DNA test kits is well founded. Single stranded reagents were present in all the samples of labelled native DNA from different commercial sources examined by Medoff.<sup>2</sup> Such defects increase binding values and decrease the specificity of the Farr test.

Since its introduction we have used routinely native <sup>14</sup>C DNA from Amersham in routine immunoassays, though both the stability and the molecular weight distribution of this material are unspecified. Of the many batches purchased, one, specially treated to remove single stranded regions, gave generally lower serum binding values, with some close to zero. As most batches appear to be incompletely double stranded we have reluctantly adopted the Farr test for antibodies to ss-DNA, using heat treated material, despite its disadvantages. Though tests for antibody to native DNA may be valuable for prognosis and treatment, their diagnostic value is uncertain. Using phage derived native DNA Swaak *et al*<sup>3</sup> found increased values were given by sera from patients who did not satisfy the diagnostic criteria for systemic lupus erythematosus.

The Crithidia luciliae immunofluorescence test is deservedly popular, but the problems of variation between observers and in the composition of the labelled antigen-serum remain. It may also detect anti-protein antibodies in some circumstances.<sup>4</sup>

The stimulation of antibody production to native DNA may no longer be considered unique to systemic lupus