

We thank Mr J Reid, Mr J McDermott, and Mr W Kirkwood for the histological staining and S Campbell for typing the paper. RAJ Spence was in receipt of an Eastern Health and Social Services Board research fellowship during this study.

#### References

- <sup>1</sup> Tompsett DH. *Anatomical techniques*. 2nd ed. London: Churchill Livingstone, 1970:9-27.
- <sup>2</sup> Butler H. The veins of the oesophagus. *Thorax* 1951;6:276-96.
- <sup>3</sup> Hales MR, Carrington CB. A pigmented gelatin mass for vascular injection. *Yale J Biol Med* 1971;43:257-70.
- <sup>4</sup> Reynolds DG, Swan KG. Intestinal microvasculature architecture in endotoxic shock. *Gastroenterology* 1972;63:601-20.
- <sup>5</sup> Drury RAB, Wallington EA. *Carleton's histological technique*. 4th ed. Oxford: Oxford University Press, 1967:167.
- <sup>6</sup> Murakami T. Application of the scanning electron microscope to the study of the fine distribution of the blood vessels. *Arch Histol Jap* 1971;32:445-54.
- <sup>7</sup> Northover JMA, Williams EDF, Terblanche J. The investigation of small vessel anatomy by scanning electron microscopy of resin casts. *J Anat* 1980;130:43-54.
- <sup>8</sup> Dollinger RK, Armstrong PB. Scanning electron microscopy of injection replicas of the chick embryo circulatory system. *J Microscopy* 1974;102:179-86.
- <sup>9</sup> Page RB, Munger BL, Bergland RM. Scanning microscopy of pituitary vascular casts. *Am J Anat* 1976;146:273-302.
- <sup>10</sup> Manenti F, Williams R. Injection studies of the splenic vasculature in portal hypertension. *Gut* 1966;7:175-80.
- <sup>11</sup> Greenway CV, Murthy VS. Effects of vasopressin and isoprenaline infusions on the distribution of blood flow in the intestine: criteria for the validity of microsphere studies. *Br J Pharmacol* 1972;46:177-88.
- <sup>12</sup> Micflikier AB, Bond JH, Sircar B, Levitt MD. Intestinal villus blood flow measured with carbon monoxide and microspheres. *Am J Physiol* 1976;230:916-9.
- <sup>13</sup> Grandison AS, Yates J, Shields R. Capillary blood flow in the canine colon and other organs at normal and raised portal pressure. *Gut* 1981;22:223-7.
- <sup>14</sup> Anderson JM. Histometry. In: Bancroft JD, Stevens A, eds. *Theory and practice of histological techniques*. 2nd ed. Edinburgh: Churchill Livingstone, 1982:548-63.
- <sup>15</sup> Aberne WA, Dunnhill MS. *Morphometry*. 1st ed. London: Edward Arnold, 1982:46-118.
- <sup>16</sup> Bradbury S. Commercial image analysers and the characterisation of microscopical images. *J Microscopy* 1983;131:203-10.
- <sup>17</sup> Laursen H. Cerebral vessels and glial cells in liver disease: a morphometric and electron microscope investigation. *Acta Neurol Scand* 1982;65:381-412.
- <sup>18</sup> Wilkinson JH, Hopewell JW, Reinhold HS. A quantitative study of age-related changes in the vascular architecture of the rat cerebral cortex. *Neuropathology and applied neurobiology* 1981;7:451-62.
- <sup>19</sup> White FH, Al-Azzawi B. Vascularity in experimental oral neoplasia: a stereological approach. *Acta Stereol* 1983;2:119-26.

Requests for reprints to: Mr RAJ Spence, Senior Surgical Registrar, The Queen's University of Belfast, Department of Surgery, Institute of Clinical Science, Grosvenor Road, Belfast, BT12 6BJ, Northern Ireland.

## Letters to the Editor

### Necrotising granulomata in prostatic resection specimens—a sequel to previous operation

We read with interest the article by Dr Lee and Dr Shepherd.<sup>1</sup> We have observed similar appearances in two patients, aged 61 and 62 years, who had second transurethral prostatectomies 22 and 25 days respectively after their first operations. There was no clinical evidence of preceding urinary infection or tuberculosis.

In both cases there were granulomata in the second specimen, whereas they were absent from the first, which showed only benign nodular hyperplasia. The giant cells, most of which were of the foreign body type and occasionally contained pigmented particles or fibrinoid material, were situated close to necrotic areas, and there was also an infiltrate of lymphocytes, plasma cells, neutrophils, and eosinophils; the last were numerous in one patient. There was frequent squamous metaplasia of prostatic ducts. We did not find distinctive linear and non-linear changes, as did Dr Lee and Dr Shepherd.

We agree that these lesions are induced by operative trauma. The squamous metaplasia may be the result of ischaemia, but the latter is probably not directly related to the granulomatous reaction since granuloma is not a feature of spontaneous prostatic infarction.

LINDA SINGH  
FE DISCHE

Department of Morbid Anatomy,  
King's College School of Medicine and  
Dentistry,  
Denmark Hill, London SE5 8RX

#### Reference

- <sup>1</sup> Lee G, Shepherd N. Necrotising granulomata in prostatic resection specimens—a sequel to previous operation. *J Clin Pathol* 1983;36:1067-70.

### Anti-DNA immunoassay

Concern recently expressed<sup>1</sup> about the quality of anti-DNA test kits is well founded. Single stranded reagents were present in all the samples of labelled native DNA from different commercial sources examined by Medoff.<sup>2</sup> Such defects increase binding values and decrease the specificity of the Farr test.

Since its introduction we have used routinely native <sup>14</sup>C DNA from Amersham in routine immunoassays, though both the stability and the molecular weight distribution of this material are unspecified. Of the many batches purchased, one, specially treated to remove single stranded regions, gave generally lower serum binding values, with some close to zero. As most batches appear to be incompletely double stranded we have reluctantly adopted the Farr test for antibodies to ss-DNA, using heat treated material, despite its disadvantages. Though tests for antibody to native DNA may be valuable for prognosis and treatment, their diagnostic value is uncertain. Using phage derived native DNA Swaak *et al*<sup>3</sup> found increased values were given by sera from patients who did not satisfy the diagnostic criteria for systemic lupus erythematosus.

The Crithidia luciliae immunofluorescence test is deservedly popular, but the problems of variation between observers and in the composition of the labelled antigen-serum remain. It may also detect anti-protein antibodies in some circumstances.<sup>4</sup>

The stimulation of antibody production to native DNA may no longer be considered unique to systemic lupus

copy right © 1984 by J Clin Pathol; first published as 10.1136/jcp.37.3.354-b on 1 March 1984. Downloaded from http://jcp.bmj.com/ on October 2, 2012 by guest. Protected by copyright.

## Letters to the Editor

erythematosus among the rheumatic diseases. Progress in the synthesis of DNA analogues and the identification of DNA determinants with monoclonal antibodies<sup>7</sup> may, however, lead to a better understanding of the factors determining these immune responses and to improvements in the diagnosis and treatment of diseases in which they are detected.

DJ LEA  
RN HOBBS  
DJ WARD

Department of Rheumatology,  
Robert Jones and Agnes Hunt  
Orthopaedic Hospital,  
Oswestry, Shropshire  
SY10 7AG

## References

- Watson KC, Kerr EJC. Antinuclear antibody-negative systemic lupus erythematosus—how common? *J Clin Pathol* 1983;**36**:1084.
- Medoff ME. Effect of incomplete double strandedness of test DNA on DNA binding results. *J Lab Clin Med* 1982;**99**:410-7.
- Barnett EV. Anti-DNA antibodies: perspectives and perplexities. *J Immunol Methods* 1979;**27**:1-7.
- Swaak AJG, Groenwald J, Aarden LA, Feltkamp TEW. Detection of anti-ds DNA as diagnostic tool. *Ann Rheum Dis* 1981;**40**:45-9.
- Jonsson J, Norberg R. Effect of enzyme treatment of Crithidia luciliae on the reactivity of its kinetoplast with anti-DNA sera. *Scand J Rheumatol* 1976;**5**:221-6.
- Saucier JM, Benard J, DaSilva J, Riou G. Occurrence of a kinetoplast DNA-protein complex in *Trypanosoma cruzi*. *Biochem Biophys Res Comm* 1981;**101**:988-94.
- Rauch J, Lafer E, Andrejewski C, Stollar BD, Schwartz RS. Monoclonal lupus auto-antibodies. *Arthr Rheum* 1982;**25**:744-6.

## Tuberculosis and "sterile" pyuria

In the past "sterile" pyuria suggested the existence of tuberculosis of the urinary tract. In common with all forms of the dis-

ease the incidence of genitourinary tuberculosis has declined but the Medical Research Council survey of the notification of tuberculosis reported 154 such cases in the six months spanning the end of 1978 and the beginning of 1979.<sup>1</sup> These contributed 14% of all cases of non-respiratory disease.

"Sterile" pyuria is still common. To investigate the local situation every specimen received from general practitioners in the Cardiff area over a three year period which showed "sterile" pyuria was cultured for *Mycobacterium tuberculosis*. Between July 1979 and September 1982 a total of 803 samples were tested, of which three (0.4%) yielded *Mycobacterium tuberculosis*.

Urinary tract tuberculosis was therefore clearly not a major cause of "sterile" pyuria. A possible alternative explanation was that many of these patients had recently taken, or were still taking, antimicrobial agents at the time of sampling. A study of the request forms showed that antimicrobial treatment was mentioned in 161 (20.0%). One of the positive isolations came from this group.

Information on request forms is notoriously unreliable. All urines showing "sterile" pyuria were therefore tested for antimicrobial activity using a modification of the method described by Pelling,<sup>2</sup> and the actual findings were compared with the expected findings. The results are shown in the Table. They were discrepant in 47 (19.7%) of the 239 cases.

The local incidence of urinary tract tuberculosis in patients whose urine showed "sterile" pyuria was low, and so the cost effectiveness of surveying all such samples must be equally low. It is not possible to reduce the number of specimens requiring screening by excluding all those reported as being on antimicrobial therapy because one of our three cases came from this group and the accuracy of the reported data on the request forms is low.

Of the 11 bacteriologically proved cases of urinary tract tuberculosis which occurred in South Glamorgan between July 1979 and September 1982, three (27.3%) were detected as a result of this survey.

Such patients are unlikely to pose a major infection hazard to other members of the community, but all three cases detected by the survey were unsuspected clinically and one suffered complications which may have been related to his infection.

It is the practice of this laboratory therefore not to examine every specimen showing "sterile" pyuria for *Mycobacterium tuberculosis* but to advise local practitioners to consider the diagnosis of urinary tract tuberculosis when persistent unexplained "sterile" pyuria is observed.

CD RIBEIRO  
PA JENKINS

Public Health Laboratory,  
University Hospital of Wales,  
Heath Park, Cardiff, CF4 4XW

## References

- Medical Research Council National Survey of tuberculosis notification in England and Wales 1978/79. *Br Med J* 1980;**281**:895-8.
- Pelling W. Inhibitory substances in urine: an addition to the routine screen. *Med Lab Sci* 1982;**39**:377-81.

## Batch screening method for detection of bacteriuria

While agreeing with Mr Kerfoot and his colleagues<sup>1</sup> that the increasing workload in urine microbiology places a considerable strain on laboratories, I believe that it is important that protocols for examining urine specimens should not give wrong, or unduly delayed, answers. I therefore have some reservations about the batch screening method which they describe. Firstly, and perhaps most important, I do not consider that it is possible for patients, especially women, to pass a mid-stream urine specimen into a small container of the size they use. If their procedure—and they do not say so—is that the urine is first passed into a container with a wide mouth, and then decanted, the cost of specimen collection must be greatly increased.

Secondly, since the boric acid preservation method was first described<sup>2</sup> further work<sup>3</sup> has shown that the percentage of false negatives may be as high as 16%. Many would not consider it a satisfactory method of urine preservation, especially when definitive culture, as described by Kerfoot *et al*, may not be undertaken for up to 48 h after the specimen is collected.

This screening method inevitably delays definitive culture and sensitivity testing by at least 24 h. Kerfoot *et al* do not say

## Comparison of expected and actual results of tests for antimicrobial activity in urine

	Antimicrobials mentioned on request form	
	Yes	No
Antimicrobials found in urine		
Yes	19	33
No	14	173