Letters to the Editor

- 9 Trowbridge EA, Slater DN, Kishk YT, *et al*. Platelet production in myocardial infarction and sudden cardiac death. *Thromb Haemostas* 1984;52:167-71.
- 10 Martin JF, Slater DN, Kishk YT, Trowbridge EA. Platelet and megakaryocyte changes in cholesterol induced experimental atherosclerosis. *Arteriosclerosis* 1985;5:604-12.
- 11 Aabo K, Bendix-Hansen K. Megakaryocytes in pulmonary blood vessels. Incidence at necropsy, clinicopathological relations, especially to disseminated intravascular coagulation. *Acta Pathol Microbiol Scand* 1978;86:285-91.

Drs Talbot and Sharma reply:

We agree with Drs Trowbridge, Slater, and Martin that our estimates of the numbers of megakaryocytes trapped in the pulmonary capillary bed would be even higher if we had applied a correction for variations in megakaryocyte size.

Although our own data cannot prove the hypothesis that the lungs are the principal site of thrombopoiesis, there are persuasive arguments for this. Evidence which has not to our knowledge been considered, is provided by the blood of lungless vertebrates such as fish,¹ which lack platelets, their role being performed by large nucleated cells known as thrombocytes, which resemble the other blood cells in cytological appearance.

IC TALBOT
G SHARMA

Department of Pathology,
Clinical Sciences Building,
Leicester Royal Infirmary LE2 5LX

Reference

- Roberts R. *Fish pathology*. London: Baillière-Tiridall, 1978.

Peripheral thrombocytopenia in human parvovirus infection

Human parvovirus (HPV) causes aplastic crises in patients with hereditary¹ or acquired² haemolytic anaemias. HPV induced red cell aplasia in patients without a shortened red cell survival is difficult to establish because of the transient nature of the erythroblastopenia. On the other hand, a moderate thrombocytopenia of central origin seems common during these aplastic crises.^{3,4}

We describe here the first documented case of peripheral thrombocytopenia with HPV associated aplastic crisis in a patient without shortened red cell survival.

Case history

A five year old girl was admitted to hospital because she had had petechial purpura on her face and legs for 24 hours after an episode of upper respiratory tract infection. Oral examination showed the presence of a superficial haemorrhagic bulla. Neither lymphadenopathy or splenomegaly was detected. The patient was febrile. The blood test showed normal haemoglobin concentration and white cell count but a platelet count of $8 \cdot 10^9/l$, although it had been $304 \times 10^9/l$ 15 months previously. Bone marrow examination performed on the day of admission showed an increased number of megakaryocytes, suggesting a peripheral origin of thrombocytopenia and an absence of erythroid precursors. The reticulocyte count was then found to be less than $5 \times 10^9/l$. This erythroblastopenia suggested a recent HPV infection, which was confirmed by the presence of anti-HPV IgM in the serum.⁵ Other recent infections were excluded by tests for infectious mononucleosis, cytomegalovirus, rubella, mumps, hepatitis A and B. Treatment with high dose intravenous gammaglobulin (daily dose of 30 g) was started immediately. Two days later the platelet count remained low (figure) and gammaglobulin was discontinued. Consequently, corticosteroids (prednisone 2 mg/kg/day) were given, and the platelet count returned to normal within three days. Haemoglobin concentration and white cell count remained normal during the evolution. The daily dose of steroids was progressively reduced after one month and the treatment finally discontinued after two months. Further controls showed normal platelet counts.

Discussion

This observation suggests that HPV may be

associated with intense peripheral platelet destruction. Haemoglobin concentration was not modified as a consequence of an aplastic crisis, as anaemia does not occur as long as red cell survival is normal; the transient nature of the erythroblastopenia is responsible for the difficulty in detecting red cell aplasia in normal people infected with HPV. To our knowledge, these findings of an HPV associated aplastic crisis in a patient free of chronic haemolytic anaemia have not been reported previously. None the less, the HPV associated aplastic crisis seems to be totally different from the "transient erythroblastopenia of childhood," whose mechanism remains to be explained.

JJ LEFRERE*

D GOT†

*Centre National de Transfusion Sanguine,
6 Rue Cabanel, 75015,
Paris, France.

†Service de Pédiatrie,
Hôpital d'Evreux, 27023,
France.

References

- 1 Pattison JR, Jones SE, Hodgson J, Davis LR, White JM, Stroud CE, Murtaza L. Parvovirus infection and hypoplastic crisis in sickle cell anaemia. *Lancet* 1981;i:664-5.
- 2 Bertrand Y, Lefrère JJ, Leverger G, Couroucé AM, Féo C, Clark M, Schaison G, Soulier JP. Autoimmune haemolytic anaemia revealed by human parvovirus linked erythroblastopenia. *Lancet* 1985;ii:382-3.
- 3 Lefrère JJ, Couroucé AM, Giro R, Bertrand Y, Soulier JP. Six cases of hereditary spherocytosis revealed by human parvovirus infection. *Br J Haematol* 1986;62:653-8.
- 4 Saunders PWG, Reid MM, Cohen BJ. Human parvovirus induced cytopenias: a report of five cases. *Br J Haematol* 1986;63:407-10.
- 5 Cohen BJ, Mortimer PP, Pereira MS. Diagnostic assays with monoclonal antibodies for the serum parvovirus-like virus (SPLV). *J Hyg (Camb)* 1983;91:113-30.

Figure Haematological data in patient studied.