Incidence of vaginal *Weeksellia virosa* (formerly group IIf)

C MARDY,* B HOLMES†

From the *Department of Pathology, The General Hospital, Altrincham, Cheshire, and the †Computer Identification Laboratory, National Collection of Type Cultures, Central Public Health Laboratory, London

SUMMARY The antimicrobial susceptibilities, biochemical properties, and cultural characteristics of six strains of *Weeksellia virosa* (formerly group IIf) were determined. The main characteristics of this non-fermentative organism were production of cytochrome oxidase, gelatinase, and indole, but a lack of saccharolytic activity.

A study was then made of the isolation rates of *Weeksellia virosa* from high vaginal swabs from 300 women: a healthy control group (n = 100), a general group with symptoms of vaginal infection (n = 100), and 100 women from a remand centre, where likelihood of exposure to sexually transmitted disease might be expected to be higher. The incidence of *Weeksellia virosa* was found to be 2% in the first two groups. This suggests that, irrespective of the presence of pathogens, the incidence of the species in the general female population is around 2%. A much higher incidence (15%) was found in the third group, suggesting sexual transmission of the organism.

In 1970 Pickett and Manclark described eight strains which they regarded as non-saccharolytic flavobacteria.¹ They reported the biochemical characteristics of the strains^{2,3} and their susceptibility to antimicrobial agents,⁴ but they did not consider any of them to be pathogenic. Twelve similar strains were reported by Olsen and Ravn⁵; they were isolated predominantly from female genital tract specimens submitted for examination for *Neisseria gonorrhoeae*. Owen and Snell examined eight such strains, five were from the urethra and one each from the cervix and vagina⁶; they found the properties of the strains to be intermediate between those of *Flavobacterium* and *Moraxella*. Tatum *et al* described 78 strains and designated them as group IIf⁷; they recently reported 87 strains primarily from urine (45%), the cervix (15%), and the vagina (15%).⁸ A single group IIf strain, from the vagina of an asymptomatic postmenopausal woman, has also been reported.⁹ A new genus and species, *Weeksellia virosa*, has now been proposed for the group.¹⁰

In the pathology department at Altrincham General Hospital an unidentified non-fermentative Gram negative rod was occasionally seen on selective chocolate blood agar plates being used for the isolation of *Neisseria gonorrhoeae* from high vaginal swabs. Representative strains of this unidentified organism

submitted to the Computer Identification Laboratory of the National Collection of Type Cultures (NCTC) were identified as *W. virosa*. Six *W. virosa* strains were selected from a number of such strains isolated from high vaginal swabs and characterised at the hospital in a wide range of biochemical tests, in the API 20E system, and in the Roche Oxi/Ferm tube.

A study was made of the rates of isolation of *Weeksellia virosa* from the high vaginal swabs of three different groups of women. Group 1 comprised well women showing no signs of bacterial infection; this group consisted mainly of volunteers and those undergoing routine preoperative high vaginal swabs. Group 2 comprised patients referred for investigation due to symptoms of possible vaginal infection. Group 3 comprised women from a special source (women's remand centre) who were possibly at greater risk of contracting sexually transmitted disease (STD). One hundred specimens were sampled from each group.

Material and methods

Charcoal swabs, transported in Stuart's medium (Medical Wire STMW 219) were used for the collection of high vaginal specimens. Selective chocolate blood agar containing colistin sulphate (7.5 µg/ml), trimethoprim (5 µg/ml), and vancomycin (3 U/ml) was used routinely for the isolation of *Neisseria gonorrhoeae*. The presence of antibiotics in the media ensures

the suppression of normal vaginal flora (*W. virosa* grows well on this medium as it is resistant to all three antibiotics at these concentrations).

Initially a smear was prepared on a sterile slide, and each high vaginal swab was subcultured on to non-selective chocolate (10% w/v) blood agar, selective chocolate (10% w/v) blood agar, and finally into *Trichomonas* medium (Medical Wire TMMW 220). The smear was fixed and stained by Gram's method (Lillie's modification).¹¹ The *Trichomonas* medium was incubated for 48 hours at 37°C aerobically; both chocolate blood agar plates were incubated in an atmosphere of about 8–10% (v/v) carbon dioxide for 48 hours.

CULTURAL CHARACTERISTICS

Commercially poured agar plates were used (Analytical and Bacteriological Media, London). The six strains chosen for detailed study were inoculated on to 10% (w/v) horse blood agar, 10% (w/v) selective chocolate horse blood agar, MacConkey agar and nutrient agar. The blood agar plates were incubated at 4°C, 22°C, 37°C and 43°C. Smears were made from colonies of each strain and stained by Gram's method.¹¹

CONVENTIONAL BIOCHEMICAL TESTS

All tests were recorded at or up to seven days' incubation, except where stated. The following tests were carried out mainly by the media and methods described by Cowan and Steel¹¹: cytochrome oxidase production (one day); acid from carbohydrates (method 1); oxidation or fermentation (or alkali production) in glucose O-F medium; citrate utilisation (method 1); dihydrolase for arginine and decarboxylases for lysine and ornithine (method 1); H₂S production (method 1); indole production (methods 2 and 3); potassium cyanide tolerance (two days); production of acetylmethylcarbinol (Voges-Proskauer test, Baritt's method); β -galactosidase production (two days); and urease production (method 1). The following tests were carried out by the specific methods indicated: catalase production,¹² methyl red reaction,¹³ nitrate reduction,¹⁴ phenylalanine deamination¹⁵ and gelatinase production.¹⁶ Pigment production was examined daily for five days on nutrient agar and on 1% (w/v) tyrosine agar.

API 20E TESTS

Each of the six strains was inoculated into an API 20E strip and incubation for 48 hours was required in accordance with the manufacturer's instructions; the six supplementary tests required were also performed. The nine digit profile number thus obtained was consulted in the *API 20E Analytical Profile Index* (July 1985 edition). Any additional tests required by the API

20E system to further the identification were also performed.

OXI/FERM TUBES

In accordance with the manufacturer's instructions the six strains were inoculated into Oxi/Ferm tubes; after incubation for 48 hours indole reagent was added to the appropriate test compartment. The profile number obtained was consulted in the Oxi/Ferm tube identification scheme (CCIS system 1979–1432) and any supplementary tests required to further the identification were performed.

ANTIMICROBIAL SUSCEPTIBILITY

Tests were performed by inoculating blood agar plates on a rotary plater from four to six hour broth cultures. After antibiotic discs were added the plates were incubated at 37°C for 18 hours. The following antibiotic discs were used: ampicillin (10 μ g); chloramphenicol (25 μ g); colomycin (25 μ g); erythromycin (5 μ g); kanamycin (30 μ g); neomycin (10 μ g); novobiocin (5 μ g); penicillin G (1 U); streptomycin (25 μ g); tetracycline (25 μ g); *Escherichia coli* NCTC 10418 was used as a control organism.

ISOLATION STUDY

The same procedure as for the routine examination of high vaginal swabs was followed except that high vaginal swabs from group 1 were also subjected to this procedure; each high vaginal swab was also subcultured on to a 10% (w/v) blood agar plate which was incubated for 48 hours at 37°C aerobically. Colonies suspected of being *W. virosa* were examined for cytochrome oxidase production and by Gram stain. Colonies giving the reactions expected for *W. virosa* had their identities confirmed by the API 20E system.

Results

CULTURAL CHARACTERISTICS

After 18 hours of incubation on 10% (w/v) blood agar all six strains appeared as cream-coloured, moist, domed colonies (1–2 mm in diameter). On further incubation the colonies became slightly larger and very glossy. A pale α -haemolysis was noted after 36 hours of incubation which intensified when the plates were left at room temperature. Growth on non-selective and selective chocolate blood agar was much better than on ordinary blood agar—very moist, cream-coloured colonies 2–3 mm in diameter were obtained after 18 hours. On further incubation the colonies became confluent, making subsequent subculture on to further chocolate blood agar plates to obtain single colonies very difficult, as confluent growth was obtained however small the inoculum. The

Table 1 Biochemical characteristics of *Weeksella virosa**Tests in which all six strains gave positive results*

Alkaline reaction in O-F test
 Catalase production
 Cytochrome oxidase production
 Gelatinase production
 Indole production (Ehrlich's reagent)
 Production of brown pigment after 24 hours on 1% (w/v) tyrosine agar

Tests in which all six strains gave negative results:

Acid from the following
 carbohydrates in peptone water medium:
 Glucose
 Adonitol
 Arabinose
 Dulcitol
 Glycerol
 Inositol
 Lactose
 Maltose
 Mannitol
 Sorbitol
 Sucrose
 Trehalose
 Arginine dihydrolase

Citrate utilisation
 Fermentative reaction in O-F test
 Hydrogen sulphide production
 Indole production (Kovac's reagent)
 Lysine decarboxylase
 Methyl red test
 Nitrate reduction
 Ornithine decarboxylase
 Oxidative reaction in O-F test
 Phenylalanine deamination
 Pigment production on nutrient agar
 Potassium cyanide tolerance
 Urease production
 Voges-Proskauer reaction
 β -galactosidase production

Table 2 Isolation rates in different groups of women

Group No (100 patients in each group)	No of isolations of <i>Weeksella virosa</i>	No of specimens in which a pathogen was also isolated
1	2	0
2	2	2
3	15	5

six strains produced luxurious growth on nutrient agar and the growth was seen to flow down nutrient agar slopes, collecting in the bottom of the bilou jar. No growth was obtained on MacConkey agar. Growth was obtained on 10% (w/v) blood agar and in nutrient broth at 22°C, 37°C, and 43°C. No growth was seen in either medium at 4°C. In a hanging drop suspension prepared from the nutrient broths at 22°C and 37°C, the strains showed no motility. In Gram stained films from blood agar the organisms appeared as large (about 3–5 μ m in length), Gram negative rods with rounded ends; some short coccobacillary forms were also seen. In the broth cultures the cells tended to form long filaments.

The results of the biochemical tests are given in table 1.

API 20E TESTS

After 48 hours of incubation in the API 20E system all six strains yielded the profile number 0002004-00. This number is listed in the *API 20E Analytical Profile Index* with the comment "low discrimination", but the most likely taxon is CDC group IIf, followed by *Moraxella* spp. The results of four additional tests

(acetate utilisation, growth at 42°C, growth on MacConkey agar and pigment production) allowed the strains to be ascribed to CDC group IIf using the appropriate table in the *API 20E Profile Index*.

OXI/FERM TUBES

An "ID value" of 0201 was obtained from the Roche Oxi/Ferm tube results for all six strains after 48 hours of incubation. The "ID value" in the CCIS system listed three possible taxa, but performance of the five supplementary tests (alkaline reaction in xylose, penicillin susceptibility, production of brown pigment, production of yellow pigment and "runny growth") permitted identification of the strains as "group 2F, *Flavobacterium*-like."

ANTIMICROBIAL SUSCEPTIBILITY

All six strains were resistant to kanamycin, neomycin, and trimethoprim; the strains were susceptible to ampicillin, chloramphenicol, colomycin, erythromycin, novobiocin, penicillin G, streptomycin, and tetracycline.

ISOLATION STUDY

The isolation rates of *Weeksella virosa* are given in table 2. A higher incidence of *Weeksella virosa* was seen in the group of women from the remand centre than in the two other groups. The presence of common pathogens and the results of the Gram stained high vaginal smears were noted for any patient from whom *Weeksella virosa* was isolated. The pathogens included *Candida albicans*, *Neisseria gonorrhoeae*, and *Trichomonas vaginalis*. No clinically important information could be formulated from the Gram stained smears, most of which contained large numbers of normal flora: Gram positive bacilli, Gram negative rods, Gram positive cocci, and in some cases, a presumptive pathogen such as yeast cells. In most smears of patients from whom *Neisseria gonorrhoeae* had been isolated, the normal flora was reduced.

Discussion

Few reports on the incidence of *Weeksella virosa* in human clinical specimens have been published. Tatum *et al* identified only 78 cultures as group IIf from among 35 000 cultures received for identification between 1949 and 1973.⁷ The incidence of this species in clinical material seems to be low, but the results of the present study suggest that the incidence of the species is higher and that the organism is probably ignored in most routine hospital laboratories, especially when isolated from clinical specimens containing a high proportion of normal flora. The mucoid nature of the colonies would often cause them to be mistakenly regarded as klebsiellas.

The clinical importance of the presence of *W virosa* is difficult to assess from the results of the isolation rate study. In group 2 *W virosa* was isolated only when a pathogen was present. This might suggest that the organism is normally suppressed in a healthy patient but becomes prominent when a patient is immuno-compromised. The same isolation rate for *W virosa*, however, was obtained in control group 1, suggesting that irrespective of the presence of pathogens the incidence of *W virosa* in the general female population is around 2%. Group 3 probably included some women with a lot of sexual partners and therefore passage of *W virosa* might be increased. This could account for the higher isolation rate in this group (15%) compared with groups 1 and 2. Interestingly, in group 3 although the incidence of pathogens present in the same patient from whom *W virosa* was isolated, was higher (5%) than that of group 2 (2%), the increase was not as high as that seen for *W virosa* (15%).

W virosa is apparently much more common in women than in men, with the highest isolation rates from urine and the genital or urethral tract.^{8 10} In this study all isolations of the organism were from the female genital tract. It would be interesting to know if a link between *W virosa* and sexual transmission could be proved, but this would require a much more extensive study (including groups of male patients), perhaps on a national scale.

W virosa is isolated with apparent ease from vaginal specimens and may be identified with a minimum amount of labour and equipment, using either the API 20E system or the Roche Oxi/Ferm tube. The conventional biochemical test results given in table 1 agree with those given in the original description of the species.¹⁰ Although almost all non-fermenters also fail to exhibit acid production in a peptone water medium, *W virosa* strains are also non-saccharolytic in ammonium salt sugar medium, which is generally more appropriate for showing acid production from carbohydrates by non-fermenters.¹⁰ Indole production by *W virosa* strains is characteristic, but the results clearly show that a sensitive technique is necessary.

Flavobacterium species are highly resistant organisms but *W virosa* has been reported to be much more susceptible, even to penicillin, with resistance shown only to kanamycin and tobramycin.¹⁷ *W virosa* has been shown to be susceptible to all β -lactam producing antibiotics tested, as well as to aztreonam, chloramphenicol, and clindamycin.^{4 18 19} The results of the present study agree with these findings.

We thank Dr R Jennings and Miss A J Kelly for their invaluable support.

References

- Pickett MJ, Manclark CR. Nonfermentative bacilli associated with man: I. Nomenclature. *Am J Clin Pathol* 1970;**54**:155-63.
- Pickett MJ, Pedersen MM. Nonfermentative bacilli associated with man: II. Detection and identification. *Am J Clin Pathol* 1970a;**54**:164-77.
- Pickett MJ, Pedersen MM. Salient features of non-saccharolytic and weakly saccharolytic nonfermentative rods. *Can J Microbiol* 1970b;**16**:401-9.
- Pedersen MM, Marso E, Pickett MJ. Nonfermentative bacilli associated with man: III. Pathogenicity and antibiotic susceptibility. *Am J Clin Pathol* 1970;**54**:178-92.
- Olsen H, Ravn T. *Flavobacterium meningosepticum* isolated from the genitals. *Acta Pathol Microbiol Scand (Sect B)* 1971;**79**:102-6.
- Owen RJ, Snell JJS. Comparison of group IIb with *Flavobacterium* and *Moraxella*. *Antonie van Leeuwenhoek* 1973;**39**:473-80.
- Tatum HW, Ewing WH, Weaver RE. Miscellaneous Gram-negative bacteria. In: Lennette EH, Spaulding EH, Truant JP, eds. *Manual of clinical microbiology*. 2nd ed. Washington, DC: American Society for Microbiology, 1974:270-94.
- Clark WA, Hollis DG, Weaver RE, Riley P. *Identification of unusual pathogenic Gram-negative aerobic and facultatively anaerobic bacteria*. Atlanta: US Department of Health and Human Services, 1984.
- Tashjian JH, Coulam CB, Washington II JA. Vaginal flora in asymptomatic women. *Mayo Clin Proc* 1976;**51**:557-61.
- Holmes B, Steigewalt AG, Weaver RE, Brenner DJ. *Weeksella virosa* gen. nov, sp. nov. (formerly group IIb), found in human clinical specimens. *System Appl Microbiol* 1986;**8**:185-90.
- Cowan ST, Steel KJ. *Manual for the identification of medical bacteria*. Cambridge: Cambridge University Press, 1965.
- Deibel RH, Evans JB. Modified benzidine test for the detection of cytochrome-containing respiratory systems in microorganisms. *J Bacteriol* 1960;**79**:356-60.
- Clark WM, Lubs HA. The differentiation of bacteria of the colon-aerogenes family by the use of indicators. *J Infect Dis* 1915;**17**:160-73.
- Cook GT. A plate test for nitrate reduction. *J Clin Pathol* 1950;**3**:359-62.
- Ewing WH, Davis BR, Reavis RW. Phenylalanine and malonate media and their use in enteric bacteriology. *Public Health Laboratory* 1957;**15**:153-67.
- Kohn J. A preliminary report of a new gelatin liquefaction method. *J Clin Pathol* 1953;**6**:249.
- Von Graevenitz A. Clinical significance and antimicrobial susceptibility of flavobacteria. In: Reichenbach H, Weeks OB, eds. *The Flavobacterium-Cytophaga group*. Proceedings of the international symposium on yellow-pigmented Gram-negative bacteria of the *Flavobacterium Cytophaga* group, Braunschweig July 8th to 11th, 1980. Weinheim: Verlag Chemie, 1981:153-64.
- Fass RJ, Barnishan J. In vitro susceptibilities of non-fermentative Gram-negative bacilli other than *Pseudomonas aeruginosa* to 32 antimicrobial agents. *Rev Infect Dis* 1980;**2**:841-53.
- Schell RF, Francisco M, Bihl JA, LeFrock JL. The activity of ceftazidime compared with those of aztreonam, newer cephalosporins and Sch 29482 against nonfermentative Gram-negative bacilli. *Chemotherapy* 1985;**31**:181-90.

Requests for reprints to: Dr B Holmes, NCTC, Central Public Health Laboratory, 61 Colindale Avenue, London NW9 5HT, England.