

Microbiology of infected pilonidal sinuses

I BROOK

From the Naval Hospital, Bethesda, Maryland, and the Walter Reed Army Hospital, Washington, DC, United States of America

SUMMARY Aspirates of pus from infected pilonidal sinuses in 75 patients showed bacterial growth. Anaerobic bacteria only were recovered in 58 (77%) specimens, aerobic bacteria only in three (4%), and mixed aerobic and anaerobic bacteria in 14 (19%). Two hundred and nine isolates were recovered: 147 anaerobes (2.0 isolates a specimen) and 62 aerobes (0.8 a specimen). The predominant anaerobes were *Bacteroides* sp (81 isolates, including 29 *Bacteroides fragilis* group) and 51 anaerobic cocci. The predominant aerobes were *Escherichia coli* (n=15), *Proteus* sp (n=9), group D streptococcus (n=7), and *Pseudomonas* sp (n=7).

This study highlights the polymicrobial nature and predominance of anaerobic bacteria in infected pilonidal sinuses.

Infection in pilonidal sinuses may be protracted, cause local irritation, and in rare cases lead to serious complications such as bacteraemia or meningitis. Most reports have described small groups of patients.¹ This report describes the microbiology of pilonidal sinuses in a group of 75 patients, which is the largest study to date, and therefore permits more precise assessment of the different organisms contributing to the infection.

Patients and methods

Between June 1975 and June 1985, 82 aspirates taken from pilonidal sinuses were processed for aerobic and anaerobic bacteria by the clinical microbiology laboratories at Walter Reed Army Medical Center in Washington, DC, and the Naval Medical Center in Bethesda, Maryland. Bacterial growth was observed in 75 (91%) cases. Mean age of the 75 patients was 27 years (range 3 to 47 years), and 61 were men. Twenty three of the patients received antibiotics before their samples were collected. Cultures were obtained either by direct needle aspiration of the purulent contents into a syringe that was immediately sealed and transported to the laboratory within 30 minutes, or by a swab that was dipped into the pus and introduced into an anaerobic transport system (Port-A-Cul, BBL, Cockeysville, Maryland) and sent to the laboratory within two hours. The specimen material was plated

on to pre-reduced vitamin K₁-enriched brucella blood agar, an anaerobic blood agar plate containing kanamycin and vancomycin, an anaerobic blood plate containing colistin and nalidixic acid, and an enriched thioglycolate broth (containing haemin, sodium bicarbonate, and vitamin K₁).² The material was then incubated in GasPak jars (BBL, Cockeysville, Maryland) and examined at 48 and 96 hours. Plates that showed growth were held until the organisms were processed and identified. All cultures that showed no growth were held for at least five days. Anaerobes were identified using the API anaerobic system (Analytab Product, Inc, Plainview, New York) at the Walter Reed Hospital and the Minitek system (Baltimore Biological Products, Cockeysville, Maryland) at the Naval Hospital. Other carbohydrate tests (Scott Laboratories, Fiskeville, Rhode Island) and gas liquid chromatography³ were performed as needed to identify the organisms. Sheep blood (5%), chocolate agar, and MacConkey agar plates were inoculated for the isolation of aerobic organisms. The plates were incubated at 37°C aerobically (MacConkey) and under 5% carbon dioxide (blood and chocolate), and examined at 24 and 48 hours. Aerobic bacteria were identified using conventional methods.³

Results

Anaerobic bacteria only were recovered in 58 (77%) specimens, aerobic bacteria only in three (4%), and mixed aerobic and anaerobic bacteria in 14 (19%). A total of 209 bacterial isolates were recovered, accounting for 2.8 isolates per specimen (2.0 anaerobes and 0.8 aerobes per specimen) (table).

Accepted for publication 12 June 1989

The opinions and assertions contained herein are those of the writers and are not to be construed as official or reflecting the views of the Department of Defense, the Navy Department, or the Naval Service at large.

Table Bacterial isolates for 75 pilonidal sinuses

Aerobic and facultative bacteria		Anaerobic bacteria	
<i>Gram positive cocci:</i>		<i>Anaerobic cocci:</i>	
Alpha-haem streptococci	6	<i>Peptostreptococcus</i> sp	15
Group D streptococci	7	<i>P asaccharolyticus</i>	5
Non-haem streptococci	3	<i>P prevotii</i>	7
<i>Streptococcus pyogenes</i>	4	<i>P anaerobius</i>	20
<i>Staphylococcus aureus</i>	3	<i>P magnus</i>	2
<i>Staphylococcus epidermidis</i>	2	<i>S constelatus</i>	1
		Microaerophilic streptococcus	1
		<i>Veillonella alcalescens</i>	2
		<i>Veillonella parvula</i>	1
<i>Gram negative bacilli:</i>		<i>Gram positive bacilli:</i>	
<i>Proteus morgani</i>	4	<i>Eubacterium lentum</i>	1
<i>Proteus mirabilis</i>	2	<i>Prop ionibacterium</i> sp	1
<i>Proteus vulgaris</i>	2	<i>P acnes</i>	2
<i>Proteus</i> sp	1	<i>Lactobacillus</i> sp	3
<i>Escherichia coli</i>	15	<i>Clostridium</i> sp	2
<i>Klebsiella pneumoniae</i>	3		
<i>Klebsiella oxytoca</i>	1	<i>Gram negative bacilli:</i>	
<i>Pseudomonas aeruginosa</i>	5	<i>F nucleatum</i>	2
<i>Pseudomonas</i> sp	2	<i>F mortiferum</i>	1
<i>Enterobacter</i> sp	2	<i>Bacteroides</i> sp	19
		<i>B fragilis</i>	15
Total	62	<i>B distasonis</i>	2
		<i>B ovatus</i>	4
		<i>B vulgatus</i>	2
		<i>B thetaiotaomicron</i>	6
		<i>B melaninogenicus</i>	13
		<i>B intermedium</i>	3
		<i>B asaccharolyticus</i>	4
		<i>B ureolyticus</i>	9
		<i>B bivius</i>	2
		<i>B oris-buccae</i>	2
		Total	147

A total of 147 anaerobic bacteria were recovered. The predominant anaerobes recovered were *Bacteroides* sp (81 isolates, including 29 *Bacteroides fragilis* group) and 51 anaerobic cocci. A total of 62 aerobic isolates were recovered. The predominant ones were *Escherichia coli* (n = 15), *Proteus* sp (n = 9), group D streptococcus (n = 7), and *Pseudomonas* sp (n = 7). There was no consistent pattern of combinations, although *Bacteroides* sp were recovered in 38 instances with anaerobic cocci.

Discussion

This study confirms previous reports in adults^{1,4,5} and children⁶ by showing the polymicrobial nature and predominance of anaerobic bacteria in infected pilonidal sinuses, where anaerobes outnumbered aerobes in a ratio of 2.4:1. In contrast to other investigators,^{1,4,5} Gram negative aerobic and facultative bacilli, especially *E coli*, *Proteus* sp, and *Pseudomonas* sp, were isolated in many instances, and like other reports, *Staphylococcus* sp were only occasionally recovered. The recovery of Gram negative bacilli is not surprising as these organisms are part of the normal gastrointestinal flora.

We have already described the microbiology of 25 infected pilonidal sinuses in children.⁶ A total of 63 anaerobes (2.5 per patient) and 13 aerobes (0.5 per patient) were recovered. The predominant anaerobic isolates were *Bacteroides* sp, anaerobic cocci, *Clostridium*, and *Fusobacterium*. The predominant aerobes were *E coli* and group D streptococcus.

Finegold reviewed 13 publications dealing with the bacteriological characteristics of infected pilonidal sinuses in adults.¹ The most common isolates reported in those studies were various *Bacteroides* sp, including *B fragilis*, anaerobic Gram positive cocci, and *Clostridium* sp. The most recent report presenting the bacteriological features of pilonidal sinuses in adults reported 11 cases.⁴ Anaerobic organisms such as *Bacteroides* sp and Gram positive anaerobic cocci were the predominant isolates. Gram negative enteric bacteria and *S aureus* were not present.

Infections with *Bacteroides* sp are generally polymicrobial where these organisms are recovered mixed with other anaerobic, facultative anaerobic, and aerobic bacteria.¹ Previous studies reported that the association between *Bacteroides* sp and its counterparts is generally synergistic.⁷⁻⁹ In studies using an intra-abdominal abscess model, Weinstein *et al* proved the need to use antibiotics that are effective against both *Enterobacteriaceae* and *Bacteroides fragilis*.¹⁰ The synergy among the different bacterial strains in mixed polymicrobial aerobic and anaerobic infections may be due to protection from phagocytosis and intracellular killing,¹¹ production of essential growth factors,¹² and lowering of oxidation reduction potential in infected host tissues.¹³

The isolation of anaerobic bacteria mixed with aerobic and facultative organisms at that site is not surprising as anaerobes are the predominant organisms in the gastrointestinal tract, where they outnumber aerobes by 1000:1.¹⁴ Because anaerobic bacteria are often associated with pilonidal sinuses, physicians should consider their presence if antimicrobial treatment is used. Gram staining of aspirated pus and appropriate aerobic and anaerobic microbiological techniques can help in the selection of proper treatment. As some of the anaerobes are resistant to penicillin, treatment should also include appropriate coverage of those organisms. Surgical drainage is still the treatment of choice. The presence of penicillin resistant anaerobic bacteria, however, such as *B fragilis*¹⁵ and some strains of the *B melaninogenicus* group,^{15,16} may warrant the administration of appropriate antimicrobial agents such as clindamycin, ceftioxin, metronidazole, imipenem, or the combination of a β lactamase inhibitor and a penicillin.

I acknowledge the efforts of the staff of the Clinical Microbiology Laboratories at Walter Reed Army

Medical Center and the Naval Medical Center, the editorial assistance of Junith Van Deusen, and the secretarial assistance of Marianne Owens.

References

- 1 Finegold SM. *Anaerobic bacteria in human disease*. New York: Academic Press Inc, 1977:389.
- 2 Sutter VL, Citron DM, Edelstein MAC, Finegold SM: *Wadsworth anaerobic bacteriology manual*. 4th ed. Belmont, California: Star Publishing Company, 1985.
- 3 Lennette EH, Balows A, Hausler Jr W, Shadomy HJ. *Manual of clinical microbiology*. 4th ed. Washington, DC: American Society for Microbiology, 1985.
- 4 Marrie TJ, Aylward D, Kerr E, *et al*. Bacteriology of pilonidal cyst abscesses. *J Clin Pathol* 1978;**31**:909.
- 5 Meislin HW, Lerner SA, Graves MH, *et al*. Cutaneous abscesses: anaerobic and aerobic bacteriology and outpatient management. *Ann Intern Med* 1977;**87**:145-9.
- 6 Brook I. Aerobic and anaerobic microbiology of pilonidal cyst abscesses in children. *Am J Dis Child* 1980;**134**:679-80.
- 7 Meloney F, Olpp J, Harvey HD, Zaysteff-Jern H. Peritonitis. II. Synergism of bacteria commonly found in peritoneal exudates. *Archives of Surgery* 1932;**25**:709-21.
- 8 Hite KE, Locke M, Hesseltine HC. Synergism in experimental infections with nonsporulating anaerobic bacteria. *J Infect Dis* 1949;**84**:1-9.
- 9 Brook I, Hunter V, Walker RI. Synergistic effect of *Bacteroides*, *Clostridium*, *Fusobacterium*, anaerobic cocci, and aerobic bacteria on mortality and induction of subcutaneous abscesses in mice. *J Infect Dis* 1984;**149**:924-8.
- 10 Weinstein WM, Onderdonk AB, Bartlett JG, Gorbach SL. Antimicrobial therapy of experimental intraabdominal sepsis. *J Infect Dis* 1975;**132**:282-6.
- 11 Ingham HR, Tharagounet D, Sission PR, Selkon JB, Codd AA. Inhibition of phagocytosis in vitro by obligate anaerobes. *Lancet* 1977;ii:1252-4.
- 12 Lev M, Lev KC, Milford AF. Succinate as a growth factor for *Bacteroides melaninogenicus*. *J Bacteriol* 1971;**108**:175-8.
- 13 Mergenhagen SE, Thonard JC, Sherp HW. Studies on synergistic infections. I. Experimental infections with anaerobic streptococci. *J Infect Dis* 1957;**103**:33-44.
- 14 Gorbach SL. Intestinal microblora. *Gastroenterology* 1971;**60**:1110-29.
- 15 Sutter VL, Finegold SM. Susceptibility of anaerobic bacteria to 23 antimicrobial agents. *Antimicrob Agents Chemother* 1976;**10**:736-52.
- 16 Murray PR, Rosenblatt JE. Penicillin resistant and penicillinase production in clinical isolates of *Bacteroides melaninogenicus*. *Antimicrob Agents Chemother* 1977;**11**:605-8.

Requests for reprints to: Dr Itzhak Brook, Armed Forces Radiobiology Research Institute, Bethesda, Maryland 20814-5145, USA.