Letters to the Editor

Tightly spiral shaped bacteria in gastric antrum: are they urease positive?

Dent *et al* recently reported the occurrence of long spiral organisms in the gastric antrum of patients attending for upper gastrointestinal endoscopy.¹ Of the 1300 patients biopsied, three were found to be infected with this organism which was not recovered by microaerophilic or anaerobic culture.¹

We have observed similar organisms in two of 600 patients attending for endoscopy at Middlemore Hospital over the past 18 months. Five antral biopsy specimens were taken from each patient. Two were sent for histological examination and two for Gram-stained tissue smears and microaerophilic culture. One biopsy specimen was inserted into a CLO-test which detects the urease activity of *Campylobacter pylori*.² Forty six per cent of patients attending for endoscopy at this hospital are infected with *C. pylori*.³

Large numbers of the tightly spiral shaped bacteria were seen in the Gram -stained biopsy smears and histological sections in both patients. *C. pylori* was neither cultured nor seen in the Gram stained biopsy smears of the histological sections. In our hands the above tests are reliable methods for detecting *C. pylori* with sensitivities of 92%, 88%, and 93%, respectively. Both patients were also seronegative for *C. pylori*.⁴ Both patient's

CLO-tests, however, were positive when inspected 24 hours after endoscopy. In the absence of *C. pylori* and because of the large numbers of organisms present we feel the positive urease results were due to this newly recognised organism. This observation differs from that of Dent *et al*, who did not observe positive urease results in their urea broth method.¹ Like Dent *et al*, we did not recover the organism by microaerophilic methods. One patient was rebiopsied but anaerobic and microaerophilic culture on lysed horse blood agar was unsuccessful. This method has been used by Lee *et al* to isolate spiral organisms from cat stomach.⁵

Electron microscopic examination showed the organisms to be similar to the ones reported by Dent *et al*¹ (fig 1). The largest organism was 6.2 µm long, average width was 0.95 µm, with one complete turn of the spiral per 0.95 µm length. The organisms also possess up to 12 sheathed flagella which are attached to the dome end of the cells (fig 2). This arrangement is similar to that seen in the spiral organism in baboon stomachs.⁶ The human and baboon spirals, however, are morphologically different to the cat spiral isolated by Lee *et al*⁵ which possesses regular paired projections along the cell wall.

On the basis of the CLO-test results we predict the human spiral will be urease positive when it is eventually cultured.

A MORRIS,
MR ALI,
LINDY THOMSEN,
M VANDERWEE,

B HOLLIS,
Department of Virology
Wallace Laboratory,
Auckland Public Hospital,
Auckland,
New Zealand

References

- 1 Dent J, McNulty C, Uff J, Wilkinson S, Gear M. Spiral organisms in the gastric antrum. *Lancet* 1987;ii:96.
- 2 Morris A, McIntyre D, Rose T, Nicholson G. Rapid diagnosis of *Campylobacter pyloridis* infection. *Lancet* 1986;i:149.
- 3 Morris A, Ali R, Brown P, Lane M, Palmer R. Treatment of *Campylobacter pylori* gastritis: a pilot study, using Pirenzepine dihydrochloride (Gastrozepin) and three formulations of Colloidal bismuth substrate (De-Nol). *NZ Med J* 1988;101.
- 4 Morris A, Nicholson G, Lloyd G, Haines D, Rogers A, Taylor D. Seroepidemiology of *Campylobacter pylori*. *NZ Med J* 1986; 99:657-9.
- 5 Lee A, Dent J, Hazell S, McNulty C. Origin of spiral organisms in human gastric antrum. *Lancet* 1988;i:300-1.
- 6 Curry A, Jones D, Eldridge J. Spiral organisms in the baboon stomach. *Lancet* 1987;ii:634.

Sensitivity of *Campylobacter pylori* to colloidal bismuth subcitrate

Exposure of *Campylobacter pylori* to some antibacterial agents has been shown to lead to the development of resistant strains; to metronidazole in vivo and in vitro,¹ and to ofloxacin in vivo.² Colloidal bismuth subcitrate (CBS, De-Nol) has been used widely in attempts to eradicate *C. pylori* from the stomach.³ Thus it is important to determine whether, after treatment with CBS, resistant strains can be found, and also whether exposure of *C. pylori* to subinhibitory concentrations of CBS can lead to strains with higher minimum inhibitory concentrations (MICs) to CBS.

A clinical study of 100 people with duodenal ulcer was recently undertaken in our hospital: 50 of the patients took four tablets daily of CBS for eight weeks.⁴ Endoscopic biopsy specimens were obtained before treatment and two, 16, and 42 weeks after treatment, and culture for *C. pylori* was by our standard method.⁴ *C. pylori* was isolated before treatment from all patients, and when CBS treatment had finished, from some patients. To determine MICs of CBS for *C. pylori*, the active component of De-Nol,

Fig 1 Spiral bacteria from human stomach, bar = 1 µm.

Fig 2 Domed end of spiral bacterium near origin of tuft of sheathed flagella, bar = 0.25 µm.