Helicobacter pylori and gastric histamine concentrations

D M M Queiroz, E N Mendes, G A Rocha, J R Cunha-Melo, A J A Barbosa, G F Lima Jr, C A Oliveira

Abstract

An enzyme isotopic assay was used to determine the histamine concentration in the gastric mucosa of patients positive for *Helicobacter pylori* with (n = 11) and without duodenal ulceration (n = 9) and in negative controls (n = 7). A significant difference was observed when the histamine content of *H pylori* negative subjects was compared with that of positive patients. On the other hand, there was no significant difference in histamine concentration between *H pylori* positive patients with duodenal ulceration and those without duodenal ulceration. *H pylori* positive patients with and without duodenal ulceration had significantly lower gastric histamine concentrations than *H pylori* negative subjects. The lower gastric histamine concentration observed in *H pylori* positive patients might be due to increased histamine release which could in turn induce increased gastric acid secretion.

The pathogenesis of duodenal ulceration is not yet fully understood, but it is thought that duodenal ulceration arises from a disturbance in the balance of acid production and mucosal resistance. *Helicobacter pylori* has been observed in up to 100% of subjects with duodenal ulceration and its elimination by antibiotics leads to healing of the ulcer.¹⁻² Although the link between *H pylori* and duodenal ulceration is complex, it has been suggested that the micro-organism could cause ulceration by damaging the duodenal mucosa³ as well as by increasing acid secretion.⁴ According to Levi *et al* the urease produced by *H pylori* transforms urea into ammonia, raising the antral surface pH, which increases serum gastrin and thus acid secretion.⁴ Their hypothesis was contested by Treiber,⁵ who affirmed that *H pylori* does not influence acid secretion.

As histamine has been proposed as the common mediator for all gastric secretions, including gastrin, this study was carried out to determine the histamine concentration in the gastric mucosa of *H pylori* positive and negative patients with and without duodenal ulceration.

Methods

Twenty seven patients presenting for endo-

scopy due to upper gastrointestinal complaints were studied. None of them had received antibiotics, histamine-2 receptor antagonists, or any medication for at least one month before the study. Eleven were *H pylori* positive with endoscopically confirmed duodenal ulceration (eight men mean age 40.8 years); nine were *H pylori* positive with histologically confirmed gastritis but no duodenal ulceration (six men mean age 41.9 years); and seven were *H pylori* negative without duodenal ulceration (six men mean age 42.2 years).

This project was approved by the ethics committee of Hospital das Clinicas, Universidade Federal de Minas Gerais, Brazil, and informed consent was obtained from all patients.

Biopsy specimens were taken from the antral and body mucosa for histological examination: preformed urease test, a smear stained by carbol fuchsin, and culture were used to identify *H pylori*.

Three fragments taken from the gastric body mucosa of each patient were immediately frozen and stored in liquid nitrogen until processing.

Histamine was extracted according to the method of Synder *et al*,⁶ but slightly modified. The tissues were weighed while thawing, immediately immersed in 0.1 M phosphate buffer saline (PBS), pH 7.9 (1 g/20 ml) in a plastic tube, heated for 10 minutes in boiling water, and homogenised with a Teflon pestle tissue homogeniser (size 0, Thomas Scientific, Swedesboro, New Jersey, USA). The tubes were then centrifuged at 5000 rpm at 4°C for 10 minutes and the supernatant was lyophilised.

The fluid was suspended in 100 µl of 0.1 M PBS, pH 7.9, for histamine assay. Aliquots of 10 µl were mixed with 35 µl of a reagent, which contained 0.05 M PBS, 12 µl of rat kidney histamine methyltransferase enzyme preparation,⁷ and 10 µl of 20 µCi/ml solution of S-[methyl-14C] adenosyl-L-methionine (specific activity 5 mCi/mmol), and incubated at 37°C for 90 minutes. Twenty microlitres of 2% N-methylhistamine dihydrochloride (1 in 4 methylhistamine dihydrochloride) (Sigma Chemical Co., St. Louis, Missouri, USA) in 1.5 M perchloric acid was used to stop the reaction. A solution containing 20 µl of 10 NaOH and 500 µl of a mixture of isoamylalcohol and toluene (20:80) was added to each tube. The tubes were capped, shaken for 10 minutes, and centrifuged for 60 seconds on a Beckman Microfuge (Beckman Instru-

Laboratory of Research in Bacteriology, Faculdade de Medicina, Universidade Federal de Minas Gerais—Av. Alfredo Balena, 190/464—CEP 30130—Belo Horizonte, Minas Gerais, Brazil
D M M Queiroz
E N Mendes
G A Rocha

Department of Surgery
J R Cunha-Melo

Department of Pathology
A J A Barbosa

Department of Medicine
C A Oliveira
G F Lima Jr

Correspondence to:
Dr D M M Queiroz

Accepted for publication
18 December 1990

Histological findings in gastric mucosa of *H pylori* positive and negative patients

Mucosal histology	Positive		Negative*
	With duodenal ulceration (n = 11)	Without duodenal ulceration (n = 9)	
Antrum			
ASCG	8	4	0
SCG	3	2	1
ACG	0	3	0
Normal	0	0	6
Body			
ASCG	0	2	0
SCG	2	3	0
ACG	0	1	0
Normal	9	3	7

ASCG: active superficial chronic gastritis; SCG: superficial chronic gastritis; ACG: atrophic chronic gastritis.

*None of them had duodenal ulceration.

ments). Three hundred microlitres of the upper layer was transferred to 5 ml of scintillation fluid for radioactive assay. The results were expressed as counts per minute and the calculation of histamine concentration/sample was based on a standard curve derived from known amounts of histamine.

Statistical analyses were carried out using the two tailed *t* test. Differences of $p < 0.05$ were regarded as significant.

Results

A significant difference was observed when the histamine content (mean, (SD)) of *H pylori* negative subjects (50.29 (14.06) $\mu\text{g/g}$ fresh weight) was compared with that of *H pylori* positive patients with duodenal ulceration (27.76 (9.17) $\mu\text{g/g}$) ($p = 0.0008$) and with gastritis but no duodenal ulceration (29.60 (11.64) $\mu\text{g/g}$) ($p = 0.006$). On the other hand, there was no significant difference in histamine concentration between *H pylori* positive patients with duodenal ulceration and those with gastritis but without duodenal ulceration ($p = 0.69$). *H pylori* positive patients with and without duodenal ulceration showed significantly lower gastric histamine concentration (28.58 (10.11) $\mu\text{g/g}$) than negative subjects ($p = 0.0002$).

The histological findings are shown in the table.

Discussion

The use of a highly sensitive and specific method for determining the histamine concentration of body-type mucosa showed that *H pylori* positive patients with duodenal ulceration as well as those with gastritis and without duodenal ulceration had a significantly lower gastric histamine concentrations than *H pylori* negative subjects. We recently observed this in *H pylori* positive children.⁸ Children with duodenal ulceration, however, presented with histamine concentrations lower than those without duodenal ulceration. We were unable to explain the observed differences between adults and children.

Several hypotheses could be advanced to explain the reduction in gastric histamine concentration that we observed in *H pylori* positive patients. The most likely is that more histamine might be released with a subsequent depletion of stores. This might be induced by an increased release of gastrin following *H pylori*

infection, as shown by Levi *et al.*⁴ Moreover, as mast cells, which store histamine in human gastric mucosa, are also involved in the immunological process, the existence of gastritis might itself be sufficient to disturb histamine metabolism. Oxyntic gastritis was a feature of most of our *H pylori* positive patients without duodenal ulceration, but most of them with duodenal ulceration had normal body mucosa. Nevertheless, all *H pylori* positive patients had antral gastritis, and we do not know how this could disrupt histamine metabolism.

A second hypothesis to explain the lower histamine concentration observed in *H pylori* positive patients is that it could be associated with a decrease in histamine production. On the other hand we found that patients with duodenal ulceration had increased gastric histamine concentrations after treatment with a histamine-2 receptor antagonist (data not shown), although they still had antral gastritis. This seems to indicate that gastritis does not decrease histamine production.

Another possibility is that histamine might have been inactivated. A previous study has shown, however, that histamine methyltransferase activity, the only enzyme able to inactivate histamine, is decreased in patients with duodenal ulceration.⁹

Therefore, our results seem to indicate that lower gastric histamine concentrations in *H pylori* positive patients might be due to the high mobilisation of endogenous mucosal histamine with increased depletion of its stores which could, indirectly, induce an increased gastric acid secretion in these patients.

- 1 Graham DY. *Campylobacter pylori* and peptic ulcer disease. *Gastroenterol* 1989;96:615-25.
- 2 Marshall BJ, Goodwin CS, Warren JR, *et al.* Prospective double-blind trial of duodenal ulcer relapse after eradication of *Campylobacter pylori*. *Lancet* 1989;iii:1437-41.
- 3 Goodwin CS. Duodenal ulcer, *Campylobacter pylori*, and the leaking roof concept. *Lancet* 1988;ii:1467-9.
- 4 Levi S, Beardshall K, Haddad G, Playford R, Ghosh P, Calam J. *Campylobacter pylori* and duodenal ulcers: the gastric link. *Lancet* 1989;ii:1167-8.
- 5 Treiber G. *Campylobacter pylori* and acid secretion. *Lancet* 1989;iii:212.
- 6 Snyder SH, Baldessarini RJ, Axelrod J. A sensitive and specific enzymatic isotopic assay for tissue histamine. *J Pharmacol Exp Ther* 1966;153:544-9.
- 7 Shaff RE, Beaven MA. Increased sensitivity of the enzymatic isotopic assay of histamine measurement of histamine in plasma and serum. *Anal Biochem* 1979;94:425-30.
- 8 Queiroz DMM, Mendes EN, Rocha GA, Barbosa AJA, Carvalho AST, Cunha-Melo JR. *Gut* 1991;32:464.
- 9 Pedem NR, Calachan H, Shepherd DM, Wormsley KG. Gastric mucosal histamine and histamine methyltransferase in patients with duodenal ulcer. *Gut* 1982;23:58-62.