Letters

Necropsy organ weights are largely useless

In 1993 the Royal College of Pathologists published *Guidelines for Post Mortem Reports*.¹ The guidelines may well have led to improvements in necropsy practice but were unreferenced and their evidence base was unstated. They have been used as a gold standard for audit of necropsy reports,² and reiterated in an editorial in this journal.³

Most of the guidelines are sensible but I question the recommendation, in adult necropsies, of routine weighing of organs. Excluding the heart, the weighing of which can provide important information (particularly when the ventricles are weighed separately), organ weights are of little or no value.

The apparent weight of an organ depends on dissection technique and on the accuracy of the weighing balance. In common with other branches of pathology, a numerical result should always be accompanied by the normal range, corrected for the patient's sex, age, and body size. I suspect very few of us comply with this basic rule. However, even if we were to provide reproducibly accurate and referenced weights, they would not be of use to our clinical colleagues, who are accustomed to clinical evaluation of organs, supplemented by imaging techniques in which, at most, a single linear measurement of an organ is given. To be of use, our necropsy practice should reflect these assessments.

The purpose of the necropsy report is to communicate the findings of the necropsy to others, be they clinicians, coroners, or others. We should stop this ritualistic, pseudoscientific practice and concentrate on providing a relevant, meaningful service to our colleagues.

T H W BARKER

Department of Histopathology and Cytopathology,
Norfolk and Norwich Hospital, Brunswick Road,
Norwich, Norfolk

- 1 The Royal College of Pathologists. *Guidelines for post mortem reports*. London: RCPATH, 1993.
- 2 Williams JO, Goddard MJ, Gresham GA, et al. Audit of necropsy reporting in East Anglia. *J Clin Pathol* 1997;50:691-4.
- 3 Start RD, Cross SS. Practice guidelines for necropsy: time for action. *J Clin Pathol* 1996;49:867-9.

We obtained the following comments on this letter:

From Dr R D Start

Dr Barker is correct in his statement that the Royal College of Pathologists' publication *Guidelines for Post Mortem Reports* has led to improvements in necropsy reporting.¹ This is clearly evident within recent reports of the National Confidential Enquiry into Perioperative Deaths (NCEPOD).² The necropsy report guidelines currently provide the only national audit standard for the quality of necropsy reports and I would agree that an appropriate evidence base is desirable. This was one reason behind my suggestion for national practice guidelines for necropsy.³

I am surprised that the routine weighing of organs is in question. Normal organ weight

ranges, corrected for patient sex, age, and body size, are available and can be used if necessary. Some variation may occur with dissection technique but this would be consistent for individual pathologists and could be addressed in any national practice guidelines. Facilities for weighing bodies and organs are fundamental requirements for any mortuary and accuracy is simple to achieve. The suggestion that we stop this "ritualistic, pseudoscientific practice" in order to communicate relevant findings to our colleagues by way of other variables such as a single linear organ measurement is illogical and unacceptable. Linear measurements of focal abnormalities complement gross necropsy findings and allow comparative audit of modern imaging techniques, many of which now give measurements in three dimensions. Supplementary information can also be provided by the dissection of organs in the planes of examination typically seen in modern imaging techniques.

Most clinicians, coroners, and (possibly more importantly) relatives are able to comprehend the concept that organs are abnormal when organ weights are put in the context of normal ranges, particularly if the organ is several times the average normal weight for an individual of similar size, sex, and weight. Although I am unable to provide a specific evidence base to support the use of organ weights, the reasoning behind not providing them is difficult to comprehend, particularly when the most recent major regional audit of necropsy reporting in East Anglia (including Norwich?) found "all pathologists agreed the value of routine weighing of heart, lungs and brain."⁴ I hesitate to suggest that organ weights may be an indirect measure of necropsy quality because through personal experience I have found not only that high quality necropsy reports can be generated from appalling necropsies but also that seemingly accurate organ weights can be determined without removal from cadavers.

More questionable than organ weight measurement is the reported desirability of histological examination in all necropsy cases. There is an increasingly vocal body of opinion within the profession which disagrees with this requirement and the same audit of necropsy reporting in East Anglia provided useful data in relation to the sensible use of histology in necropsy practice.⁵ Such evidence must be incorporated into any future national guidelines for necropsy practice in order that the guidelines reflect the views of most practising pathologists, rather than those of a few interested individuals who perform few, if any, necropsies.

Our principal aim should not be for more necropsies but for better quality necropsies, which are fully supported by a system of formal quality assurance. The Royal College of Pathologists has a major part to play in this process and in recent years has overlooked the necropsy in favour of diagnostic histopathology and cytopathology. The intermittent production of necropsy related guidelines has been useful but a complete reappraisal of all necropsy related matters is urgently required at a national level. Other countries have already addressed these issues⁶ and this is presumably one reason for the commissioning of a Royal College of Pathologists Working Party to examine the current status of necropsy in the United Kingdom.

R D START

Pathology Directorate, Royal Hospital, Calow,
Chesterfield, Derbyshire

- 1 The Royal College of Pathologists. *Guidelines for post mortem reports*. London: RCPATH, 1993.
- 2 The Royal College of Surgeons of London. *National enquiry into perioperative deaths (NCEPOD)*. London: RCS, 1998.
- 3 Start RD, Cross SS. Practice guidelines for necropsy: time for action. *J Clin Pathol* 1996;49:867-9.
- 4 Williams JO, Goddard MJ, Gresham GA, et al. Audit of necropsy reporting in East Anglia. *J Clin Pathol* 1997;50:691-4.
- 5 Williams JO, Goddard MJ, Gresham GA, et al. The use of histopathology in the practice of necropsy. *J Clin Pathol* 1997;50:695-8.
- 6 Hutchins GM, Glenn GC. Practice guidelines for autopsy pathology. *Arch Pathol Lab Med* 1994;118:19-25.

From Dr M J Goddard

In our paper,¹ we included the measurements of major organ weights (heart, lungs, brain, liver, and kidneys) as one of the audit criteria. We do not consider this a gold standard but set this audit standard on the basis of the guidelines of the Royal College of Pathologists, together with a consensus view of the pathologists whose reports were used in the audit.

It is interesting to note that in the audit, this was the area of the internal examination which was done least well, with only 73% of reports recording all five weights in the initial audit, improving to 84% at the time of re-audit. This could be taken to suggest that at least a proportion of pathologists are unwilling to record data that they perceive as meaningless.

As one of the pathologists who reviewed the necropsy reports at the time of the audit, I would have to say that there were very few if any instances where the weight of the organ other than the heart contributed to the quality of the report. I would have to agree that from my own personal viewpoint, together with my impressions gleaned from the audit, the routine weighing of organs serves no useful purpose, provides no information to clinical colleagues, and should cease.

M J GODDARD

Department of Histopathology, Papworth Hospital,
Cambridge University Teaching Hospitals Trust,
Papworth Everard, Cambridge

- 1 Williams JO, Goddard MJ, Gresham GA, et al. Audit of necropsy reporting in East Anglia. *J Clin Pathol* 1997;50:691-4.

Book reviews

The Science of Laboratory Diagnosis. Edited by J Crocker and D Bumett. (£59.95.) Isis Medical Media, 1998. ISBN 1899066 62 4.

The editors have done a marvellous job, more than fulfilling their stated aim of producing a volume describing the multidisciplinary state of modern pathology which will be of interest to a wide range of readers. The book is beautifully produced with excellent colour photographs and line diagrams which clearly explain the practicalities described in the text. I was particularly

impressed by the many tables and flow charts, which can be used as aids to decision making.

All aspects of pathology are covered and it is very easy to find the specific information one needs; the first page of each chapter has its own index, there are summaries of topics at the top of each page, and the general index is detailed and comprehensive. Each subject is clearly introduced with relevant background information. Practical details are easy to follow and alternative methods discussed. Attention is paid to the interpretation of results and to the use of quality controls. At the end of each chapter future directions are covered and there is a useful further reading list. The unifying theme to the book is the application of similar methods to different disciplines. Microscopy and molecular techniques are referred to frequently and it is obvious that a common language is developing between pathologists. This welcome situation will be helped considerably by this work, which should be on the bookshelf of every laboratory. The availability of a version on CD-ROM will increase its appeal.

D M BARNES

Oxford Handbook of Clinical Immunology. By G Spickett. (£19.95.) Oxford University Press, 1999. ISBN: 0 19 262721 X.

Immunologists may well wonder whether there is any need for another volume to complement the many excellent texts that already exist on clinical immunology. Gavin Spickett explains why he wrote the book, saying it was for selfish reasons in that the book that he would like to have had was not available when he was a trainee. Therefore the volume has to be assessed from two points of view: first, does it fill a need in the market, and second, does it meet Dr Spickett's own requirements?

In my opinion the answer to both questions is yes. The volume is up to date, comprehensive, and easily referenced. The text includes information on both clinical and laboratory

immunology and has useful appendices on quality and managerial issues which laboratory trainees will find particularly valuable. Skipping through the volume, I find it is generally very up to date, for example the section on X linked lymphoproliferative disease identifies the cloned gene, information about which was only published in *Nature* in October 1998.

In a volume such as this, one's attention tends to get drawn to areas of one's own particular interest and not surprisingly not all contributions are similarly up to date. I found it surprising that the section on HIV infection did not mention combination antiretroviral therapy, particularly protease inhibitor therapy, or viral load testing in any detail, and also indicated that there was no value in sequential monitoring of CD4 counts once they had fallen below 0.05—information that is clearly incorrect following the advent of new treatments.

However, such criticisms are minor in what is otherwise an excellent volume, and I applaud Dr Spickett's omission of fundamental immunology which has little or no place in such a book. I think that immunology trainees will find this volume invaluable, as will most of their trainers, but whether the market will be larger than the 150 or so individuals that this group comprises remains to be seen. I certainly cannot see many SHOs in general medicine carrying this book alongside the many other small slim volumes that currently sit in RMO's pockets, but I may be proved wrong on this too.

GRAHAM BIRD

Correction

Because of an error in the publishing process, Dr Ibrahim's Best Practice article in the February issue (Guidelines for handling oesophageal biopsies and resection speci-

mens and their reporting, vol 53, no 2, pp 89–94) has been numbered 155 instead of 156. Reprints of this article will be numbered correctly, as will references to the article in our in-house Best Practice advert that appears at the end of each issue. We apologise for this error.

Notice

St Mary's Hospital Campus of ICSM, London

Haematology Courses, Spring 2000

1. Two day course in haematology morphology

St Mary's Hospital, 3–4 April 2000
Suitable for consultants and career grade post holders in haematology. Also valuable for trainees in haematology. CME approved (6+7 CME credits)
Cost £140 including light lunch

2. One day course in histopathology of the bone marrow

St Mary's Hospital, 5 April 2000
Suitable for consultants, career grade post holders, and trainees in haematology and histopathology. CME approved (7 CME credits)
Cost £85 including light lunch

Inquiries in writing or by fax only, to:
The Postgraduate Course Organiser
Academic Surgical Unit
10th Floor, QEQM Wing
St Mary's Hospital
London W2 1NY
Fax No +44 (0)171 886 6314.

Instructions for Authors

Papers for publication should be sent to the Editor, *Journal of Clinical Pathology*, BMA House, Tavistock Square, London WC1H 9JR (tel: 020 7383 6209/6154; fax: 020 7383 6668; email: ndavies@bmjgroup.com). Receipt of manuscripts will be acknowledged by the editorial office.

Submission of a paper will be held to imply that it contains original work not being offered elsewhere or published previously. Manuscripts should be prepared in accordance with the Vancouver style.¹ The Editor retains the right to shorten the article or make changes to conform with style and to improve clarity. All authors must sign the electronic licence form after acceptance.

Failure to adhere to any of these instructions may result in delay in processing the manuscript and it may be returned to the authors for correction before being submitted to a referee.

General

- Authors must submit four copies of the original manuscript typed in double line spacing plus covering letter signed by all authors, consenting to publication. Revised manuscripts should be submitted as printed copy and on disk. A guide to submitting an article on disk will be sent when requesting a revision or on notification of an acceptance. Authors should not submit the original paper on disk.
- The names of the authors, with initials, should be followed by the name of the institution where the work was carried out. An indication of the position held by each author should be given in an accompanying letter to the Editor, and manuscripts should bear the name of one author to whom correspondence should be addressed. If available, a fax number and an email address should be supplied.
- Identifying information should not be published in written descriptions, photographs, or pedigrees unless the information is essential for scientific purposes and the patient (or parent or guardian) gives written informed consent for publication; but patient data should never be altered or falsified in an attempt to attain anonymity. Informed consent should be obtained if there is any doubt. Masking the eyes in photographs of patients is inadequate protection of anonymity (for the full statement see the *BMJ*®).
- Authors should include the names and addresses of four experts whom the authors consider suitable to peer review their work.
- When submitting original manuscripts authors should send a copy of any of their other papers on a similar subject to assure the editors that there is no risk of duplicate publication.
- If requested, authors should produce the data upon which the manuscript is based for examination by the Editor.
- The number of authors should be kept to a minimum and should include only those who have made a contribution to the research: justification should be made for more than five authors. Acknowledgments should be limited to workers whose courtesy or assistance has extended beyond their paid work, and to supporting organisations.
- Please inform us of any grant, business interest, or consultancy that could lead to a conflict of interest.
- Authors should provide up to four keywords/phrases for the index.
- All measurements must be in SI units apart from blood pressure measurements, which

should be in mm Hg, and drugs in metric units.

- Abbreviations should be used rarely and should be preceded by the words in full before the first appearance.
- In the statistical analysis of data 95% confidence intervals should be used wherever appropriate.
- Any article may be submitted to outside peer review and for statistical assessment.
- Animal studies will be considered only if the clinical implications are significant.
- Research studies should have stated local ethical committee approval.
- No free offprints will be provided; reprints may be ordered when the proof is returned.

Original articles

- Papers should be no more than 2000 words long and should report original research of relevance to the understanding and practice of clinical pathology. They should be written in the standard form: abstract; introduction; methods; results; and discussion.
- The journal uses a structured form of abstract in the interests of clarity. This should be short (no more than 250 words) and include four headings: *Aims*—the main purpose of the study; *Methods*—what was done, and with what material; *Results*—the most important results illustrated by numerical data with p values or confidence intervals; and *Conclusions*—the implications and relevance of the results.

Leaders/Editorials

- Leaders and Editorials are published by editorial invitation; unsolicited reviews or commentaries are unlikely to be accepted, though the Editor is always pleased to receive suggestions.

Short reports

- Single case reports of outstanding interest or clinical relevance, short technical notes, and brief investigative studies are welcomed and usually published in the form of a Short/Technical report.
- Length must not exceed 1000 words, including a structured abstract of less than 150 words (unstructured for case reports), up to two figures (together occupying half a column of space) or tables (or one of each) and up to 10 references. If more illustrations are required the text must be reduced accordingly.

Letters

- Letters must be typed in double line spacing, should normally be no more than 500 words, have no more than five references, and must be signed by all authors. Two copies should be provided.

Tables and illustrations

Tables should be presented separately in double line spacing without ruled lines; when presented on disk they should be in a separate file from the text.

- Letters and other marks which are to appear on the face of a photomicrograph should be made on a photocopy: they will be added in the Journal style in the editorial office when the manuscript is accepted.
- Legends for illustrations should be typed with double spacing on a separate sheet. The staining technique used should be stated. Magnifications should be given for electron micrographs but not for light micrographs except in cases where this is important.
- Photographs and photomicrographs should be on glossy paper for half tone reproduction. The printing process requires that prints are unmounted and unbacked, and of high quality, with full tonal scale. Illustrations that will not reproduce well will be returned and this may delay

publication. Areas in which tissue does not appear ("background") should be as near white as possible. Three sets of prints must be supplied with each manuscript. Only salient features should be included to preserve detail.

- Colour reproduction of figures in papers is encouraged and is heavily subsidised by the Journal. Advice on costs and material to be submitted for colour work should be sought from the editorial office. The journal can accept colour images as TIFF files in the following media: zipped or unzipped files on floppy disks, compact disks, or optical disks. A hard copy of the image should be provided.
- If any tables or illustrations submitted have been published elsewhere, written consent to republication should be obtained by the author from the copyright holder (usually the publisher) and the authors. A copy of the letter giving consent must be included.

Descriptions of laboratory methods

- When a manufacturer's method is used in a study with a particular item of equipment or kit of reagents, the source of this method and reference to the scientific literature on which it was based should be given.
- For quantitative methods, information on the sensitivity, precision, and accuracy in the hands of the authors should always be provided. When a well recognised method is used, these requirements could be met simply by providing the references to the methodology and discussing the performance in a recognised current quality assurance scheme. Modifications to methods that have not been previously published should be detailed in the text and supported by evidence of their efficacy.
- It is useful to indicate, either from personal observations or by reference, the working range of an assay and the normal reference range. When information is expressed as mean $\pm$ 2SD, the distribution of the range (normal, skew, or logarithmic) should be stated.

References

- References must be numbered in the order they appear in the text and include all information (Vancouver style; references with more than three authors should give only the first three followed by *et al*):
 - 1 Fletcher CDM, McKee H. Sarcomas - a clinicopathological guide with particular reference to cutaneous manifestations. I. Dermatofibrosarcoma protuberans, malignant fibrous histiocytoma and the epithelial sarcoma of Enzinger. *Clin Exp Dermatol* 1984;9:451-65.
 - 2 Washington JA. Conventional approaches to blood culture. In: Washington JA, ed. *The detection of septicemia*. West Palm Beach, Florida: CRP Press, 1978:41-87.
- References in the text should be identified by arabic numerals in brackets—for example, [1] [2].
- Information from manuscripts not yet accepted, or personal communications may be cited only in the text and not included in the references. References are not checked by us; authors must verify references against the original documents before submitting the article.
 - 1 International Committee of Medical Journal Editors. Uniform requirements for manuscripts submitted to biomedical journals. *BMJ* 1991;302:338-41.
 - 2 International Committee of Medical Journal Editors. Protection of patients' rights to privacy. *BMJ* 1995;311:1272.

Manuscript checklist:

- Is there an abstract?
- Are the abbreviations spelt out?
- Are the measurements in SI units?
- Are the references in Vancouver style?

Revised Mar 2000

Best Practice articles (formerly "Broadsheets") prepared by the Association of Clinical Pathologists

Just published

156 Handling oesophageal biopsies and resection specimens and their reporting 2000 NBN IBRAHIM

Recent Publications

155 Pathological investigation of deaths following surgery, anaesthesia, and medical procedures 1999
RD START, SS CROSS

154 *Helicobacter pylori* 1999 CAM McNULTY, JI WYATT (with correction in June issue)

Other Best Practice articles are still available for purchase

153 The laboratory investigation of vaginal
discharge 1998 KF MACSWEEN, GL RIDGWAY

152 Clinical implications of plasma homocysteine
measurement in cardiovascular disease 1998
RA STILL, IFW MCDOWELL

151 Investigation of dyslipidaemias 1997 AF WINDER,
W RICHMOND, DT VALLANCE

150 Antenatal serological testing and prevention of
haemolytic disease of the newborn 1997 JKM DUGUID

149 Serological diagnosis of gluten sensitive enteropathy
1996 DJ UNSWORTH

148 Laboratory diagnosis of malaria 1996
DC WARHURST, JE WILLIAMS

147 Mycological techniques 1996 KG DAVEY,
CK CAMPBELL, DW WARNOCK

146 Macroscopic examination of prostatic specimens
1995 P HARNDEN, MC PARKINSON

145 Investigation of patients with autoimmune haemolytic
anaemia and provision of blood for transfusion 1995
RJ SOKOL, DJ BOOKER, R STAMPS

*Earlier Broadsheets may still be available
from the author. A full list can be obtained
from the Publications Secretary,
Association of Clinical Pathologists,
189 Dyke Road, Hove, East Sussex
BN3 1TL.*

Prices

INLAND One copy, £2.50; 2–10 copies
(of any one broadsheet or reprint),
£2.00 each; 11–100 copies (of any
one), £1.75 each; 101 plus copies (of
any one), price to be agreed; authors
(over 50 free copies), £1.25 each.
OVERSEAS One copy, \$6.75; 2–10 copies
(of any one broadsheet or reprint),
\$5.25; 11–100 copies (of any one),
\$3.75; 101 plus copies (of any one),
price to be agreed.
Authors \$2.25. Prices include postage
but air mail will be charged extra. Trade
discount 10%. All orders (and all
changes of address of regular
subscribers) should be sent to the
Publishing Manager • *Journal of Clinical
Pathology*, BMJ Publishing Group,
BMA House, Tavistock Square,
London WC1H 9JR.