

p21^{waf} correlates with DNA replication but not with prognosis in invasive breast cancer

U-J Göhring, A Bersch, M Becker, W Neuhaus, T Schöndorf

Abstract

Background/Aims—p21^{waf} plays a central role both in the regulation of the cell cycle and in DNA replication. Accordingly, p21^{waf} is a putative tumour suppressor. The role of p21^{waf} expression in breast cancer is still unclear, particularly with respect to the clinical situation. Therefore, this retrospective study was designed to investigate the value of immunohistochemically detected p21^{waf} expression in invasive breast cancer.

Methods—Cellular expression of p21^{waf} was assessed in 307 breast cancer tissues by immunohistochemistry using the monoclonal antibody, clone 4D10. The data were correlated to established and functional factors of prognosis (age, menopausal status, tumour size, nodal status, tumour grade, receptor status, proliferating cell nuclear antigen (PCNA) expression, Her-2/neu expression, and p53 expression), and to clinical follow up (median observation time, 82 months).

Results—Ninety nine of 307 (32.2%) tumour tissues were considered p21^{waf} positive (nuclear staining). In the entire study group, p21^{waf} expression correlated only with increased PCNA expression (χ^2 test: $p = 0.029$), and with none of the other investigated markers. In node negative patients ($n = 134$), p21^{waf} expression correlated with increased tumour size and increased PCNA expression, whereas the node positive subgroup ($n = 161$) showed no correlation with these parameters (lymphonodectomy was done in 295 women). With respect to clinical outcome, p21^{waf} expression showed a definite favourable trend in both subgroups (N0: p21^{waf} negative, 23 of 87; p21^{waf} positive, nine of 43. N+: p21^{waf} negative, 63 of 107; p21^{waf} positive, 23 of 52), but this observation was not significant ($p > 0.05$). Multivariate analysis for disease free survival as indicated by Cox regression analysis included all factors investigated. The most striking parameters were nodal status (relative risk (RR), 1.74; $p = 0.00001$), receptor status (RR, 0.59; $p = 0.0085$), tumour size (RR, 1.42; $p = 0.02$), and Her2/neu expression (RR, 1.56; $p = 0.033$). p21^{waf} expression was not significant in the multivariate analysis ($p > 0.05$).

Conclusions—p21^{waf} expression is an independent factor but fails to be of prognostic or predictive value in multivariate analysis. These data confirm the hypothesis of a

p53 independent p21^{waf} induction and suggest a functional role in the inhibition of PCNA mediated DNA replication.

(J Clin Pathol 2001;54:866–870)

Keywords: p21^{waf}; proliferating cell nuclear antigen; p53; breast cancer; immunohistochemistry; prognosis

Disorders of cell cycle control are the major causes of cancer. The defective function of regulatory cell cycle elements leads towards increased cell proliferation and, in addition, expansion of genome damaged cells.^{1,2} Aberrations of the cell cycle are often accompanied by overexpressed nuclear kinases, such as the cyclin dependent kinases (CDKs).³ Alternatively, the abnormal function of CDK inhibitors (for example, p16 and p27) or the retinoblastoma gene product (Rb), a central regulatory effector, causes the deregulated proliferative activities of the tumour cell.⁴

p21^{waf} (also termed CIP1 or SDI1) is a nuclear protein with a pivotal role in cell cycle regulation.⁵ It acts as a universal inhibitor of CDKs,⁶ thus directly arresting the cell cycle at the G1/S phase checkpoint. In particular, p21^{waf} mediates p53 induced cell cycle arrest resulting from DNA damage after irradiation.^{7,8} This arrest is important in the process of DNA repair or, alternatively, the switch to apoptosis. Moreover, p21^{waf} is induced independently of p53.⁸ BRCA1 mediated growth arrest operates through p21^{waf} expression.⁹ Differentiation inducing agents such as trans-retinoic acid,¹⁰ growth factors,¹¹ or prostaglandin A2¹² can also initiate p21^{waf} transcription. Cyclin D1 is associated with p21^{waf} expression¹³ and acts as an p21^{waf} inducer.¹⁴ In addition, p21^{waf} is involved in arrest of the cell cycle at the G2 phase, by inhibiting the c-myc oncogene.¹⁵ Thus, p21^{waf} executes its numerous regulatory functions both intrinsic to and separated from the core cell cycle machinery. Accordingly, the p21^{waf} protein possesses tumour suppressive properties.^{16,17}

These basic facts hold true for all cancer types, including carcinomas of the breast. However, the clinical value of p21^{waf} detection in breast tumour tissue remains unknown. In clinical practice, there are no determinants to separate patients with an unfavourable prognosis from those whose tumours are not prone to form occult metastases. Those with an unfavourable prognosis need intensified systemic treatment, whereas for those with a more favourable prognosis chemotherapy should be less intensive or avoided. At present, a large number of patients with node negative tumours are treated unnecessarily, and patients who

Department of Gynecology and Obstetrics, University of Cologne, 50924 Cologne, Germany
U-J Göhring
A Bersch
M Becker
W Neuhaus
T Schöndorf

Correspondence to:
Dr Schöndorf
thomas.schoendorf@medizin.uni-koeln.de

Accepted for publication
1 May 2001




Figure 1 Positive p21^{waf} staining in the nuclei of tumour cells in an infiltrating ductal breast cancer (immunoreactive score, 6).

need dose intensification cannot be identified precisely. More specific tumour characteristics are needed so that patients can be offered individualised treatments according to the phenotype of their tumours.

Because of its functional properties, p21^{waf} is a potential marker. Our retrospective study on primary breast cancer tissues was designed to investigate whether p21^{waf} has prognostic impact and whether it correlates with other markers.

Material and methods

PATIENTS

Our study comprised 307 women with primary infiltrating breast carcinomas (T1–4, N0–2). The patients were treated between 1983 and

Table 1 Clinical, morphological, and biological data of the entire study group T1–4 N0–2 M0 (n = 307)

	p21 ^{waf} negative		p21 ^{waf} positive		Total		p Value
	n	%	n	%	n	%	
Age (years)							
< 50	73	23.8	35	11.4	108	35.2	
> 50	135	44.0	64	20.8	199	64.8	NS
Menopausal status							
Pre	59	19.2	28	9.1	87	28.3	
Peri	28	9.1	16	5.2	44	14.3	
Post	121	39.4	121	39.4	176	57.3	NS
Tumour size (cm)							
< 2	80	26.1	35	11.4	115	37.5	
> 2–5	102	33.2	45	14.7	147	49.9	
> 5	26	8.5	19	6.2	45	14.7	NS
Nodal status							
N0	90	30.5	44	14.9	134	45.4	
1–3 nodes	69	23.4	27	9.2	96	32.5	
> 4 nodes	40	13.6	25	8.5	65	22.0	NS
Tumour grade							
GI	33	10.7	17	5.5	50	16.3	
GII	110	35.8	53	17.3	163	53.1	
GIII	65	21.2	29	9.4	94	30.6	NS
Receptor status							
Negative	66	21.9	27	8.9	93	30.8	
Positive	139	46.0	70	23.2	209	69.2	NS
PCNA							
< 10%	95	31.6	41	13.6	136	45.2	
10–49%	71	23.6	26	8.6	97	32.2	
> 50%	37	12.3	31	10.3	68	22.6	0.029
Her2/neu							
Negative	153	51.3	79	26.5	232	77.9	
Positive	49	16.4	17	5.7	66	22.1	NS
p53							
Negative	123	41.1	57	19.1	180	60.2	
Positive	81	27.1	38	12.7	119	39.8	NS
p21 ^{waf}	208	67.8	99	32.2			

Univariate statistical calculation using χ^2 test.

Receptor status: positive, ER positive and/or PR positive; negative, ER negative and PR negative; detection by immunohistochemistry.

Lymph nodes were removed in 295 cases, steroid hormone receptor status was determined in 302 women, PCNA in 301, Her2/neu expression in 298, and p53 expression in 299 samples.

ER, oestrogen receptor; NS, not significant; PCNA, proliferating cell nuclear antigen; PR, progesterone receptor.

1989 at our department. All women had no evidence of metastases (M0) at the time of diagnosis. The patients underwent mastectomy (225 patients) or tumorectomy with postoperative irradiation (linear accelerator: 50 Gy and 10 Gy boost to tumour bed) of the breast (82 patients). Two hundred and ninety five patients underwent axillary lymphonodectomy with removal of more than nine (≥ 10) nodes to rule out nodal metastatic disease. No adjuvant systemic treatment was administered to node negative patients (n = 134). In the node positive patient subgroup, premenopausal or postmenopausal women without steroid hormone receptors received six cycles of a chemotherapy regimen consisting of a combination of either cyclophosphamide, methotrexate, and 5-fluorouracil (600/40/600 mg/m²) or epirubicin and cyclophosphamide (60/600 mg/m²). Postmenopausal women with positive steroid hormone receptors were treated with 20–30 mg tamoxifen for up to five years.

Physical, laboratory, and apparative checks (mammography, thorax x ray, and abdominal ultrasonography) on patients were carried out regularly as a part of an organised follow up programme. Follow up ranged from 24 to 114 months (median, 82). Histological classification and grading were based on the World Health Organisation criteria (1981) and the suggestions of Bloom and Richardson, respectively. Tumour stage followed the TNM system of UICC. Oestrogen receptor (ER) and progesterone receptor (PR) status were determined by immunohistochemistry (ER: ERICA-System, Abbott, Wiesbaden, Germany; PR: monoclonal antibody mPR1, Dianova, Hamburg, Germany). Steroid hormone receptor status was considered to be positive when ER and/or PR were positive, and negative when both ER and PR were negative. The proliferation marker proliferating cell nuclear antigen (PCNA; NA03; Dianova), the tumour suppressor p53 (MAb 1801; Dianova), and the Her2/neu protein (OPA 01/1; Medac, Hamburg, Germany) were determined immunohistochemically in adjacent sections from the same tumour tissue block. Technical procedures and the immunoreactive scoring system have been described previously.^{18–20}

IMMUNOHISTOCHEMISTRY

Immunohistochemical analyses of p21^{waf} were performed on routinely processed blocks of formalin fixed, paraffin wax embedded surgical specimens of the primary carcinomas. The 3–4 μ m sections of carcinoma tissues were mounted on 3-aminopropyltriethoxysilane (APES) covered glass slides. After drying, paraffin wax was removed with xylene (30 minutes), the sections were rehydrated, and the tissues digested with 0.1% trypsin (15 minutes). A modified three step avidin–biotin complex method to detect the p21^{waf} protein was used. All incubations with antibodies were performed in a moist chamber. The primary monoclonal antibody (clone 4D10, mouse IgG1 subtype; Novocastra, Newcastle upon Tyne, UK) was incubated at 4°C for 24 hours. Second and third antibodies were incubated at room temperature

Table 2 Clinical, morphological, and biological data of node negative women T1–4 N0 M0 (n = 134)

	p21 ^{waf} negative		p21 ^{waf} positive		Total		p Value
	n	%	n	%	n	%	
Age (years)							
< 50 years	34	25.4	13	9.7	47	35.1	
> 50	56	41.8	31	23.1	87	64.9	NS
Menopausal status							
Pre	25	18.7	10	7.5	35	26.1	
Peri	14	10.4	7	5.2	21	15.7	
Post	51	38.1	27	20.1	78	58.2	NS
Tumour size (cm)							
< 2	48	35.8	17	12.7	65	48.5	
2–5	41	30.6	19	14.2	60	44.8	
> 5	1	0.7	8	6.0	9	6.7	0.001
Tumour grade							
GI	15	11.2	10	7.5	25	18.7	
GII	53	39.6	21	15.7	74	55.2	
GIII	22	16.4	13	9.7	35	26.1	NS
Receptor status							
Negative	25	18.9	13	9.8	38	28.8	
Positive	63	47.7	31	23.5	94	71.2	NS
PCNA							
< 10%	45	34.4	20	15.3	65	49.6	
10–49%	33	25.2	11	8.4	44	33.6	
> 50%	9	6.9	13	9.9	22	16.8	0.017
Her2/neu							
Negative	67	51.5	37	28.5	104	80.0	
Positive	20	15.4	6	4.6	26	20.0	NS
p53							
Negative	59	45.0	26	19.8	85	64.9	
Positive	30	22.9	16	12.2	46	35.1	NS
p21 ^{waf}	90	67.2	44	32.8			

Univariate statistical calculation using χ^2 test.

Receptor status: positive, ER positive and/or PR positive; negative, ER negative and PR negative; detection by immunohistochemistry.

Steroid hormone receptor status was determined in 132 women, PCNA in 131, Her2/neu expression in 130, and p53 expression in 131 samples.

ER, oestrogen receptor; NS, not significant; PCNA, proliferating cell nuclear antigen; PR, progesterone receptor.

for 30 minutes. At intervals of different incubations, slides were washed with phosphate

Table 3 Clinical, morphological, and biological data of node positive women T1–4 N+ M0 (n = 161)

	p21 ^{waf} negative		p21 ^{waf} positive		Total		p Value
	n	%	n	%	n	%	
Age (years)							
< 50	34	21.1	22	13.7	56	34.8	
> 50	75	46.6	30	18.6	105	65.2	NS
Menopausal status							
Pre	29	18.0	18	11.2	47	29.2	
Peri	12	7.5	8	5.0	20	12.4	
Post	68	42.2	26	16.1	94	58.4	NS
Tumour size (cm)							
< 2	30	18.6	17	10.6	47	29.2	
> 2–5	57	35.4	25	15.5	82	50.9	
> 5	22	13.7	10	6.2	32	19.9	NS
Tumour grade							
GI	18	11.2	6	3.7	24	14.9	
GII	49	30.4	32	19.9	81	50.3	
GIII	42	26.1	14	8.7	56	34.8	NS
Receptor status							
Negative	39	24.7	14	8.9	53	33.5	
Positive	69	43.7	36	22.8	105	66.5	NS
PCNA							
< 10%	46	29.1	19	12.0	65	41.1	
10–49%	35	22.2	14	8.9	49	31.0	
> 50%	26	16.5	18	11.4	44	27.8	NS
Her2/neu							
Negative	78	50.0	39	25.0	117	75.0	
Positive	28	30.1	11	7.1	39	25.0	NS
p53							
Negative	59	37.8	29	18.6	88	56.4	
Positive	47	30.1	21	13.5	68	43.6	NS
p21 ^{waf}	109	67.7	52	32.3			

Univariate statistical calculation using χ^2 test.

Receptor status: positive, ER positive and/or PR positive; negative, ER negative and PR negative; detection by immunohistochemistry.

Steroid hormone receptor status was determined in 158 women, PCNA in 158, Her2/neu expression in 156, and p53 expression in 156 samples.

ER, oestrogen receptor; NS, not significant; PCNA, proliferating cell nuclear antigen; PR, progesterone receptor.

buffered saline. Antigen–antibody complexes were visualised using 3,3'-diaminobenzidine tetrahydrochloride (10 minutes). Cell nuclei were counterstained with haematoxylin (two minutes). A highly positive breast cancer and a negative control (non-specific immunoglobulins) were used in each run.

Specific staining was evaluated independently by two investigators semiquantitatively, yielding an immunoreactive score (IRS) ranging from 0 to 9. The IRS was calculated by multiplying the number of positive nuclear staining tumour cells (0, none; 1, < 10%; 2, 10–49%; 3, \geq 50% positive tumour cells) by the staining intensity (1, weak; 2, moderate; 3, strong). Tumours were considered p21^{waf} positive with IRS \geq 2. When the scores of the two investigators differed, consensus was reached after examination using a teaching microscope. To demonstrate the reproducibility of p21^{waf} detection, consecutive sections from 40 carcinomas were stained at one week intervals.

STATISTICS

We used SPSS 9.0 for Windows (Statistical Package for the Social Sciences; Munich, Germany) for statistical analysis. The χ^2 test was used for univariate comparison of data, whereas follow up data were analysed using the log rank test. Multivariate analyses were based on the Cox proportional hazards model and calculated relative risks.

Results

The p21^{waf} protein was detected in the nuclei of tumour cells. Intracytoplasmic reactions were very rare and much weaker than nuclear staining. The staining pattern in tumours was heterogeneous, revealing a mixture of positive and negative cells. Among the specimens with specific p21^{waf} staining, most of the tumours expressed p21^{waf} only in up to 20% of the cells. Non-diseased lobular or ductal epithelia found in the tumour periphery did not express detectable amounts of p21^{waf} (fig 1). The staining of different consecutive series showed similar results. Thus, the immunohistochemical detection of p21^{waf} expression was considered reproducible and reliable.

Ninety nine of 307 tumours (32.2%) were considered p21^{waf} positive. This observation was independent of the nodal status (N0, 32.3%; N+, 32.8%).

Table 1 summarises the clinical, morphological, and biological data of the study group. The study group showed the expected distribution of established parameters. There was no significant correlation between p21^{waf} and the parameters studied ($p > 0.05$), except for PCNA expression ($p = 0.029$). One of the major subdivisions of the patients concerned their nodal status. Therefore, the group was further divided into patients with node negative and node positive tumour tissues (tables 2 and 3). There was no significant correlation between p21^{waf} and the other parameters in the node positive subgroup ($p > 0.05$), although p21^{waf} expression correlated with PCNA expression ($p = 0.017$) and larger tumour size ($p = 0.001$) in the node negative subgroup.

Table 4 Number of events (relapses) in the node negative (known follow up $n = 130$) and node positive ($n = 159$) subgroups

	Node negative patients (T1-4 N0 M0)				Node positive patients (T1-4 N1-2 M0)			
	Total	Events	%	<i>p</i> Value	Total	Events	%	<i>p</i> Value
Age (years)								
< 50	47	9	19.2	NS	55	25	45.5	NS
> 50	83	23	27.7		104	61	58.7	
Menopausal status								
Pre	35	6	17.1	NS	46	22	47.8	NS
Peri	21	4	19.1		20	10	50.0	
Post	74	22	29.7		93	54	58.1	
Tumour size (cm)								
< 2	63	14	13.0	NS	47	19	40.4	< 0.00001
2-5	59	16	27.1		80	43	53.8	
> 5	8	2	25.0		32	24	75.0	
Tumour grade								
GI	23	3	13.0	NS	23	9	39.1	< 0.00001
GII	72	20	27.8		80	35	43.8	
GIII	35	9	25.7		56	42	75.0	
Receptor status								
Negative	38	12	31.6	NS	52	38	73.1	0.00001
Positive	90	19	21.1		104	47	45.2	
PCNA								
< 10%	62	13	21.0	NS	65	27	41.5	0.037
10-49%	43	10	23.4		47	28	59.6	
> 50%	22	8	36.4		44	29	65.9	
Her2/neu								
Negative	102	24	23.5	NS	116	56	48.3	0.0017
Positive	25	7	28		38	29	76.3	
p53								
Negative	82	18	22.0	NS	87	40	46.0	0.005
Positive	46	13	28.3		67	44	65.7	
p21 ^{waf}								
Negative	87	23	26.4	NS	107	63	58.9	NS
Positive	43	9	20.9		52	23	44.2	
Total	130	32	24.6		159	86	54.1	

Median observation time for disease free survival, 82 months. Statistical calculation using log rank test.

Univariate statistical calculation using χ^2 test.

Receptor status: positive, ER positive and/or PR positive; negative, ER negative and PR negative; detection by immunohistochemistry.

ER, oestrogen receptor; NS, not significant; PCNA, proliferating cell nuclear antigen; PR, progesterone receptor.

Using the log rank test, tumour size, tumour grade, and the expression of steroid hormone receptors, PCNA, Her2/neu, and p53 correlated significantly with clinical outcome (table 4). In node negative patients, all markers failed to indicate recurrence. Disease free survival was calculated according to Kaplan and Meier (fig 2). There was a tendency to a favourable outcome for node positive patients with detectable p21^{waf} expression, but this was not significant. Similarly, node negative patients seemed to benefit moderately from p21^{waf} expression, although again the results were not significant ($p > 0.05$). Data for overall survival revealed similar results (not shown).

Multivariate analysis for disease free survival, as indicated by Cox regression analysis, found nodal status to be the strongest parameter (relative risk (RR), 1.74; $p = 0.00001$), followed by receptor status (RR, 0.59; $p = 0.0085$), tumour size (RR, 1.42; $p = 0.02$), and Her2/neu expression (RR, 1.56; $p = 0.033$) in the entire study group. In the subgroup of node positive patients, tumour size (RR, 1.83; $p = 0.0004$), steroid hormone receptor status (RR, 0.43; $p = 0.0002$), p53 expression (RR, 1.8; $p = 0.009$), and Her2/neu expression (RR, 1.74; $p = 0.019$) were independent parameters. Node negative patients showed no independent marker ($p > 0.05$).




Figure 2 Kaplan-Meier curves for disease free survival (event = relapse) for (A) node negative ($n = 130$) and (B) node positive ($n = 159$) subgroups. Only trends could be observed (log rank test: $p > 0.05$).

Discussion

In selected breast cancer tissues, p21^{waf} expression rates range from 32% to 57%.^{13 21 22} In our study group, 32.2% of tumours had p21^{waf} expression. With respect to the scoring system used we excluded spotted, weakly stained tumour cells from the p21^{waf} positive group. Thus, p21^{waf} detection in our study confirmed earlier results.

The role of p21^{waf} expression in breast cancer is controversial: whereas some studies reveal an inverse correlation between p21^{waf} expression and the apoptotic marker bcl-2,^{23 24} the tumour suppressor p53,^{22 25} and to histological grading,^{22 26} other reports demonstrate p53 independent p21^{waf} expression^{13 27} and association with high histological grading,^{13 28} positive nodal status,²⁸ and large tumour size.²⁸ It is postulated that p21^{waf} expression is a prognostic marker for relapse free survival and improved overall survival.^{22 25 28} In combination with nodal status, p21^{waf} expression is thought to have predictive value. On the contrary, in more recent studies p21^{waf} expression was not a prognostic factor²⁹ and did not correlate with clinical outcome.^{13 30} Significant correlations were restricted to a lobular subtype.¹³ Elledge and Allred summarised numerous clinical studies with respect to p21^{waf} and concluded that p21^{waf} expression plays a subordinate role in breast cancer prognosis.³¹ Our study confirms these latter results. We show here that p21^{waf} expression is an independent factor in

breast carcinoma progression, but lacks a clear predictive and prognostic relevance. Only trends could be seen with regard to clinical outcome.

Interestingly, in node negative women, p21^{waf} expression correlated significantly with tumour size and PCNA expression (table 2). The subgroup of tumours > 5 cm showed an opposite distribution of p21^{waf} positive and p21^{waf} negative cells. Although more patients need to be investigated, this suggests that p21^{waf} might have a protective effect on nodal involvement during tumour development. This is confirmed by earlier results that show a correlation between p21^{waf} expression and negative nodal status,²⁶ and vice versa.²² Further studies should be designed to prove the hypothesis of p21^{waf} mediated node protection.

One of the main functions of p21^{waf} uncoupled from cell cycle regulation is its inhibitory effect on the proliferation of tumour cells.¹⁸ p21^{waf} expression strongly correlates with Ki67 expression.³⁰ Furthermore, p21^{waf} can affect DNA replication via physically binding to PCNA.³² p21^{waf} disrupts the PCNA-Fen1 complex, thereby prohibiting DNA replication.³³ This inhibitory role is concentrated on the operation of PCNA in DNA replication but not in DNA repair.³⁴ In our study, p21^{waf} expression correlated with increased PCNA expression in node negative patients. To our knowledge, this is the first clinical study to deal with both parameters. We assume that increased PCNA expression induces p21^{waf} expression in primary node negative breast cancer. Consistent with the reports discussed above, this might be a protective event that results in decreased nodal involvement. However, these effects are abolished if tumour cells are capable of inducing nodal spread.

In conclusion, the immunohistochemical detection of p21^{waf} expression is of no use when making decisions about the treatment of breast cancer, although it is useful for understanding the biology of breast carcinogenesis.

We thank the department of pathology, University of Cologne for providing tissue specimens. We are grateful to Mrs J Ruestemeyer for technical assistance. This study was supported by "Köln Fortune" programme, University of Cologne, Faculty of Medicine.

- 1 Sherr CJ. Cancer cell cycles. *Science* 1996;274:1672-7.
- 2 Michalides RJ. Cell cycle regulators: mechanisms and their role in aetiology, prognosis and treatment of cancer. *J Clin Pathol* 1999;52:555-68.
- 3 Fernández PL, Jares P, Rey MJ, et al. Cell cycle regulators and their abnormalities in breast cancer. *J Clin Pathol: Mol Pathol* 1998;51:305-9.
- 4 Jacks T, Weinberg RA. The expanding role of cell cycle regulators. *Science* 1998;280:1035-6.
- 5 El-Deiry WS, Tokino T, Velculescu VE, et al. WAF1, a potential mediator of p53 tumour suppression. *Cell* 1993;75:817-25.
- 6 Xiong Y, Hannon GJ, Zhang H, et al. p21 is a universal inhibitor of cyclin kinases. *Nature* 1993;366:701-4.
- 7 El-Deiry WS, Harper JW, O'Connor PM, et al. WAF1/CIP1 is induced in p53-mediated G₁ arrest and apoptosis. *Cancer Res* 1994;54:1169-74.
- 8 El-Deiry WS, Tokino T, Waldmann T, et al. Topological control of p21^{WAF1/CIP1} expression in normal and neoplastic tissues. *Cancer Res* 1995;55:2910-19.

- 9 Somasundaram K, Zhang H, Zeng Y-X, et al. Arrest of the cell cycle by the tumour-suppressor BRCA1 requires the CDK-inhibitor p21^{WAF1/CIP1}. *Nature* 1997;389:187-90.
- 10 Steinmann RA, Hoffmann B, Iro A, et al. Induction of p21 (WAF1/CIP1) during differentiation. *Oncogene* 1994;9:3389-96.
- 11 Michieli P, Li W, Lorenzi MV, et al. Inhibition of oncogene-mediated transformation by ectopic expression of p21^{Waf1} in NIH3T3 cells. *Oncogene* 1996;12:775-84.
- 12 Gorospe M, Liu Y, Xu Q, et al. Inhibition of G₁ cyclin-dependent kinase activity during growth arrest of human breast carcinoma cells by prostaglandin A₂. *Mol Cell Biol* 1996;16:762-70.
- 13 Lallemand F, Courilleau D, Buquet-Fagot C, et al. Sodium butyrate induces G₂ arrest in the human breast cancer cells MDA-MB-231 and renders them competent for rereplication. *Exp Cell Res* 1999;247:432-40.
- 14 de Jong JS, van Diest PJ, Michalides RJ, et al. Concerted overexpression of the genes encoding p21 and cyclin D1 is associated with growth inhibition and differentiation in various carcinomas. *J Clin Pathol: Mol Pathol* 1999;52:78-83.
- 15 Mitchell KO, El-Deiry WS. Overexpression of c-myc inhibits p21^{WAF1/CIP1} expression and induces S-phase entry in 12-*o*-tetradecanoylphorbol-13-acetate (TPA)-sensitive human cancer cells. *Cell Growth Differ* 1999;10:223-30.
- 16 Chen YQ, Cipriano SC, Arenkiel JM, et al. Tumour suppression by p21^{WAF1}. *Cancer Res* 1995;55:4536-9.
- 17 Jones JM, Cui X-S, Medina D, et al. Heterozygosity of p21^{WAF1/CIP1} enhances tumour cell proliferation and cyclin D1-associated kinase activity in a murine mammary cancer model. *Cell Growth Differ* 1999;10:213-22.
- 18 Göhring U-J, Vierbuchen M, Scharl A. Immunohistochemical detection of p185^{erbB2}—a marker for poor prognosis in primary breast cancer. *Tumourdiagnostik und Therapie* 1993;14:26-31.
- 19 Göhring U-J, Scharl A, Stoffl M, et al. Detection of proliferating cell nuclear antigen in primary breast carcinoma—a proliferation marker as a prognostic factor in nodal-positive patients. *Tumourdiagnostik und Therapie* 1994;15:171-5.
- 20 Göhring U-J, Scharl A, Heckel C, et al. p53 protein in primary breast carcinoma—immunohistochemical detection and clinical value as prognostic factor. *Arch Gynecol Obstet* 1995;256:139-46.
- 21 Jiang M, Shao ZM, Wu J, et al. P21/waf1/cip1 and mdm-2 expression in breast carcinoma patients as related to prognosis. *Int J Cancer* 1997;74:529-34.
- 22 Rey MJ, Fernández PL, Jares P, et al. p21^{WAF1/CIP1} is associated with cyclin D1^{CND1} expression and tubular differentiation but is independent of p53 overexpression in human breast carcinoma. *J Pathol* 1998;184:265-71.
- 23 Upadhyay S, Li G, Liu H, et al. bcl-2 suppresses expression of p21^{WAF1/CIP1} in breast epithelial cells. *Cancer Res* 1995;55:4520-4.
- 24 Bukholm IK, Nesland JM, Kåresen R, et al. Interaction between bcl-2 and p21 (WAF1/CIP1) in breast carcinomas with wild-type p53. *Int J Cancer* 1997;73:38-41.
- 25 McClelland RA, Gee JM, O'Sullivan L, et al. p21(WAF1) expression and endocrine response in breast cancer. *J Pathol* 1999;188:126-32.
- 26 Wakasugi E, Kobayashi T, Tamaki Y, et al. P21 (Waf1/Cip1) and p53 protein expression in breast cancer. *Am J Clin Pathol* 1997;107:684-91.
- 27 Barbareschi M, Pelosio P, Caffo O, et al. Cyclin-D1-gene amplification and expression in breast carcinoma: relation with clinicopathologic characteristics and with retinoblastoma gene product, p53 and p21^{WAF1} immunohistochemical expression. *Int J Cancer* 1997;74:171-4.
- 28 Caffo O, Doglioni C, Veronese S, et al. Prognostic value of p21(WAF1) and p53 expression in breast carcinoma: an immunohistochemical study in 261 patients with long-term follow-up. *Clin Cancer Res* 1996;2:1591-9.
- 29 Reed W, Florenes VA, Holm R, et al. Elevated levels of p27, p21 and cyclin D1 correlate with positive oestrogen and progesterone receptor status in node-negative breast carcinoma patients. *Virchows Arch* 1999;435:116-24.
- 30 Mathoulin-Portier MP, Viens P, Cowen D, et al. Prognostic value of simultaneous expression of p21 and mdm2 in breast carcinomas treated by adjuvant chemotherapy with anthracycline. *Oncol Rep* 2000;7:675-80.
- 31 Elledge RM, Allred DC. Prognostic and predictive value of p53 and p21 in breast cancer. *Breast Cancer Res Treat* 1998;52:79-98.
- 32 Flores-Rozas H, Kelman Z, Dean FB, et al. Cdk-interacting protein 1 directly binds with proliferation cell nuclear antigen and inhibits DNA replication catalyzed by the DNA polymerase δ holoenzyme. *Proc Natl Acad Sci U S A* 1994;91:8655-9.
- 33 Chen J, Chen S, Saha P, et al. p21^{Cip1/Waf1} disrupts the recruitment of human Fen1 by proliferating-cell nuclear antigen into DNA replication complex. *Proc Natl Acad Sci U S A* 1996;93:11597-602.
- 34 Li R, Waga S, Hannon GJ, et al. Differential effects by the p21 CDK inhibitor on PCNA-dependent DNA replication and repair. *Nature* 1994;371:534-7.