2. **Groisman GM**, Amar M, Meir A. Expression of the intestinal marker Cdx2 in the columnar-lined oesophagus with and without intestinal (Barrett's) metaplasia. *Mod Pathol* 2004;**17**:1282–8.
3. **Suh E**, Traber PG. An intestine-specific homeobox gene regulates proliferation and differentiation. *Mol Cell Biol* 1996;**16**:619–25.
4. **Silberg DG**, Swain GP, Suh ER, *et al*. Cdx1 and cdx2 expression during intestinal development. *Gastroenterology* 2000;**119**:961–71.
5. **Phillips RW**, Frierson HF, Moskaluk CA. CDX2 as a marker of epithelial intestinal differentiation in the esophagus. *Am J Surg Pathol* 2003;**27**:1442–7.
6. **Kazumori H**, Ishihara S, Kinoshita Y. Roles of caudal-related homeobox gene Cdx1 in oesophageal epithelial cells in Barrett's epithelium development. *Gut* 2009;**58**:620–8.
7. **Moons LMG**, Bax DA, Kuipers EJ, *et al*. The homeodomain protein CDX2 is an early marker of Barrett's oesophagus. *J Clin Pathol* 2004;**57**:1063–8.
8. **Eda A**, Osawa H, Satoh K, *et al*. Aberrant expression of CDX2 in Barrett's epithelium and inflammatory esophageal mucosa. *J Gastroenterol* 2003;**38**:14–22.
9. **Odze RD**. Diagnosis and grading of dysplasia in Barrett's oesophagus. *J Clin Pathol* 2005;**59**:1029–38.
10. **Morris CD**, Armstrong GR, Bigley G, *et al*. Cyclooxygenase-2 expression in the Barrett's metaplasia-dysplasia-adenocarcinoma sequence. *Am J Gastroenterol* 2001;**96**:990–6.
11. **Kerkhof M**, van Dekken H, Steyerberg EW, *et al*. Grading of dysplasia in Barrett's oesophagus: substantial interobserver variation between general and gastrointestinal pathologists. *Histopathology* 2007;**50**:920–7.
12. **Montgomery E**, Bronner MP, Goldblum JR, *et al*. Reproducibility of the diagnosis of dysplasia in Barrett esophagus: a reaffirmation. *Hum Pathol* 2001;**32**:368–78.
13. **Reid BJ**, Blount PL, Rubin CE, *et al*. Predictors of progression to malignancy in Barrett's esophagus: endoscopic, histologic and flow cytometric follow-up of a cohort. *Gastroenterology* 1992;**102**:1212–19.
14. **Lord RVN**, Brabender J, Wickramasinghe K. Increased CDX2 and decreased PITX1 homeobox gene expression in Barrett's oesophagus and Barrett's-associated adenocarcinoma. *Surgery* 2005;**138**:924–30.
15. **Villanacci V**, Rossi E, Zambelli C, *et al*. COX-2, CDX2, and CDC2 immunohistochemical assessment for dysplasia-carcinoma in Barrett's esophagus. *Digest Liver Dis* 2007;**39**:305–11.
16. **Mallo GV**, Rechreche H, Frigerio J-M, *et al*. Molecular cloning, sequencing and expression of the mRNA encoding human cdx1 and cdx2 homeobox: down-regulation of cdx1 and cdx2 mRNA expression during colorectal carcinogenesis. *Int J Cancer* 1997;**74**:35–44.
17. **Yun-Qing B**, Hiroshi Y, Yoshimitsu A, *et al*. Ectopic expression of homeodomain protein cdx2 in intestinal metaplasia and carcinomas of the stomach. *Cancer Lett* 2002;**176**:47–55.
18. **Bonhomme C**, Duluc I, Martin E, *et al*. The Cdx2 homeobox gene has a tumour suppressor function in the distal colon in addition to a homeotic role during gut development. *Gut* 2003;**52**:1465–71.
19. **Guo RJ**, Suh ER, Lynch JP. The role of Cdx proteins in intestinal development and cancer. *Cancer Biol Ther* 2004;**3**:593–601.
20. **Liu Q**, Teh M, Ito K, *et al*. CDX2 expression is progressively decreased in human gastric intestinal metaplasia, dysplasia and cancer. *Mod Pathol* 2007;**20**:1286–97.

Correction

Ze-Peng Du, Zhuo Lv, Bing-Li Wu, *et al*. Neutrophil gelatinase-associated lipocalin and its receptor: independent prognostic factors of oesophageal squamous cell carcinoma. *J Clin Pathol* 2011;**64**:69–74. Correspondence for this article should be addressed to: Professor Li-Yan Xu, Institute of Oncologic Pathology, Medical College of Shantou University, No 22, Xinling Road, Shantou 515041, China; liyanxu1130@yahoo.com.cn.

J Clin Pathol 2011;**64**:113. doi:10.1136/jcp.2010.083907corr1