



Editor's choice  
Scan to access more  
free content

# Differentiation of oncocytoma from chromophobe renal cell carcinoma (RCC): can novel molecular biomarkers help solve an old problem?

Keng Lim Ng,<sup>1,2,3</sup> Retnagowri Rajandram,<sup>1,3,4</sup> Christudas Morais,<sup>1</sup> Ning Yi Yap,<sup>3</sup> Hema Samaratunga,<sup>5</sup> Glenda C Gobe,<sup>1</sup> Simon T Wood<sup>2</sup>

<sup>1</sup>Centre for Kidney Disease Research, School of Medicine, The University of Queensland, Translational Research Institute, Brisbane, Australia

<sup>2</sup>Department of Urology, Princess Alexandra Hospital, Brisbane, Australia

<sup>3</sup>Department of Surgery, Faculty of Medicine, University of Malaya, Kuala Lumpur, Malaysia

<sup>4</sup>University Malaya Cancer Research Institute, Kuala Lumpur, Malaysia

<sup>5</sup>Aquesta Pathology, Brisbane, Australia

## Correspondence to

A/Professor Glenda Gobe, Centre for Kidney Disease Research, School of Medicine, Translational Research Institute, Kent Street, Woolloongabba, Brisbane 4102, Australia; g.gobe@uq.edu.au

Received 1 August 2013

Revised 30 September 2013

Accepted 1 October 2013

Published Online First

29 October 2013

## ABSTRACT

Standard treatment of renal neoplasms remains surgical resection, and nephrectomy for localised renal cell carcinoma (RCC) still has the best chance of cure with excellent long-term results. For smaller renal masses, especially stage T1a tumours less than 4 cm, nephron-sparing surgery is often employed. However, small incidentally detected renal masses pose an important diagnostic dilemma as a proportion of them may be benign and could be managed conservatively. Renal oncocytoma is one such lesion that may pose little risk to a patient if managed with routine surveillance rather than surgery. Additionally, lower-risk RCC, such as small chromophobe RCC, may be managed in a similar way, although with more caution than the renal oncocytomas (RO). The ability to differentiate ROs from chromophobe RCCs, and from other RCCs with a greater chance of metastasis, would guide the physician and patient towards the most appropriate management, whether nephron-sparing surgical resection or conservative surveillance. Consistent accurate diagnosis of ROs is likely to remain elusive until modern molecular biomarkers are identified and applied routinely. This review focuses on the differentiation of renal oncocytomas and chromophobe RCCs. It summarises the history, epidemiology and clinical presentation of the renal neoplasms, explains the diagnostic dilemma, and describes the value, or not, of current molecular markers that are in development to assist in diagnosis of the renal neoplasms.

## INTRODUCTION

The incidence of renal tumours has been increasing steadily in Europe, USA and Australia over the past three decades.<sup>1</sup> The widespread use of cross-sectional imaging has increased the detection of incidental smaller tumours,<sup>2</sup> while the 20–30% incidence of advanced tumours has remained relatively constant.<sup>3</sup> Despite current imaging techniques and the availability of renal lesion biopsy, most contemporary surgical series continue to report significant rates of benign lesions among resected small renal masses.<sup>2–4</sup> Preoperative biopsy of these small lesions is not widely employed, and one contributing factor is potential diagnostic uncertainty in the differentiation of benign renal oncocytomas (RO) from malignant chromophobe renal cell carcinomas (chRCC)<sup>5</sup> and, as an added difficulty, eosinophilic clear-cell RCCs (ccRCC). Consequently, there is a group of small renal lesions where increased confidence in characterisation may defer or obviate the need for surgical

intervention. ROs and small chRCCs are two such lesions.

ChRCCs, although having a more favourable prognosis than other RCC subtypes, is a malignant tumour with the potential for metastatic spread and death. By comparison, there appears to be only one confirmed case report of metastases from ROs.<sup>6</sup> Thus, due to its benign nature, ROs can usually be monitored and treated expectantly. Similarly, small renal masses found to be chRCCs may, in some situations, be suitable for active surveillance rather than immediate resection or ablation. ROs and chRCCs are often considered to be extremities of the same morphological spectrum.<sup>7</sup> Proper differentiation largely relies on H&E histochemistry of sections, and an experienced histopathologist to discern the characteristic histomorphologic features between the two entities. Immunohistochemistry is used in selected instances. Electron microscopy was commonly performed in the past, but is done only in rare cases now, given the prominent overlap of staining patterns. There is also the coexistence of ROs with chRCCs seen in sporadic cases of hybrid tumours, renal oncocytosis and Birt-Hogg-Dube (BHD) syndrome. Differentiation of ROs and chRCCs, especially as small renal masses, from other more sinister forms of RCCs, like ccRCCs, is also important for the appropriate management of these patients.

## HISTORY

RO was first described by Zippel in 1942 as a neoplasm entirely composed of large eosinophilic cells called oncocytes.<sup>8</sup> Later, in 1976, Klein and Valensi<sup>9</sup> identified another 13 cases as a distinct clinical pathological entity with a typical benign histological presentation and clinical course. RO was originally thought to derive from renal proximal tubules, but most pathologists now suggest a distal origin,<sup>10</sup> most likely arising from intercalated cells of collecting ducts. The first description of chRCCs, as distinct from ccRCCs, was made by Theones *et al* in 1985,<sup>11</sup> and a year later, they added the chRCC subtype to the classification of renal tumours.<sup>12</sup> The cell characteristic had been described prior to the 1985 publication but only in experimentally induced adenomas in animals. The chromophobe cells had slightly opaque or finely reticular cytoplasm that resisted staining with haematoxylin and eosin. They were able to be distinguished from ccRCCs by a strongly positive reaction within their cytoplasm to Hale's colloidal iron, and a weaker positive reaction with Alcian Blue, a

**To cite:** Ng KL, Rajandram R, Morais C, *et al.* *J Clin Pathol* 2014;**67**:97–104.

distinction that has since been found to be unreliable. The authors, however, made a step forward for classification of RCCs by suggesting that the descriptive term 'light cell' RCC should be discarded and replaced by either 'clear cell' or 'chromophobe cell' as appropriate. They pointed out that chromophobe cell tumours were likely to have a different derivation from ccRCCs and other RCCs, and that they may also have a different prognosis, a fact that has since been established. Since the description of chRCCs came a decade later than ROs, there were many instances in that era where renal tumours, which were likely to be chRCCs, were described as ROs. This may have contributed to the confusion surrounding the original recognition of the benign nature of ROs.

## EPIDEMIOLOGY

Renal tumours are highly heterogeneous with at least 16 known subtypes, of which four subtypes predominate.<sup>13–14</sup> CcRCCs, arising from the proximal tubular epithelial cells, is the most common subtype constituting 70–80% of RCCs, followed by papillary (10–15%), chRCCs (5%) and collecting duct RCCs (<1%).<sup>15–16</sup> RO accounts for approximately 3–7% of all adult renal neoplasms. The peak age of incidence for detection of ROs tends to be in the seventh decade of life. For chRCCs, the peak incidence occurs in the sixth decade. For cases of RO, men seem to be affected twice as often as women. For chRCCs, the disease tends to affect men and women equally.<sup>17</sup>

ROs and chRCCs can develop as either sporadic or familial forms, and both can be associated with distinct genetic mutations. The majority of ROs and chRCCs occur as sporadic cases.<sup>18–19</sup> There is also the occasional occurrence of familial renal cancers of oncocytoma with BHD syndrome. Familial oncocytoma is due to partial or complete loss of multiple chromosomes. BHD syndrome is an autosomal dominant inherited syndrome with the BHD gene locus located in the short arm of chromosome 17.<sup>20–21</sup> This syndrome is characterised by fibrofolliculomas, lung cysts that can lead to spontaneous pneumothoraxes, and various subtypes of renal tumours including hybrid tumours, ROs, chRCCs and ccRCCs.


Sometimes in rare instances, patients can present with renal oncocytosis. Renal oncocytosis was first described in 1982<sup>22</sup>: multiple and bilateral oncocytic nodules and a spectrum of oncocytic changes are found diffusely throughout the renal parenchyma. A large series investigating renal oncocytosis revealed that hybrid development of ROs and chRCCs was most common.<sup>23</sup>

## CLINICAL PRESENTATION

Generally, patients with ROs tend to be asymptomatic and present incidentally following cross-sectional imaging for an unrelated complaint. Similarly, the majority of patients with chRCCs present incidentally with asymptomatic renal masses.<sup>24</sup> Less commonly, chRCCs may present with local symptoms of haematuria, flank mass and loin pain, and constitutional symptoms of weight loss and loss of appetite.<sup>18</sup> chRCCs can also present with paraneoplastic syndrome and metastases with predilection to the liver.<sup>25</sup> In the largest published series to date, chRCCs present with metastases at a rate of 1.3%.<sup>24</sup> Generally, patients with chRCCs tend to present in less advanced stages (I and II), less frequently with metastases and are usually of better performance status<sup>25</sup> compared with other subtypes of RCCs. It should be noted, however, that the local and constitutional symptoms for chRCCs are similar to those seen for other RCCs.

ROs will almost always follow a benign clinical course with no significant risk of metastases, whereas malignant chRCCs can subsequently progress to metastases. There have been a few isolated case reports of metastatic ROs on initial presentation or following resection of the ROs,<sup>13–26</sup> however, these case reports have not been substantiated with proper histopathological confirmation of the metastatic deposits, except for one liver metastasis.<sup>6</sup> Therefore, the distinction of almost exclusively benign ROs from malignant, potentially metastatic, chRCCs is needed to guide the management of these often difficult-to-separate entities.

Renal tumours can be detected by radiological imaging using ultrasonography, CT, MRI and positron emission tomography (PET). Usually, following the suspicion of a renal mass, either clinically or via ultrasound, a 4-phase CT scan will be performed to delineate its nature. Multiphase CT scans can clearly delineate the renal tumour, its local extension to surrounding tissues and detect any metastases to regional lymph nodes or other organs. Cases of small renal masses (lesions <4 cm) detected incidentally are increasing in incidence largely owing to the widespread use of ultrasound and CT scans. Generally, there is no accurate differentiation between benign and malignant renal lesions using CT scans (except angiomyolipoma), but retrospectively, about 20% of these small renal masses will be found to be benign lesions. Percutaneous biopsy of these small renal masses provides an enticing strategy to identify lesions of no or low malignant



**Figure 1** (A) H&E-stained section of an example of eosinophilic variant of chromophobe renal cell carcinoma, showing typical large, pale, polygonal cells with prominent cell membranes. Nuclei tend to be irregular and wrinkled, and cells are sometimes binucleated (asterisks). Perinuclear clearing can be prominent. (B) H&E stained section of an example of renal oncocytoma, showing large oncocytes with densely eosinophilic cytoplasm. Cells are round to polygonal and nuclei are round and monotonous. Nucleoli are small and inconspicuous.

**Table 1** Comparison of macroscopic, microscopic and ultrastructural features for oncocytoma and chromophobe renal cell carcinoma

Features	Oncocytoma	Chromophobe RCC
Macroscopic	Well circumscribed, tan or mahogany brown, sometimes with a central stellate scar <sup>30</sup>	Usually circumscribed, homogenous, light brown, beige, yellow or tan colour. <sup>46</sup>
Microscopic	Cells arranged in a nested or organoid pattern, but tubular, trabecular or solid structure can also be seen. <sup>54</sup>	Variants: classic, eosinophilic and mixed.
Cytoplasm	Granular eosinophilic cytoplasm	Cells arranged in sheets, with distinct or accentuated cell borders. <sup>56</sup>
Nuclei	Round, uniform nuclei <sup>55</sup>	Granular eosinophilic (eosinophilic variant) or pale, reticular and almost transparent appearance (classic). <sup>57</sup> Presence of perinuclear halos, wrinkled nuclei. <sup>57</sup>
Ultrastructural	Abundant mitochondria with lamellar or focally stacked cristae. Absent or sparse vesicles. <sup>55</sup>	Scant mitochondria with tubule-vesicular cristae. Abundant microvesicles between mitochondria. <sup>46</sup>

potential, however, widespread uptake of biopsy into clinical practice has been limited.

Predicting whether a small renal mass is malignant, based on its growth velocity, has been reported, but there is no good correlation of malignancy with growth rate.<sup>16</sup> A recent meta-analysis of small renal masses which included benign and

malignant lesions, showed a mean growth rate of 0.28 cm annually (range 0.09 to 0.86) for small renal masses followed with imaging.<sup>14</sup> ROs increase in size with variable velocity, with one case series reporting an observed growth rate of 0.20 cm annually.<sup>15</sup> The largest pool of 33 biopsy-proven benign ROs demonstrated a growth rate similar to reported growth rates for RCCs, thus highlighting again that observation of growth cannot distinguish between the benign or malignant nature of such lesions.<sup>16</sup> The locality and size of tumours may also be variable. Uncommonly, there have been case reports of large ROs (25×15×12 cm),<sup>27</sup> but the average size is normally around 4.9±2.7 cm.<sup>28</sup> Published reports worldwide show that ROs can be multifocal in 6–11%,<sup>29–30</sup> and bilaterality was reported in about 3–5%.<sup>29–31</sup> By comparison, the median size of chRCCs is about 6.0 cm,<sup>18</sup> which is larger compared with other subtypes of RCCs.<sup>32</sup> Multifocality of chRCCs is usually around 10–12%.<sup>33</sup>

### DIAGNOSTIC DILEMMA

The increasing use of CT scans for small renal masses has led to a diagnostic dilemma of accurately characterising the nature of these renal lesions and their subsequent management. Typically, on CT scans, RCCs are solid heterogeneous masses with contrast enhancement showing areas of patchy uptake of contrast. Locally advanced tumours may directly invade the adrenal gland, renal vein, inferior vena cava and regional lymph nodes. ChRCCs usually demonstrate homogeneous enhancement, whereas ccRCCs, papillary and collecting duct RCCs tended to show heterogeneous or predominantly peripheral enhancement. Calcification was seen more commonly in chRCCs than in

**Table 2** Histochemistry and immunohistochemistry (IHC) to differentiate chromophobe renal cell carcinomas (RCC) and renal oncocytomas

Method	Number of patients	Success as biomarker	Reference
Hale's colloidal iron stain	28 cases (11 chRCC, 12 RO, 6 ccRCC)	Colloidal iron was diffusely and strongly positive in 9/11 of chRCC, focally and weakly positive in 5/12 of RO, and negative in all granular cell variants of ccRCC (0/6).	58
Modified Mowry's colloidal iron stain better characterised chRCC	62 cases (14 chRCC, 19 RO, 11 ccRCC, 7 eosinophilic variants of pRCC)	Positive colloidal iron stain was not limited to chRCC, however a diffuse and strong, reticular staining pattern was observed only in chRCC (100%). Staining patterns less consistent in all other renal neoplasms. Most RO (84%) had focal, weak, fine dust-like positivity. 100% ccRCC had focal, coarse, droplet-like positivity.	59
	76 cases (30 ccRCC, 16 pRCC, 21 chRCC, 8 RO, 1 cdRCC)	Fine reticular cytoplasmic pattern with perinuclear halo (87.5% chRCC; 16% ccRCC). 12.5% RO had focal, coarse, cytoplasmic staining without perinuclear halo.	45
CD10	76 cases (30 ccRCC, 16 pRCC, 21 chRCC, 8 RO, 1 cdRCC)	CD10 positive, 79% ccRCC, 6.3% chRCC and 0% RO. CD10 reactivity favours ccRCC, and the absence of CD10 in RO shows CD10 could differentiate between chRCCs and RO in a panel of biomarkers.	45
Outcome of CD10 to distinguish between chRCC and ROs is variable.	83 cases (22 chRCC, 17 RO, and 45 ccRCC)	CD10 positive, ccRCC (91%), chRCC (45%) and RO (29%).	47
	28 cases (11 chRCC, 12 RO, 6 ccRCC)	CD10 positive, 100% ccRCC, 72% chRCC and 58% RO. Not useful as a biomarker.	58
RCC marker (RCCma)	76 cases (30 ccRCC, 16 pRCC, 21 chRCC, 8 RO, 1 cdRCC)	RCCma, positive in 62.5% ccRCC, 12.5% RO, but negative in chRCC. Holds potential as part of a panel to differentiate between chRCC and RO.	45
RCCma is a relatively new IHC marker that has variable results.	Renal cell neoplasm TMA (30 RO, 18 chRCC, 64 ccRCC, 50 pRCCs, 31 RO)	RCCma, positive in most RCC with granular/eosinophilic features. ccRCC (71%), pRCC (76%), negative in RO.	60
	328 samples (256 ccRCC, 27 pRCC, 28 chRCC, 5 cdRCC, 5 unclassified RCC, 7 RO)	RCCma was negative in chRCC but was positive in 3/7 RO.	61
	29 cases (11 chRCC, 12 RO, 6 ccRCC)	RCCma was observed in more than 80% of ccRCCs but was negative in all chRCCs and RO.	58
Vimentin	76 cases (30 ccRCC, 16 pRCC, 21 chRCC, 8 RO, 1 cdRCC)	Vimentin positive, 95% ccRCC, 6.3% chRCC, 12.5% RO. Negative staining for vimentin, chRCC or RO.	45
	83 cases (22 chRCC, 17 RO, 45 ccRCC)	Vimentin positive exclusively in ccRCCs.	47
	Renal cell neoplasm TMAs (30 RO, 18 chRCC, 64 ccRCC, 50 pRCC, 31 RO)	Positive in most RCC with granular/eosinophilic features (ccRCC 78%, pRCC 85%). Negative in RO and chRCC.	60

ccRCC, clear cell renal cell carcinoma; cdRCC, collecting duct renal cell carcinoma; chRCC, chromophobe renal cell carcinoma; pRCC, papillary renal cell carcinoma; RO, renal oncocytoma.

**Table 3** Biomarkers used to differentiate chromophobe renal cell carcinoma (RCC) (chRCC) from oncocytoma (RO)

Method	Number of patients	Significance of success as biomarker	Reference
BCA2	158 patients (104 ccRCC, 8 chRCC, 2 pRCC, 38 RO, 6 oncocytic neoplasms)	All RO and oncocytic neoplasms, which favour RO, were positive for BCA2 while all RCC were negative, including chRCC.	50
C-kit (encodes the membrane-bound tyrosine kinase KIT)	mRNA levels, 17 chRCC, 20 RO from cDNA microarrays IHC analysis, 226 renal tumours in TMAs (40 chRCC, 41 RO, 40 ccRCC, 29 renal angio-myolipoma, 21 pRCC).	Significant increment of c-kit mRNA and overexpression of KIT protein by IHC in chRCC and RO, hence low potential for differentiating between the two types. However, there was potential for differentiating chRCC/RO from the other renal cell tumours (ccRCC and pRCC).	62
EMA	86 retrospective nephrectomy specimens (15 ccRCCs, 15 pRCCs, 15 chRCCs, 10 ROs, 6 cdc)	EMA was positive in chRCC (75–100%), ccRCC (50–77%) and oncocytomas (51–86%), showing no major promise as a marker. (Comparison made with 3 tubulocystic carcinoma, 3 renal medullary carcinoma, 3 mucinous tubular and spindle cell carcinoma, 4 metanephric adenoma, 12 invasive high-grade urothelial carcinoma)	53
	76 cases (30 ccRCC, 16 pRCC, 21 chRCC, 8 RO, 1 cdc)	EMA was positive in 100% of ChRCCs, 100% of ROs and 75% of ccRCC. So, we concluded that EMA is not a good marker for the differentiation of renal tumours.	45
Carbonic anhydrase IX (CA IX)	TMAs, 20 cases of each ccRCC, chRCC, pRCC and RO	CA IX was highly sensitive for ccRCCs (90% positivity) and was negative in all other renal epithelial tumours except for 1 chRCC.	48
Galectin-3	TMAs, 20 cases of each ccRCC, chRCC, pRCC and RO	Galectin-3 found mostly in renal tumours with oncocytic features, including RO (100%) and chRCCs (89%). May hold small promise to distinguish these from other RCC.	48
Glutathione S-transferase alpha (GST- $\alpha$ )	22 chRCC, 17 RO, 45 ccRCC	GST- $\alpha$ exclusively observed in ccRCCs.	47
KIT (CD117)	256 ccRCC, 29 chRCC, 25 pRCC, 6cdc, 6 unclassified RCC, 7 RO, 20 UC, 7 NB, 2 AM	83% chRCCs and 71% RO had membranous immunoreactivity for KIT, while none of the other RCC or the angiomyolipomas expressed. Cannot be used to differentiate chRCC and RO.	63
	11 chRCCs, 12 RO, 6 ccRCC	KIT was a very sensitive marker for both chRCC and RO, but not useful to differentiate between the two. KIT with RCCma may be useful when trying to differentiate ccRCCs from chRCCs or ROs.	58
	22 chRCC, RO & ccRCC	CD117, strongly expressed in chRCC (82%) and RO (100%), whereas none of the ccRCCs were immunoreactive.	47
CD15	10 ccRCC, pRCC, chRCC and RO	CD15 was able to distinguish between chRCCs and RO. 7/10 RO (70%) stained positive for CD15 and none of the chRCC stained for CD15.	64
MAGE-A3/4 cancer testis antigen/CTA	35 patients (17 RO, 18 chRCC)	88% RO stained positively for MAGE-A3/4; 39% chRCC stained positively	51
RON proto-oncogene, encoding a receptor tyrosine kinase,	TMAs (55 RO, 52 chRCCs). 15 & 5 conventional sections of RO & chRCC were also analysed 11 chRCC, 12 RO, 6 ccRCCs	69 of 70 RO and 55 of 57 chRCC had strong, diffuse cytoplasmic stain. 11/11 chRCCs, 12/12 RO, but only 3/6 of ccRCC.	65 58
NY-ESO-1 CTA	35 patients (17 RO, 18 chRCC)	15/17 RO stained positive, and 6/18 chRCC were positive.	51
Interphase fluorescence in situ hybridisation (FISH)	11 chRCC, 12 RO, compared with conventional metaphase cytogenetics by karyotyping.	RO often show normal DNA content by interphase and metaphase analyses. The loss of 2 or more of chromosomes 1, 2, 6, 10, and 17 favours the diagnosis of chRCC over RO. FISH analysis is shown to be a useful tool that helps identify differences between these 2 tumour types.	66
Endogenous avidin-binding activity (EABA)	Renal TMAs (30 RO, 18 chRCC, 64 ccRCC, 50 pRCC, 31 benign renal tissues)	97% RO, 26% ccRCC, 35% pRCC with granular/eosinophilic (GE) features and 6% of chRCCs positive for EABA. RCC without GE features were negative. EABA is an excellent marker for RO, and so useful in differentiating RO from chRCC.	60
PAX8 and MUC-1	TMAs of 36 chRCC, 20 RO	Expression of PAX8 more frequent in RO than in chRCC (55% vs 25%). MUC1 expressed more diffusely and frequently in chRCC than RO (94% vs 55%).	48

AM, angiomyolipomas; BCA2, a RING H2 finger protein of RING E3 ligases; ccRCC, clear cell renal cell carcinoma; cdRCC, collecting duct renal cell carcinoma; chRCC, chromophobe renal cell carcinoma; NB, nephroblastomas; pRCC, papillary renal cell carcinoma; RO, renal oncocytoma; UC, urothelial carcinomas.

papillary or conventional RCCs.<sup>34</sup> On MRI, chRCCs typically have heterogeneous T2 signal intensity and enhancement.

By contrast, on CT scanning, ROs typically show a well-defined, smooth, relatively homogeneous solid mass with a central area of hypoattenuation due to the presence of a central stellate scar, and rarely show any extension to the renal vein, inferior vena cava or the adrenals. MRI scan will typically reveal low to moderate homogeneous intensity on T1-weighted images and relatively high signal intensity on T2-weighted images.<sup>35</sup> Classically, if renal angiography on ROs were performed, it would show a typical spoke-wheel pattern, highlighting the marked peripheral vascularity in contrast with the relatively hypovascular central part of the tumour. However, classical

hypoattenuation of the central stellate scar on CT scan is seen in less than one-third of ROs, and although characteristic of ROs, it is not diagnostic.<sup>14–20</sup> Moreover, there are no consistently reliable pathognomic CT scan features that can safely differentiate ROs from RCCs.<sup>36</sup> Therefore, most ROs are treated as suspicious of RCCs based on imaging, and thereafter, are subjected to surgical resection.

A recent study on the ability of MRI to discriminate ROs from chRCCs showed that these two entities exhibited similar findings, and no MRI features were reliable in distinguishing between the two.<sup>37</sup> The ability of any renal lesion to uptake the 18-fluorodeoxyglucose (FDG) is the basis of 18-FDG PET/CT scans. However, in detection of renal tumours, the role of FDG

PET is limited as there are high false negative rates<sup>38</sup> Benign ROs are also often FDG-avid, and thus, this cannot be used in separating them from malignant renal tumours.<sup>39</sup> Recently, multiphasic multidetector CT scans have helped to discriminate ccRCCs from ROs, papillary RCCs and chRCCs, by using the different enhancements at various phases of the scans.<sup>40</sup> This will aid somewhat in the distinction of ccRCCs from ROs, but not the discrimination of ROs from chRCCs. Arterial phase enhancements >500% and washout values >50% in Hounsfield units obtained in multiphasic CT scans can be seen exclusively in ROs and can aid in distinguishing ROs from other subtypes of RCCs.<sup>41</sup>

## PATHOLOGY

Despite the non-invasive discriminatory features of multiphasic CT scans, renal mass biopsy provides the best opportunity for preoperative diagnosis. However, there are numerous potential shortcomings for this procedure, leading to the inevitability of surgical excision. One of the main drawbacks of renal mass biopsy is the relative difficulty faced by pathologists to accurately and conclusively diagnose renal tumour subtype from the limited tissue biopsy samples, as usually an entire range of cyto-architectural features is necessary for examination to arrive at a diagnosis.<sup>42</sup> However, as a general rule, if the lesion looks like chRCCs on needle biopsy, it can be confidently reported as such. By comparison, a lesion that looks like an RO may be incompletely sampled, with other areas merging into the eosinophilic variant of chRCCs. This may be a hybrid tumour or simply oncocyoma-like areas in a chRCC. Therefore, most pathologists would not diagnose an RO outright on a needle biopsy, and make a comment as to the possibility of having chRCCs elsewhere in the tumour. In addition to the difficulties in differentiating ROs from chRCCs clinically, the pathological features following surgical resection of these tumours often overlap and pose a diagnostic challenge to pathologists.

ChRCCs are well-circumscribed encapsulated tumours which have a light-brown to tan cut surface. These are typically solid but cystic areas can be found. Central scarring may be seen. Histologically, there are two types. The classic type has large polygonal cells with finely granular cytoplasm. These have prominent plant-like thick cell membranes. The eosinophilic variant is composed of polygonal cells with abundant eosinophilic cytoplasm. Nuclei are irregular, crinkled and angulated,

often with perinuclear clearing. Binucleation is common. A solid sheet-like pattern with poor cellular cohesion is commonly found. ROs are also well circumscribed, but unencapsulated tumours which are typically mahogany brown but sometimes tan coloured. A central stellate scar is present in about one-third of cases. Rarely, cystic change or haemorrhage can be found. Histologically, there are large round polygonal cells with abundant eosinophilic cytoplasm and round nuclei. Nucleoli are inconspicuous. Cells form nests, tubules, acini and microcysts. Focal degenerative nuclear atypia may be seen. Figure 1 demonstrates histopathology of chRCCs and ROs.

Hybrid tumours have zones classic for ROs and chRCCs as described above. Some cases of chRCCs have features overlapping with ccRCCs in which the component cells have granular cytoplasm. In ccRCCs, at least some areas have cells with completely clear cytoplasm with high vascularity and typical, delicate, thin-walled, vascular structure throughout the tumour, contrasting with thick-walled blood vessels present in chRCCs. Also, a ccRCC lacks the plant-like architecture seen in chRCCs.

Table 1 describes the macroscopic and microscopic features of ROs and chRCCs. Despite having some subtle distinguishing macroscopic, microscopic and ultrastructural differences, there is often need to use ancillary histochemical and immunohistochemical (IHC) stains to differentiate these two entities. Recently, a new oncocytic variant of a chRCC was described, that morphologically resembles RO, but has the biological characteristics of a chRCC.<sup>43</sup> This further adds to the difficulties for pathologists to discern ROs from all these variants of chRCCs.

To date, none of the histochemical, IHC or cytogenetic features has been proven to be reliable and specific.<sup>44</sup> However, IHC markers may be a cost-effective and valuable form of information for monitoring disease for both prognosis and treatment-planning regimens. Tables 2–4 list some of the histochemical and IHC markers that have been published. Hale's colloidal iron staining is still used. Currently, the most useful IHC markers for the differentiation of renal tumours are vimentin, cytokeratin (CK)7, CD10 and marker for RCC (RCCma). Vimentin has been shown to be positive in ccRCCs and negative in chRCCs and ROs, and CK7 is positive in chRCCs and negative in ROs and ccRCCs. RCCma and CD10 are positive in ccRCCs and negative in both chRCCs and ROs. Hale's colloidal iron staining with diffuse reticular pattern and perinuclear halo is present in chRCCs but non-existent in ROs and ccRCCs.<sup>45</sup>

**Table 4** Biomarkers from the cadherin family (also known as calcium-dependent adhesion)

Method	Number of patients	Significance of success as biomarker	Ref
Kidney-specific cadherin (Ksp-cad)	102 ccRCC, 46 pRCC, 30 chRCC, 3 cdRCC, 31 RO	Ksp-cad was expressed almost exclusively in chRCCs (97.7% of cases). Ksp-cad offers a quick, dependable approach for differentiating between RO and chRCCs.	44
	42 ccRCC, 30 pRCC, 13 chRCC, 20 RO using whole sections	By contrast with Mazal <i>et al</i> , 2004, here both chRCC (13/13) and RO (19/20) were positive for Ksp-cad. Ksp-cad not a useful marker for differentiating.	67
	15 chRCC, 15 RO for mRNA analysis and IHC on TMAs containing 36 chRCC, 41 RO	Ksp-cad differentiate RO from chRCC. Ksp-cad was present in chRCCs and ROs at mRNA (89% chRCC and 64% RO) and IHC (31/36 chRCCs and 31/41 RO)	68
N-Cadherin	21 Japanese cases chRCC, ccRCC, RO.	chRCC and RO were positive for E-cadherin but not for N-cadherin. All ccRCCs were negative for E-cadherin, and 58% were positive for N-cadherin. Useful to distinguish chRCC from ccRCC but not between chRCC and RO.	69
E-Cadherin			
Ep-CAM (epithelial cell adhesion molecule)	22 chRCC, 17 RO, 45 ccRCC	Expressed in all chRCC in more than 90% of cells. EpCAM-positive RO (5/17; 29%) had single cell or small cell cluster positivity. The homogeneous EpCAM expression assists to diagnosis chRCC from RO.	47
	10 each of ccRCC, pRCC, chRCCs, RO	EpCAM distinguished between RO and chRCC. RO were negative for EpCAM but positive in 8/10 (80%) of chRCC.	64

ccRCC, clear cell renal cell carcinoma; pRCC, papillary renal cell carcinoma; chRCC, chromophobe renal cell carcinoma; RO, renal oncocyoma; cdRCC, collecting duct renal cell carcinoma.

**Table 5** Biomarkers from the cytokeratin family

Method	Number of patients	Significance of success as biomarker	Reference
CK7 (Basic or neutral cytokeratin)	6 chRCC, 11 RO	All chRCC, strong cytoplasmic staining with peripheral cell accentuation. 8/11 RO, negative, 3 weakly staining.	70
	21 chRCC, 26 RO	chRCCs (100%) and almost all RO (96%) were positive for CK7.	71
	11 chRCC, 21 RO from 4 hospitals	73% chRCC, 25% RO positive for CK7; 33% RO focally positive for CK7. No consistency in differentiating the 2 neoplasms.	72
	22 chRCC, 17 RO, 45 ccRCC	Positive in 100% chRCC, 8% ccRCC and negative in RO.	45
	TMA (20 each ccRCC, chRCC, pRCC, RO)	Positive in 80% chRCC, 0% RO.	47
	TMA (36 chRCC, 20 RO)	Positive in pRCC (90%), chRCC (89%), and RO (90%). Expressed significantly more often in chRCC than RO, both diffusely (53% vs 10%) and focally (42% vs 15%).	48
CK8 (Basic or neutral cytokeratins)	TMA (30 RO, 18 chRCC, 64 ccRCC, 50 pRCC)	81% pRCC, 63% chRCC, essentially negative in ccRCC and RO	60
	10 each ccRCC, pRCC, chRCC, RO	Distinguished RO and chRCC. RO were not stained 80% chRCCs were positive.	64
CK18 (Acidic cytokeratin)	76 cases (30 ccRCC, 16 pRCC, 21 chRCC, 8 RO, 1 cdRCC)	Positive in 70% ccRCC, 93% chRCC and 87.5% RO.	45
CK19 (Acidic cytokeratin)	76 cases (30 ccRCC, 16 pRCC, 21 chRCC, 8 RO, 1 cdRCC)	Positive in 87% ccRCC, 100% chRCC and 87.5% RO.	45
CK20 (Acidic cytokeratin)	76 cases (30 ccRCC, 16 pRCC, 21 chRCC, 8 RO, 1 cdRCC)	Positive in 41% ccRCC, 37.5% chRCC and 62.5% RO. Not a useful marker for differentiation among these subtypes.	45
	15 RO only from archives	12/15 RO were positive for CK20	73
	11 chRCC, 21 RO from 4 hospitals	chRCC and RO were uniformly negative for CK20	72
	76 cases (30 ccRCC, 16 pRCC, 21 chRCC, 8 RO, 1 cdRCC)	Positive in only 8% ccRCCs, 12.5% chRCCs, negative in RO. Not a useful marker for differentiation among these subtypes.	45

ccRCC, clear cell renal cell carcinoma; cdRCC, collecting duct renal cell carcinoma; chRCC, chromophobe renal cell carcinoma; pRCC, papillary renal cell carcinoma; RO, renal oncocytoma; TMA, tissue microarray.

Colloidal iron and widespread CK7 positivity have been suggested to be useful in distinguishing chRCCs from ROs. In ROs, colloidal iron staining is usually negative and CK7 shows only focal positivity. However, there is overlap in the staining patterns, preventing these stains to be of much practical value. Negative staining for vimentin and widespread staining for CK7 versus negative staining for CK7 and positive staining for vimentin can be useful in distinguishing chRCCs from ccRCCs.

However, as seen in table 2, these IHC markers still have their pitfalls in distinguishing between chRCCs and ROs. For example, the problems with Hale's colloidal iron in certain instances is its failure to stain adequately, or the staining pattern (diffuse cytoplasmic vs luminal) could not be adequately assessed.<sup>46</sup> However, vimentin may be useful in discriminating chRCCs from other RCCs, and a panel of vimentin with GST- $\alpha$  and EpCAM may achieve 100% sensitivity and specificity for the differential diagnosis of chRCCs, ROs and ccRCCs.<sup>47</sup>

ROs and chRCCs share histologic and cytologic features, and also share IHC markers for S100A1 and CD117 (KIT).<sup>48</sup> Several other studies with IHC markers, including kidney-specific cadherin, CK7, EMA, CD10, RCC, c-KIT, and RON proto-oncogene have been used to distinguish chRCCs from ROs, but the results of these studies are inconsistent and unsatisfactory.<sup>49</sup> Table 3 illustrates current biomarkers used to differentiate chRCCs from ROs, directly or indirectly. BCA2, a RING H2 finger protein RING E3 ligase, holds potential as a tool to distinguish ROs from its mimickers, like chRCCs.<sup>50</sup> Additionally, RO has significantly higher expression of the cancer-testis antigens (CTAs), such as MAGE-A3/4 and NY-ESO-1.<sup>51</sup> Further investigation is needed to evaluate the potential diagnostic implications for these markers.

The cadherins comprise of a family of transmembrane glycoproteins that function as calcium-dependent homotypic adhesion molecules and are expressed by the majority of epithelium. Currently, over 20 different tissue-specific cadherins have been

identified.<sup>52</sup> The promise of cadherin proteins in distinguishing chRCCs from ROs is shown in table 4. CKs are a family of intermediate filaments that are characteristic markers of epithelial differentiation. Currently, 20 distinct CKs have been identified. They can be useful in the differential diagnosis of neoplasms of epithelial origin, and consequently, several CKs have been investigated in renal neoplasms.<sup>53</sup> The CKs that have been trialled to discriminate chRCCs from other RCCs and also ROs, are listed in table 5, but none holds major promise, including CK7. Caveolin-1 is a scaffolding protein encoded by the Cav-1 protein. This has demonstrated better promise in differentiating chRCCs from ROs than CK7.<sup>47</sup>

## CONCLUSION

The standard treatment of localised renal tumours remains surgical resection via complete or partial nephrectomy. The increasing detection of small renal masses with a significant chance of benign aetiology provides a diagnostic and management challenge. ROs, and to a lesser extent small chRCCs, are two lesions that could be managed conservatively in many situations, avoiding the morbidity inherent to resection of renal lesions. However, a very high level of diagnostic certainty is required if surgical intervention is to be avoided. Current imaging and biopsy techniques do not always provide this certainty as evidenced by the number of benign small renal lesions reported in contemporary surgical series. If confident diagnosis of renal lesions with low or no malignant potential can be achieved, then active surveillance will usually be appropriate, with intervention reserved for tumours demonstrating excessive growth or symptoms. The ability to diagnose ROs and chRCCs with a high level of confidence may lead to improved use of preoperative diagnostic techniques and reduced intervention rates for indolent renal lesions. Achieving consistent accurate diagnosis of ROs and chRCCs via non-surgical means will remain elusive until modern molecular biomarkers are identified.

## Key messages

- ▶ Clinical diagnostic dilemma and difficult histopathological differentiation of renal oncocytoma from chromophobe renal cell carcinoma still persist.
- ▶ The ability to achieve confident accurate diagnosis of these renal tumors via non-surgical means remain elusive until new specific molecular biomarkers are discovered.
- ▶ Better preoperative non-invasive characterisation of specific biomarkers for renal oncocytoma and chromophobe renal cell carcinoma may lead to reduced rates of surgical intervention for benign renal lesions.

**Contributors** KLN, RR and YNY performed the literature search. KLN, RR, YNY, HS and CM drafted the manuscript, while GCG and STW helped in the critical revision of the manuscript. HS provided the picture of histology slides. All authors contributed actively in this manuscript, and all approved the final revised version of this manuscript.

**Funding** This review article was partly supported by the University Malaya Research Grant (UMRG RG348/11HTM).

**Competing interests** None.

**Provenance and peer review** Not commissioned; externally peer reviewed.

## REFERENCES

- 1 Ljungberg B, Campbell SC, Choi HY, *et al.* The epidemiology of renal cell carcinoma. *Eur Urol* 2011;60:615–21.
- 2 Duchene DA, Lotan Y, Cadeddu JA, *et al.* Histopathology of surgically managed renal tumors: analysis of a contemporary series. *Urology* 2003;62:827–30.
- 3 Gupta K, Miller JD, Li JZ, *et al.* Epidemiologic and socioeconomic burden of metastatic renal cell carcinoma (mRCC): a literature review. *Cancer Treat Rev* 2008;34:193–205.
- 4 Schachter LR, Cookson MS, Chang SS, *et al.* Second prize: frequency of benign renal cortical tumors and histologic subtypes based on size in a contemporary series: what to tell our patients. *J Endourol* 2007;21:819–23.
- 5 Yusenko MV. Molecular pathology of renal oncocytoma: A review. *Int J Urol* 2010;17:592–12.
- 6 Oxley JD, Sullivan J, Mitchelmore A, *et al.* Metastatic renal oncocytoma. *J Clin Pathol* 2007;60:720–2.
- 7 Delongchamps NB, Galmiche L, Eiss D, *et al.* Hybrid tumour 'oncocytoma-chromophobe renal cell carcinoma' of the kidney: a report of seven sporadic cases. *BJU Int* 2009;103:1381–4.
- 8 Zippel . Zur Kenntnis der Oncocytome. *Virchows Arch A Pathol Anat* 1942;308:360–82.
- 9 Klein MJ, Valensi QJ. Proximal tubular adenomas of kidney with so-called oncocytic features. A clinicopathologic study of 13 cases of a rarely reported neoplasm. *Cancer* 1976;38:906–14.
- 10 Storkel S, Pannen B, Thoenes W, *et al.* Intercalated cells as a probable source for the development of renal oncocytoma. *Virchows Arch B Cell Pathol Incl Mol Pathol* 1988;56:185–9.
- 11 Thoenes W, Storkel S, Rumpelt HJ. Human chromophobe cell renal carcinoma. *Virchows Arch B Cell Pathol Incl Mol Pathol* 1985;48:207–17.
- 12 Thoenes W. SSaRHJ. Histopathology and classification of renal cell tumours (adenomas, oncocytomas, carcinomas). The basic cytological and histopathological elements and their use for diagnostics. *Pathol Res Pract* 1986;181:125–43.
- 13 Amin R, Anthony P. Metastatic renal oncocytoma: A case report and review of the literature. *Clin Oncol (R Coll Radiol)* 1999;11:277–9.
- 14 Chawla SN, Crispen PL, Hanlon AL, *et al.* The natural history of observed enhancing renal masses: meta-analysis and review of the world literature. *J Urol* 2006;175:425–31.
- 15 Kawaguchi S, Fernandes KA, Finelli A, *et al.* Most renal oncocytomas appear to grow: observations of tumor kinetics with active surveillance. *J Urol* 2011;186:1218–22.
- 16 Kurup AN, Thompson RH, Leibovich BC, *et al.* Renal oncocytoma growth rates before intervention. *BJU Int* 2012;110:1444–8.
- 17 Cindolo L, de ITA, Schips L, *et al.* Chromophobe renal cell carcinoma: comprehensive analysis of 104 cases from multicenter European database. *Urology* 2005;65:681–6.
- 18 Vera-Badillo FE, Conde E, Duran I. Chromophobe renal cell carcinoma: a review of an uncommon entity. *Int J Urol* 2012;19:894–900.
- 19 Lopez-Beltran A, Scarpelli M, Montironi R, *et al.* 2004 WHO classification of the renal tumors of the adults. *Eur Urol* 2006;49:798–805.
- 20 Khoo SK, Bradley M, Wong FK, *et al.* Birt-Hogg-Dube syndrome: mapping of a novel hereditary neoplasia gene to chromosome 17p12-q11.2. *Oncogene* 2001;20:5239–42.
- 21 Nickerson ML, Warren MB, Toro JR, *et al.* Mutations in a novel gene lead to kidney tumors, lung wall defects, and benign tumors of the hair follicle in patients with the Birt-Hogg-Dube syndrome. *Cancer Cell* 2002;2:157–64.
- 22 Warfel KA, Eble JN. Renal oncocytomatosis. *J Urol* 1982;127:1179–80.
- 23 Adamy A, Lowrance WT, Yee DS, *et al.* Renal oncocytosis: management and clinical outcomes. *J Urol* 2011;185:795–801.
- 24 Volpe A, Novara G, Antonelli A, *et al.* Chromophobe renal cell carcinoma (RCC): oncological outcomes and prognostic factors in a large multicentre series. *BJU Int* 2012;110:76–83.
- 25 Tobias Klatt K-rH, Jonathan W. Said, Malte Bohm. Pathobiology and prognosis of chromophobe renal cell carcinoma. *Urol Oncol-Semin Orig Investig* 2008;26:604–09.
- 26 PerezOrdonez B, Hamed G, Campbell S, *et al.* Renal oncocytoma: A clinicopathologic study of 70 cases. *Am J Surg Pathol* 1997;21:871–83.
- 27 Akbulut S, Senol A, Cakabay B, *et al.* Giant renal oncocytoma: a case report and review of the literature. *J Med Case Rep* 2010;4:52.
- 28 Romis L, Cindolo L, Patard JJ, *et al.* Frequency, clinical presentation and evolution of renal oncocytomas: multicentric experience from a European database. *Eur Urol* 2004;45:53–7; discussion 57
- 29 Dechet CB, Bostwick DG, Blute MJ, *et al.* Renal oncocytoma: Multifocality, bilateralism, metachronous tumor development and coexistent renal cell carcinoma. *J Urology* 1999;162:40–2.
- 30 Trpkov K, Yilmaz A, Uzer D, *et al.* Renal oncocytoma revisited: a clinicopathological study of 109 cases with emphasis on problematic diagnostic features. *Histopathology* 2010;57:893–906.
- 31 Davis CJ MF, Sesterhenn IA, Ho CK. Renal Oncocytoma. Clinicopathological study of 166 patients. *J Urogenital Pathol* 1991;1:41–52.
- 32 Cheville JC, Lohse CM, Zincke H, *et al.* Comparisons of outcome and prognostic features among histologic subtypes of renal cell carcinoma. *Am J Surg Pathol* 2003;27:612–24.
- 33 Yusenko MV. Molecular pathology of chromophobe renal cell carcinoma: a review. *Int J Urol* 2010;17:592–600.
- 34 Kim JK, Kim TK, Ahn HJ, *et al.* Differentiation of subtypes of renal cell carcinoma on helical CT scans. *AJR Am J Roentgenol* 2002;178:1499–506.
- 35 Remark RR, Berquist TH, Lieber MM, *et al.* Magnetic resonance imaging of renal oncocytoma. *Urology* 1988;31:176–9.
- 36 Choudhary S, Rajesh A, Mayer NJ, *et al.* Renal oncocytoma: CT features cannot reliably distinguish oncocytoma from other renal neoplasms. *Clin Radiol* 2009;64:517–22.
- 37 Rosenkrantz AB, Hindman N, Fitzgerald EF, *et al.* MRI features of renal oncocytoma and chromophobe renal cell carcinoma. *AJR Am J Roentgenol* 2010;195:W421–7.
- 38 Aide N, Cappelletti O, Bottet P, *et al.* Efficiency of [(18)F]FDG PET in characterising renal cancer and detecting distant metastases: a comparison with CT. *Eur J Nucl Med Mol Imaging* 2003;30:1236–45.
- 39 Ramdave S, Thomas GW, Berlangieri SU, *et al.* Clinical role of F-18 fluorodeoxyglucose positron emission tomography for detection and management of renal cell carcinoma. *J Urol* 2001;166:825–30.
- 40 Young JR, Margolis D, Sauk S, *et al.* Clear Cell Renal Cell Carcinoma: Discrimination from Other Renal Cell Carcinoma Subtypes and Oncocytoma at Multiphasic Multidetector CT. *Radiology* 2013;267:444–53.
- 41 Bird VG, Kanagarajah P, Morillo G, *et al.* Response by authors re: Differentiation of oncocytoma and renal cell carcinoma in small renal masses (<4 cm): the role of 4-phase computerized tomography. *World J Urol* 2013;31:1011–2.
- 42 Barocas DA, Rohan SM, Kao J, *et al.* Diagnosis of renal tumors on needle biopsy specimens by histological and molecular analysis. *J Urol* 2006;176:1957–62.
- 43 Kuroda N, Tanaka A, Yamaguchi T, *et al.* Chromophobe renal cell carcinoma, oncocytic variant: a proposal of a new variant giving a critical diagnostic pitfall in diagnosing renal oncocytic tumors. *Med Mol Morphol* 2013;46:49–55.
- 44 Mazal PR, Exner M, Haitel A, *et al.* Expression of kidney-specific cadherin distinguishes chromophobe renal cell carcinoma from renal oncocytoma. *Human pathology* 2005;36:22–8.
- 45 Geramizadeh B, Ravanshad M, Rahsaz M. Useful markers for differential diagnosis of oncocytoma, chromophobe renal cell carcinoma and conventional renal cell carcinoma. *Indian J Pathol Microbiol* 2008;51:167–71.
- 46 Latham B, Dickensin GR, Oliva E. Subtypes of chromophobe cell renal carcinoma: an ultrastructural and histochemical study of 13 cases. *Am J Surg Pathol* 1999;23:530–5.
- 47 Liu L, Qian J, Singh H, *et al.* Immunohistochemical analysis of chromophobe renal cell carcinoma, renal oncocytoma, and clear cell carcinoma: an optimal and practical panel for differential diagnosis. *Arch Pathol Lab Med* 2007;131:1290–7.
- 48 Bing Z, Lal P, Lu S, *et al.* Role of carbonic anhydrase IX, alpha-methylacyl coenzyme a racemase, cytokeratin 7, and galectin-3 in the evaluation of renal neoplasms: a tissue microarray immunohistochemical study. *Ann Diagn Pathol* 2013;17:58–62.

- 49 Lee HW, Lee EH, Lee CH, *et al.* Diagnostic Utility of Caveolin-1 and MOC-31 in Distinguishing Chromophobe Renal Cell Carcinoma from Renal Oncocytoma. *Korean J Urol* 2011;52:96–103.
- 50 Ehsani L, Seth R, Bacopulos S, *et al.* BCA2 is differentially expressed in renal oncocytoma: an analysis of 158 renal neoplasms. *Tumour Biol* 2013;34:787–91.
- 51 Demirovic A, Dzombeta T, Tomas D, *et al.* Immunohistochemical expression of tumor antigens MAGE-A3/4 and NY-ESO-1 in renal oncocytoma and chromophobe renal cell carcinoma. *Pathol Res Pract* 2010;206:695–9.
- 52 Langner C, Ratschek M, Rehak P, *et al.* Expression of MUC1 (EMA) and E-cadherin in renal cell carcinoma: a systematic immunohistochemical analysis of 188 cases. *Mod Pathol* 2004;17:180–8.
- 53 Skinnider BF, Folpe AL, Hennigar RA, *et al.* Distribution of cytokeratins and vimentin in adult renal neoplasms and normal renal tissue: potential utility of a cytokeratin antibody panel in the differential diagnosis of renal tumors. *Am J Surg Pathol* 2005;29:747–54.
- 54 Gudbjartsson T, Hardarson S, Petursdottir V, *et al.* Renal oncocytoma: a clinicopathological analysis of 45 consecutive cases. *BJU Int* 2005;96:1275–9.
- 55 Tickoo SK, Amin MB. Discriminant nuclear features of renal oncocytoma and chromophobe renal cell carcinoma. Analysis of their potential utility in the differential diagnosis. *Am J Clin Pathol* 1998;110:782–7.
- 56 Abrahams NA, MacLennan GT, Khoury JD, *et al.* Chromophobe renal cell carcinoma: a comparative study of histological, immunohistochemical and ultrastructural features using high throughput tissue microarray. *Histopathology* 2004;45:593–602.
- 57 Crotty TB, Farrow GM, Lieber MM. Chromophobe cell renal carcinoma: clinicopathological features of 50 cases. *J Urol* 1995;154:964–7.
- 58 Wang HY, Mills SE. KIT and RCC are useful in distinguishing chromophobe renal cell carcinoma from the granular variant of clear cell renal cell carcinoma. *Am J Surg Pathol* 2005;29:640–6.
- 59 Tickoo SK, Amin MB, Zarbo RJ. Colloidal iron staining in renal epithelial neoplasms, including chromophobe renal cell carcinoma: emphasis on technique and patterns of staining. *Am J Surg Pathol* 1998;22:419–24.
- 60 Huang W, Kanehira K, Drew S, *et al.* Oncocytoma can be differentiated from its renal cell carcinoma mimics by a panel of markers: an automated tissue microarray study. *Appl Immunohistochem Mol Morphol* 2009;17:12–7.
- 61 Pan CC, Chen PC, Ho DM. The diagnostic utility of MOC31, BerEP4, RCC marker and CD10 in the classification of renal cell carcinoma and renal oncocytoma: an immunohistochemical analysis of 328 cases. *Histopathology* 2004;45:452–9.
- 62 Huo L, Sugimura J, Tretiakova MS, *et al.* C-kit expression in renal oncocytomas and chromophobe renal cell carcinomas. *Hum Pathol* 2005;36:262–8.
- 63 Pan CC, Chen PC, Chiang H. Overexpression of KIT (CD117) in chromophobe renal cell carcinoma and renal oncocytoma. *Am J Clin Pathol* 2004;121:878–83.
- 64 Ray ER, Goodwill J, Chandra A, *et al.* Exploring the Potential of Immunohistochemistry to Identify Renal Oncocytoma. *Br J Med Surg Urol* 2011;4:8–12.
- 65 Patton KT, Tretiakova MS, Yao JL, *et al.* Expression of RON Proto-oncogene in Renal Oncocytoma and Chromophobe Renal Cell Carcinoma. *Am J Surg Pathol* 2004;28:1045–50.
- 66 Brunelli M, Delahunt B, Gobbo S, *et al.* Diagnostic usefulness of fluorescent cytogenetics in differentiating chromophobe renal cell carcinoma from renal oncocytoma: a validation study combining metaphase and interphase analyses. *Am J Clin Pathol* 2010;133:116–26.
- 67 Shen SS, Krishna B, Chirala R, *et al.* Kidney-specific cadherin, a specific marker for the distal portion of the nephron and related renal neoplasms. *Mod Pathol* 2005;18:933–40.
- 68 Adley BP, Gupta A, Lin F, *et al.* Expression of kidney-specific cadherin in chromophobe renal cell carcinoma and renal oncocytoma. *Am J Clin Pathol* 2006;126:79–85.
- 69 Taki A, Nakatani Y, Misugi K, *et al.* Chromophobe renal cell carcinoma: an immunohistochemical study of 21 Japanese cases. *Mod Pathol* 1999;12:310–7.
- 70 Leroy X, Moukassa D, Copin MC, *et al.* Utility of cytokeratin 7 for distinguishing chromophobe renal cell carcinoma from renal oncocytoma. *Eur Urol* 2000;37:484–7.
- 71 Garcia E, Li M. Caveolin-1 immunohistochemical analysis in differentiating chromophobe renal cell carcinoma from renal oncocytoma. *Am J Clin Pathol* 2006;125:392–8.
- 72 Wu SL, Kothari P, Wheeler TM, *et al.* Cytokeratins 7 and 20 immunoreactivity in chromophobe renal cell carcinomas and renal oncocytomas. *Mod Pathol* 2002;15:712–7.
- 73 Stopyra GA, Warhol MJ, Mulhaupt HA. Cytokeratin 20 immunoreactivity in renal oncocytomas. *J Histochem Cytochem* 2001;49:919–20.