

## Correction

Quesada AE, Tholpady A, Wanger A, *et al.* Utility of bone marrow examination for workup of fever of unknown origin in patients with HIV/AIDS. *J Clin Pathol* 2015;68:241–5. doi:10.1136/jclinpath-2014-202715

The following sentence has been amended to read: The average white blood cell count for all patients was  $\times 10^9/L$  (range 0.1–23.7  $\times 10^9/L$ ). The average haemoglobin was 91 g/L (range 54–135 g/L). The average platelet count was 146  $\times 10^9/L$  (range 4–642  $\times 10^9/L$ ).

Table 2 has also been amended to read:

	BM culture negative	BM culture positive
White blood cell count	4 $\times 10^9/L$	2.8 $\times 10^9/L$
Reference range: 4.5–11.0 $\times 10^9/L$	0.1–23.7 $\times 10^9/L$	1.2–5.6 $\times 10^9/L$
Haemoglobin	93 g/L	83 g/L
Reference range: 54–135 g/L	57–135 g/L	54–117 g/L
Platelets	148 $\times 10^9/L$	140 $\times 10^9/L$
Reference range: 150–400 $\times 10^9/L$	4–642 $\times 10^9/L$	12–390 $\times 10^9/L$



CrossMark

*J Clin Pathol* 2015;68:487. doi:10.1136/jclinpath-2014-202715corr1

## Correction

---

Quesada AE, Tholpady A, Wanger A, *et al.* Utility of bone marrow examination for workup of fever of unknown origin in patients with HIV/AIDS. *J Clin Pathol* 2015;68:241–5. doi:10.1136/jclinpath-2014-202715

The following sentence has been amended to read: The average white blood cell count for all patients was  $3.8 \times 10^9/L$  (range  $0.1\text{--}23.7 \times 10^9/L$ ). The average haemoglobin was 91 g/L (range 54–135 g/L). The average platelet count was  $146 \times 10^9/L$  (range  $4\text{--}642 \times 10^9/L$ ).



CrossMark

*J Clin Pathol* 2015;68:577. doi:10.1136/jclinpath-2014-202715corr2