Genetics and genomics of breast fibroadenomas

Benjamin Nathanael Loke,^{1,2} Nur Diyana Md Nasir,² Aye Aye Thike,^{2,3}
Jonathan Yu Han Lee,⁴ Cheek Soon Lee,^{5,6,7,8,9} Bin Tean Teh,^{3,10,11,12}
Puay Hoon Tan^{2,3,13}

¹Department of Biological Sciences, National University of Singapore, Singapore

²Department of Anatomical Pathology, Singapore General Hospital, Singapore

³Duke–NUS Medical School, Singapore

⁴School of Science and Health, Western Sydney University, Sydney, New South Wales, Australia

⁵Discipline of Pathology, School of Medicine, Western Sydney University, Sydney, New South Wales, Australia

⁶Department of Anatomical Pathology, Liverpool Hospital, Liverpool, New South Wales, Australia

⁷South Western Sydney Clinical School, University of New South Wales, Sydney, New South Wales, Australia

⁸Department of Tissue Pathology, Royal Prince Alfred Hospital, Sydney, New South Wales, Australia

⁹Cancer Pathology Laboratory, Bosch Institute, University of Sydney, Sydney, New South Wales, Australia

¹⁰Laboratory of Cancer Epigenome, National Cancer Centre Singapore, Singapore

¹¹Cancer Science Institute of Singapore, National University of Singapore, Singapore

¹²Institute of Molecular and Cell Biology, Singapore

¹³Division of Pathology, Singapore General Hospital, Singapore

Correspondence to

Dr Puay Hoon Tan, Division of Pathology, Singapore General Hospital, Singapore; tan.puay.hoon@singhealth.com.sg

Received 12 October 2017

Revised 16 November 2017

Accepted 17 November 2017

Published Online First

16 December 2017

To cite: Loke BN, Md Nasir ND, Thike AA, et al. *J Clin Pathol* 2018;**71**:381–387.

ABSTRACT

Fibroadenomas of the breast are benign fibroepithelial tumours most frequently encountered in women of reproductive age, although they may be diagnosed at any age. The fibroadenoma comprises a proliferation of both stromal and epithelial components. The mechanisms underlying fibroadenoma pathogenesis remain incompletely understood. In the clinical setting, distinguishing cellular fibroadenomas from benign phyllodes tumours is a common diagnostic challenge due to subjective histopathological criteria and interobserver differences. Recent sequencing studies have demonstrated the presence of highly recurrent mutations in fibroadenomas, and also delineated the genomic landscapes of fibroadenomas and the closely related phyllodes tumours, revealing differences at the gene level, which may be of potential adjunctive diagnostic use. The present article provides an overview of key studies uncovering genetic and genomic abnormalities in fibroadenomas, from initial karyotype reports revealing myriad cytogenetic aberrations to next-generation sequencing-based approaches that led to the discovery of highly recurrent *MED12* mutations. A thorough understanding of these abnormalities is important to further elucidate the mechanisms by which fibroadenomas arise and to refine diagnostic assessment of this very common tumour.

INTRODUCTION

Breast fibroadenomas (FAs) are common, benign fibroepithelial lesions composed of stromal and epithelial components, with intralobular stroma enclosing glandular spaces lined by luminal epithelium and myoepithelium. This tumour is most frequently encountered in adolescent girls and young women within the first three decades of life, although it may also be diagnosed at any age.^{1–3} FAs have been clinically observed to be hormone dependent—stimulated by oestrogen, progesterone and lactation during pregnancy, as well as shrinking during menopause.^{4–6} Typically, they present as slow-growing, firm and solitary nodules, although synchronous and metachronous multifocal lesions have also been described in the literature.⁷ While FAs are benign neoplasms, one study suggested that they are an independent risk factor for breast cancer,⁸ while another report provided evidence that they are associated with a twofold increase in relative risk of developing invasive breast carcinoma after 20 years.⁹ Although the genetic risk factors for FAs are currently unknown, patients with Carney complex may develop bilateral or multiple myxoid FAs, and bilateral and/or multiple FAs have been

significantly associated with family history.^{10 11} In renal transplant patients, the use of cyclosporin has been linked to the development of bilateral and/or multiple FAs.^{12–14} In this review, we appraise the literature on the genetics and genomics of breast FAs.

CYTOGENETIC ABERRATIONS

In the early 1990s, structural abnormalities including rearrangements of chromosomes 1, 6, 7, 8, 12 and 15 were described in isolated case reports of breast FAs, together with monosomy of chromosomes X, 12 and 21 and trisomy of chromosomes 5, 6, 11, 17 and 20.^{15–20} Of particular interest was the 12q13–15 region, which was also implicated in several other benign tumours in different anatomical sites such as lipomas, uterine leiomyomas and pleomorphic adenomas.^{21–23} Non-random involvement of alleles found in chromosomes 1 and 17 were previously reported in breast cancer,^{24–31} and it was thought that there could be overlapping mutations driving the development of FAs as well.³² However, no consensus could be achieved regarding specific mutations that result in the development of FAs.

This changed in 1995 when Schoenmakers *et al* identified a candidate gene named HMGI-C within the multiple aberration region on chromosome 12q15 that frequently harbours breakpoints in benign solid tumours, which was mutated in FA together with seven other tumour types.³³ In a later clinical case report of an intracanalicular FA showing a clonal chromosomal aberration t(4;12), Staats *et al* used fluorescence in situ hybridisation (FISH) analysis for the chromosomal region 12q14–15 and narrowed down the breakpoint to a 230 kb fragment that belonged to the HMGI-C gene.³⁴ However, this was only shown to be true in a single case of FA, and thus the involvement of this gene was weakly supported at best. Given the technologies of the 1990s, the approaches available to understand the genetics of FAs were largely limited to karyotype studies, positional cloning and FISH analysis, which could not search the entire genome efficiently to pinpoint specific genetic loci involved in FAs.

MICROSATELLITE INSTABILITY AND LOSS OF HETEROZYGOSITY

As technological advancements in biological assays emerged towards the later part of the 1990s, Deng *et al* reported that loss of heterozygosity (LOH) was detected in morphologically normal lobules found beside malignant breast tumours, with the most frequent aberration found in chromosome

3p22–25. Of 10 breast carcinomas with this aberration, 6 cases showed the same LOH within normal lobules, suggesting the presence of a tumour suppressor gene at 3p22–25.³⁵ Yet, it was shown by Lizard-Nacol *et al* that in FAs, there was an absence of many genetic alterations such as gene amplification commonly found in breast carcinomas.³⁶ Combining this with previous studies that reported associations between FAs and breast cancer risk,^{37–40} McCulloch *et al* decided to study the incidence of microsatellite instability (MSI) and LOH in FAs and showed that it was low at 11 different loci: TPOX, D3S1514, D3S1675, FAPB, D4S243, D9S254, TH01, D13S289, SCA3, D17S559 and D17S855.⁴¹ Additionally, epithelial cells in a single case of FA in a series of benign breast diseases preceding invasive breast cancer were shown to be negative for LOH at chromosome 3p.⁴² The markers used for this subsequent study included D3S1766, D3S1612, D3S1244, ITIH-1, D3S4102 and LUCA2.1. Later reports, such as by Franco *et al*, also verified that LOH was indeed a rare feature in FAs,⁴³ as opposed to phyllodes tumours that displayed extensive LOH on genome-wide single-nucleotide polymorphism (SNP) array analysis.⁴⁴ Although more specific regions of the genome could be assayed to search for abnormalities, MSI and LOH were features hardly found in FAs, and hence studies focusing on them generated little knowledge in terms of the pathological mechanisms that underlie FAs.

TP53 AND RAS MUTATIONS

The unequivocal importance of the tumour suppressor gene *TP53* is clearly demonstrated in its abrogation in over half of all human sporadic cancers, in spontaneous tumour development in Trp53-knockout mice as well as in patients with Li-Fraumeni syndrome who inherit a mutant *TP53* allele.^{45–46} Over the course of tumour development, sporadic *TP53* mutations may arise, which are quickly followed by LOH, leading to total p53 deficiency. This accelerates both initiation and progression of tumour growth, and tumours displaying a lack of p53 are often characterised by more aggressive characteristics such as local invasiveness and metastatic ability.^{47–50}

TP53 was first examined in FAs over two decades ago by Koutselini *et al* who studied expression of this gene in cytological specimens from both benign and malignant breast tumours using a monoclonal antibody (mAb) that detected p53. In 38 out of 39 cases (97.4%), benign epithelial breast cells obtained from FAs, fibrocystic change as well as smears from nipple discharge stained negative for p53.⁵¹ This was contradictory to a study 4 years later, which showed that one in eight FAs exhibited a non-silent *TP53* mutation.⁵² This was then opposed by Franco *et al* who demonstrated that there was an absence of *TP53* mutations in FAs from women who remained unaffected by breast cancer over an average follow-up period of 10 years.⁴³ Franco *et al* went on to demonstrate that *TP53* gene mutations were absent in FAs, regardless of whether or not they were diagnosed in women affected by breast carcinoma.⁵³ Lastly, the findings by Schneider *et al* showed that p53 mRNA and protein levels in FAs were higher compared with adjacent normal breast tissue in women of reproductive age, providing further conflicting evidence about p53 in this disease.⁵⁴ Taken together, it can be concluded that while the role of p53 has been thoroughly established in human cancers, its implication in the pathogenesis and development of FAs was unproven.

Mutations in the *ras* oncogenes *H-ras*, *K-ras* and *N-ras* have long been known to contribute towards uncontrolled cell proliferation ultimately leading to tumour development.⁵⁵ As small GTPases, *ras* proteins alternate between two states: the inactive

guanosine diphosphate-bound and the active guanosine triphosphate (GTP)-bound conformations.^{56–57} They regulate cell fates through the coupling of receptor activation with downstream effector pathways that elicit cellular responses such as increased proliferation, differentiation and survival.^{58–60}

Only 2 out of 10 (20%) FAs were scored positively when stained using a *ras*-sensitive mAb, as compared with 19 out of 30 (63.3%) infiltrating ductal carcinomas. Interestingly, these two were multiple FA cases.⁶¹ Using a combination of PCR amplification and allele-specific oligonucleotide hybridisation techniques, Kumar *et al* found that out of seven rat FAs induced by neonatal exposure to nitrosomethylurea, *H-ras* was mutated in three of them (42.9%), while *K-ras* was mutated in one case (14.3%).⁶² Although FAs exhibited *ras* oncogene activation, it was proven to be within the basal levels of *ras* activation in normal breast, brain and peripheral nerve tissues.^{63–65} On a related note, DNA microarray analysis revealed that the Ras homologue enriched in brain (RHEB) protein, also involved in cell growth and differentiation, was expressed at levels more than twofold higher in FAs than in normal tissues. The average RHEB:β-actin ratio of FAs was found to be two times greater than that in infiltrating ductal carcinomas, suggesting RHEB immunohistochemical staining as a possible secondary diagnostic tool to distinguish between the two.⁶⁶ However, further supporting literature on the role of RHEB in FAs is scant. In summary, the *ras* oncogenes and related homologues have not been shown to play consequential roles in FAs.

HORMONE INVOLVEMENT

Hormones such as oestrin, now more commonly known as oestrogen, were implicated in the development of breast lesions including gynaecomastia and FAs as early as over 80 years ago.⁶⁷ When nine benign breast conditions including eight fibrocystic lesions and one FA were processed into slices and cultured in medium, the addition of prolactin (PRL) caused increased DNA synthesis as observed by a higher rate of incorporation of radioactive thymidine, elegantly showing PRL as a possible mitogenic stimulant in FAs.⁶⁸ This stood in contrast to a study by Manni *et al* using the soft agar clonogenic assay: the only case of FA included in the study did not respond to PRL administration, resulting in no colony formation.⁶⁹ This was supported by the findings of Kumar *et al* who observed that 2 cases of FA stained negative for tissue-bound PRL, as opposed to 14 out of 15 cases of apocrine metaplasia of the breast, which exhibited strong PRL staining.⁷⁰

Compared with normal breast tissue, FAs were also rigorously proven to possess a higher expression level of the prolactin receptor (PRLR) gene by multiple methods including quantitative PCR, immunocytochemistry and immunohistochemistry.^{71–73} As the role of PRL receptors became gradually established in FAs, it started a hunt for mutations in the gene encoding these transmembrane proteins.

In a study involving 74 patients with multiple fibroadenoma (MFA) and 170 control subjects, Bogorad *et al* identified four patients harbouring a heterozygous SNP in exon 6 of the PRLR gene, which resulted in a substitution mutation changing isoleucine to leucine at position 146 of the PRLR extracellular domain. This single mutation was enough to confer constitutive activity to the PRLR variant, confirmed by PRL-independent functions of PRLR tyrosine phosphorylation, activation of signal transducer and activator of transcription 5 signalling, transcriptional activity towards a PRL-responsive reporter gene, and cell proliferation and protection from cell death.⁷⁴ A further

Figure 1 Fibroadenoma that on genomic analysis was found to have a single *MED12* frameshift deletion. The fibroadenoma shows circumscribed borders and both pericanalicular and intracanalicular growth patterns (A). Higher magnification (B) shows hyalinised stroma around epithelial elements that are stretched into arc-like formations.

study of 95 women with MFA by Courtillot *et al* from the same research group reported another SNP in exon 5 that encoded the amino acid substitution of isoleucine to valine at position 76 of the PRLR. However, in their study cohort, only 15% of the MFA cases harboured PRLR that showed basal activity in vitro, raising questions about whether PRLR was widely involved in the aetiology of MFAs and whether it is a relevant inhibitory target for these patients.⁷⁵

The third and definitive study by this group involving another 71 women affected by MFA provided conclusive clinical evidence that gain-of-function PRL receptor variants PRLR_{T146L} and PRLR_{T76V} were indeed not associated with MFA risk. One gripping point raised was that although these variants' gain-of-function properties were ascertained in reconstituted cell systems, they were not replicable in ex vivo analysis in mammary tissues of mouse or human origin, which suggested that the cellular effect of these variants in vivo was not enough to result in mammary tumour development.⁷⁶

MED12 MUTATIONS

In a landmark study, the use of next-generation sequencing (NGS)-based exome sequencing led to the discovery of highly recurrent Mediator complex subunit 12 (*MED12*) somatic mutations in FAs of the breast. These mutations were found in 4 out of an initial batch of 8 (50%) FA samples, before being further ascertained in another 54 out of 90 (60%) FAs (figure 1). Out of these 98 FAs sequenced, 41 of them (42%) had point mutations in codon 44 of exon 2 of the *MED12* gene.⁷⁷

The results from this paper were met with multiple studies in quick succession globally, confirming that *MED12* mutations are found in up to 65% of FAs, with an overwhelming majority of them being located in codon 44 of exon 2.^{78–80} Laser microdissection studies have also revealed that the *MED12* mutations are limited to the stromal compartment,^{77–81} with mutation status correlating with stromal but not epithelial protein expression.⁸² This is supported by the fact that the histological pattern that most frequently harbours *MED12* mutations is the intracanalicular FA,^{79–81} which exhibits stroma growing into and compressing the epithelial compartment, stretching the ducts into elongated, arc-like formations.¹ Collectively, this suggests that the initiation of FA development may likely be due to a *MED12* mutation

event within the stromal compartment, much like the uterine leiomyoma (UL) in which *MED12* mutations were first found and have been established as integral for UL tumourigenesis.^{83–84} In both FAs and ULs, there is a stark absence of recurrent point mutations other than in *MED12*, which suggests that *MED12* mutations might be sufficient for tumourigenesis in both tumour types. However, although there is high mutation frequency and extremely similar mutational patterns in the *MED12* gene between FAs and ULs, *MED12* mutations were only found in 3 of 41 (7.3%) uterine leiomyosarcomas, a malignant tumour closely related to ULs.⁸⁵ This contrasts against the persistently higher mutation frequency of *MED12* in phyllodes tumours of all grades (although some studies have found a relatively lower frequency of *MED12* mutations in malignant phyllodes tumours), the less innocuous fibroepithelial tumour regarded to be closely related to FAs. This suggests that *MED12* mutations are key drivers behind the spectrum of breast fibroepithelial tumours from benign FAs to phyllodes tumours.⁷⁸

Akin to the uterus, the breast is one of the main target sites for oestrogen,⁸⁶ and given that the Mediator complex has been demonstrated to have cellular interactions with oestrogen receptors α and β ,⁸⁷ *MED12* mutations in intralobular stroma may be implicated in aberrant oestrogen signalling that ultimately leads to FA pathogenesis. A recent study by Lozada *et al*, however, has shown that myxoid FAs lack *MED12* mutations, suggesting that this particular variant may arise through mechanisms independent of *MED12*.⁸⁸ The interaction of cellular *MED12* with Cyclin C has been demonstrated to be dependent on the first 100 amino acids in the primary structure of *MED12*, including those encoded by the entire exon 2. Somatic mutations in *MED12* exon 2 have been shown to adversely affect interactions between *MED12* and Cyclin C-CDK8/CDK19, causing loss of Mediator-associated, Cyclin-dependent kinase (CDK) activity, resulting in RAD51B expression and eventual tumourigenesis.^{89–90} Within the CDK module of Mediator, Cyclin C, CDK8 and *MED12* are three subunits that are exclusively recruited in the presence of strong p21 transcriptional upregulation, which occurs downstream of the p53-dependent cell cycle arrest response.⁹¹ This further suggests that *MED12* exon 2 mutations can be deleterious with regard to p53-dependent cell cycle arrest. Therefore, the tumourigenic impact of somatic *MED12* mutations is likely

Table 1 Key publications on genomic and genetic abnormalities reported in breast fibroadenomas

Calabrese <i>et al</i> 1991 ¹⁵	Stephenson <i>et al</i> 1992 ¹⁷	Cavalli <i>et al</i> 1992 ²⁰	Rohen <i>et al</i> 1993 ¹⁹	Leuschner <i>et al</i> 1994 ¹⁸	Ozisik <i>et al</i> 1994 ³²	Schoenmakers <i>et al</i> 1995 ³³	Staats <i>et al</i> 1996 ³⁴
Lizard-Nacol <i>et al</i> 1995 ³⁶	Deng <i>et al</i> 1996 ³⁵	McCulloch <i>et al</i> 1998 ⁴¹	Euhus <i>et al</i> 1999 ⁴²	Franco <i>et al</i> 2001 ⁴³	Wang <i>et al</i> 2006 ⁴⁴	Hand <i>et al</i> 1984 ⁶¹	Kumar <i>et al</i> 1990 ⁶²
Koutselini <i>et al</i> 1991 ⁵¹	Millikan <i>et al</i> 1995 ⁵²	von Lintig <i>et al</i> 2000 ⁶³	Franco <i>et al</i> 2003 ⁵³	Eom <i>et al</i> 2008 ⁶⁶	Schneider <i>et al</i> 2009 ⁵⁴	Welsch <i>et al</i> 1979 ⁶⁸	Manni <i>et al</i> 1986 ⁶⁹
Kumar <i>et al</i> 1987 ⁷⁰	Touraine <i>et al</i> 1998 ⁷¹	Mertani <i>et al</i> 1998 ⁷²	Gill <i>et al</i> 2001 ⁷³	Bogorad <i>et al</i> 2008 ⁷⁴	Courtillot <i>et al</i> 2010 ⁷⁵	Chakhtoura <i>et al</i> 2016 ⁷⁶	Lim <i>et al</i> 2014 ⁷⁷

Publications are colour coded according to the type reported: red—cytogenetic aberrations; yellow—microsatellite instability and loss of heterozygosity; blue—*TP53* and *ras* mutations; purple—prolactin receptor mutations; green—*MED12* mutations.

Figure 2 Key genomic and genetic abnormalities reported in breast fibroadenomas. Findings are colour coded according to the type reported: red—cytogenetic aberrations; yellow—microsatellite instability and loss of heterozygosity; blue—*TP53* and *ras* mutations; purple—prolactin receptor mutations; green—*MED12* mutations. FA, fibroadenoma; HMGI-C, high mobility group protein; LOH, loss of heterozygosity; mAb, monoclonal antibody; qPCR, quantitative PCR; RHEB, Ras homology enriched in brain; SNP, single-nucleotide polymorphism.

Figure 3 Cellular fibroadenoma at low magnification (A) shows a typical appearance of a proliferation of both epithelial and stromal components. At medium (B) and high (C) magnifications, the increased stromal cellularity is observed. Cellular fibroadenomas may be difficult to distinguish from benign phyllodes tumours when there is a predominant intracanalicular growth pattern.

to be multifaceted, involving aberrant oestrogen signalling as well as attenuated cell cycle arrest, and thus warrants further investigations to better understand its full scope.

OTHER MUTATIONS

Previously, somatic missense mutations in *RARA* were only reported in acute promyelocytic leukaemia that was refractory to treatment, as well as in other solid tumours at low frequencies below 5%.⁹² Targeted deep sequencing of 21 FAs revealed mutations in the retinoic acid receptor alpha (*RARA*) gene to be highly clustered within the region that encodes the ligand-binding domain, in 3 (14.3%) of these cases. On top of this, missense mutations were detected in *FLNA*, *PCLO*, *CHD8* and *ROS1*, while frameshift mutations were found in *MLL2* and *PCNXL4*.⁹³ Taken together, the identification of recurrent *MED12* and *RARA* mutations within the stromal compartment of FAs points towards clonality. This counters earlier investigations into the clonality of FAs, which used PCR and comparative genomic hybridisation-based approaches to arrive at the conclusion that FAs are polyclonal.^{94–97} This disparity may be attributed to low stromal cellularity within the FAs that may have masked their true clonality, supported by the fact that FAs with high stromal cellularity or ‘phyllodal features’ were demonstrated to be monoclonal.^{98,99}

Among breast fibroepithelial tumours, mutations in *FLNA*, *SETD2*, *MLL2*, *BCOR* and *MAP3K1* were observed to be more common in phyllodes tumours than FAs. In addition, borderline and malignant phyllodes tumours disclosed mutations in *NF1*, *RB1*, *PIK3CA*, *EGFR*, *TP53* and *ERBB4*, which were largely absent in FAs and benign phyllodes tumours.⁹³ These genomic differences may be of potential clinical use in distinguishing FAs from phyllodes tumours, an important separation as there are divergent management approaches. It is postulated that similar mutational profiles exist for cellular FAs, which represent a diagnostic difficulty in their distinction from phyllodes tumours, though specific data on cellular FAs remain pending.

Among breast papillary lesions that may occasionally display slight morphological similarity to fibroepithelial tumours, a high prevalence of *PIK3CA*/*AKT* pathway mutations has been found.¹⁰⁰ There are currently no papers in the literature highlighting frequent mutations of *PIK3CA* and related genes in FAs. On the other hand, studies describing the presence of *MED12* mutations in intraductal papillomas are scant, with one report published this year showing a 0% mutation rate of *MED12* in six intraductal papillomas.¹⁰¹ Furthermore, another recent publication demonstrated tubular, lactating and ductal adenomas to lack mutations in exon 2 of the *MED12* gene, and that ductal

adenomas showed recurrent mutations in *GNAS* and the *PI3K*–*AKT* pathway.¹⁰² Ductal adenomas of the breast have been previously postulated to be closely related to or are in fact sclerosed intraductal papillomas.¹⁰³

CONCLUSION

Knowledge surrounding FA genetics and genomics has progressed rapidly over the last 20 years—from mostly karyotype studies and FISH analyses at the beginning to the NGS-based approaches today. Table 1 and figure 2 show a summary of the genetic and genomic discoveries in FAs. The next step forward will be to capture their genomic features for correlation with clinicopathological features, which will allow us to evaluate whether genomic data may be harnessed as a secondary adjunct to refine diagnostic assessment of fibroepithelial tumours. In particular, distinguishing cellular FA (figure 3) from benign phyllodes tumour is a common diagnostic challenge, which was brought to greater attention by an interobserver study in which consensus among all pathologists was achieved in only 2 (9.5%) of 21 difficult fibroepithelial lesions. This can be attributed to subjective histopathological criteria such as stromal cellularity, stromal condensation, atypia and well-developed leaf-like growths.^{99,104} Studies that sought to differentiate FAs from phyllodes tumours using protein immunohistochemistry and other methods have so far not been able to consistently assist in resolving this diagnostic dilemma.^{105–108} As such, further investigations are warranted to understand whether genomic features can be clinically useful. Finally, a recent study involving massively parallel sequencing analysis and formal clonality analysis of FAs and phyllodes tumours has shown that *MED12* mutations are extremely common among them and that a clonal relationship likely exists between FAs and phyllodes tumours that share identical *MED12* mutations.¹⁰⁹ Therefore, additional studies are also required to shed greater light on the progression of fibroepithelial lesions and whether FAs do constitute the substrate from which phyllodes tumours develop. We are gradually arriving at a more comprehensive understanding of the molecular underpinnings driving the formation of fibroepithelial tumours.

Handling editor Runjan Chetty.

Contributors BNL performed the literature review and wrote the review paper. NDMN took the photographs of the fibroadenoma cases harbouring *MED12* mutations. AAT, JYHL, CSL and BTT edited the manuscript. PHT coordinated the above and edited the manuscript.

Competing interests None declared.

Provenance and peer review Not commissioned; externally peer reviewed.

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2018. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- Tan PH, Sahin AA. *Atlas of differential diagnosis in breast pathology*. New York: Springer Science, Business Media LLC, 2017.
- Krishnamurthy S, Ashfaq R, Shin HJ, et al. Distinction of phyllodes tumor from fibroadenoma. *Cancer* 2000;90:342–9.
- Fine RE, Whitworth PW, Kim JA, et al. Low-risk palpable breast masses removed using a vacuum-assisted hand-held device. *Am J Surg* 2003;186:362–7.
- Greenberg R, Skornick Y, Kaplan O. Management of breast fibroadenomas. *J Gen Intern Med* 1998;13:640–5.
- Bernardes JR, Seixas MT, Lima GR, et al. The effect of tamoxifen on PCNA expression in fibroadenomas. *Breast J* 2003;9:302–6.
- Yu JH, Kim MJ, Cho H, et al. Breast diseases during pregnancy and lactation. *Obstet Gynecol Sci* 2013;56:143–59.
- Loda M, Mucci LA, Mittelstadt ML, et al. *Pathology and epidemiology of cancer*. Switzerland: Springer International Publishing, 2017.
- McDivitt RW, Stevensm JA, Lee NC, et al. Histologic types of benign breast disease and the risk for breast cancer. *Cancer* 1992;69:1408–14.
- Dupont WD, Page DL, Parl FF, et al. Long-term risk of breast cancer in women with fibroadenoma. *N Engl J Med* 1994;331:10–15.
- Carney JA, Toorkey BC. Myxoid fibroadenoma and allied conditions (myxomatosis) of the breast. A heritable disorder with special associations including cardiac and cutaneous myxomas. *Am J Surg Pathol* 1991;15:713–21.
- Williamson ME, Lyons K, Hughes LE. Multiple fibroadenomas of the breast: a problem of uncertain incidence and management. *Ann R Coll Surg Engl* 1993;75:161–3.
- Iaria G, Pisani F, De Luca L, et al. Prospective study of switch from cyclosporine to tacrolimus for fibroadenomas of the breast in kidney transplantation. *Transplant Proc* 2010;42:1169–70.
- Seo YL, Choi CS, Yoon DY, et al. Benign breast diseases associated with cyclosporine therapy in renal transplant recipients. *Transplant Proc* 2005;37:4315–9.
- Alkhunaizi AM, Ismail A, Yousif BM. Breast fibroadenomas in renal transplant recipients. *Transplant Proc* 2004;36:1839–40.
- Calabrese G, Di Virgilio C, Cianchetti E, et al. Chromosome abnormalities in breast fibroadenomas. *Genes Chromosomes Cancer* 1991;3:202–4.
- Fletcher JA, Pinkus GS, Weidner N, et al. Lineage-restricted clonality in biphasic solid tumors. *Am J Pathol* 1991;138:1199–207.
- Stephenson CF, Davis RI, Moore GE, et al. Cytogenetic and fluorescence in situ hybridization analysis of breast fibroadenomas. *Cancer Genet Cytogenet* 1992;63:32–6.
- Leuschner E, Meyer-Bolte K, Caselitz J, et al. Fibroadenoma of the breast showing a translocation (6;14), a ring chromosome and two markers involving parts of chromosome 11. *Cancer Genet Cytogenet* 1994;76:145–7.
- Rohen C, Bonk U, Staats B, et al. Two human breast tumors with translocations involving 12q13–15 as the sole cytogenetic abnormality. *Cancer Genet Cytogenet* 1993;69:68–71.
- Cavalli LR, Rogatto SR, Cavalli JJ, et al. Characterization of chromosome alterations in fibroadenomas of the breast. *Cancer Genet Cytogenet* 1992;63:156.
- Sreekantaiah C, Leong SP, Karakousis CP, et al. Cytogenetic profile of 109 lipomas. *Cancer Res* 1991;51:422–33.
- Nibert M, Heim S. Uterine leiomyoma cytogenetics. *Genes Chromosomes Cancer* 1990;2:3–13.
- Sandros J, Stenman G, Mark J. Cytogenetic and molecular observations in human and experimental salivary gland tumors. *Cancer Genet Cytogenet* 1990;44:153–67.
- Ferti-Passantonopoulou AD, Panani AD. Common cytogenetic findings in primary breast cancer. *Cancer Genet Cytogenet* 1987;27:289–98.
- Geleick D, Müller H, Matter A, et al. Cytogenetics of breast cancer. *Cancer Genet Cytogenet* 1990;46:217–29.
- Pandis N, Heim S, Bardi G, et al. Whole-arm t(1;16) and i(1q) as sole anomalies identify gain of 1q as a primary chromosomal abnormality in breast cancer. *Genes Chromosomes Cancer* 1992;5:235–8.
- Pandis N, Heim S, Bardi G, et al. Chromosome analysis of 20 breast carcinomas: cytogenetic multiclone and karyotypic-pathologic correlations. *Genes Chromosomes Cancer* 1993;6:51–7.
- Thompson F, Emerson J, Dalton W, et al. Clonal chromosome abnormalities in human breast carcinomas. I. Twenty-eight cases with primary disease. *Genes Chromosomes Cancer* 1993;7:185–93.
- Lu YJ, Xiao S, Yan YS, et al. Direct chromosome analysis of 50 primary breast carcinomas. *Cancer Genet Cytogenet* 1993;69:91–9.
- Larsson C, Byström C, Skoog L, et al. Genomic alterations in human breast carcinomas. *Genes Chromosomes Cancer* 1990;2:191–7.
- Borg A, Zhang QX, Olsson H, et al. Chromosome 1 alterations in breast cancer: allelic loss on 1p and 1q is related to lymphogenic metastases and poor prognosis. *Genes Chromosomes Cancer* 1992;5:311–20.
- Ozisk YY, Meloni AM, Stephenson CF, et al. Chromosome abnormalities in breast fibroadenomas. *Cancer Genet Cytogenet* 1994;77:125–8.
- Schoenmakers EF, Wanschura S, Mols R, et al. Recurrent rearrangements in the high mobility group protein gene, HMGI-C, in benign mesenchymal tumours. *Nat Genet* 1995;10:436–44.
- Staats B, Bonk U, Wanschura S, et al. A fibroadenoma with a t(4;12) (q27;q15) affecting the HMGI-C gene, a member of the high mobility group protein gene family. *Breast Cancer Res Treat* 1996;38:299–303.
- Deng G, Lu Y, Zlotnikov G, et al. Loss of heterozygosity in normal tissue adjacent to breast carcinomas. *Science* 1996;274:2057–9.
- Lizard-Nacol S, Lidereau R, Collin F, et al. Benign breast disease: absence of genetic alterations at several loci implicated in breast cancer malignancy. *Cancer Res* 1995;55:4416–9.
- Krieger N, Hiatt RA. Risk of breast cancer after benign breast diseases. Variation by histologic type, degree of atypia, age at biopsy, and length of follow-up. *Am J Epidemiol* 1992;135:619–31.
- Carter CL, Corle DK, Micozzi MS, et al. A prospective study of the development of breast cancer in 16,692 women with benign breast disease. *Am J Epidemiol* 1988;128:467–77.
- Hutchinson WB, Thomas DB, Hamlin WB, et al. Risk of breast cancer in women with benign breast disease. *J Natl Cancer Inst* 1980;65:13–20.
- Levi F, Randimbison L, Te VC, et al. Incidence of breast cancer in women with fibroadenoma. *Int J Cancer* 1994;57:681–3.
- McCulloch RK, Sellner LN, Papadimitrou JM, et al. The incidence of microsatellite instability and loss of heterozygosity in fibroadenoma of the breast. *Breast Cancer Res Treat* 1998;49:165–9.
- Euhus DM, Maitra A, Wistuba II, et al. Loss of heterozygosity at 3p in benign lesions preceding invasive breast cancer. *J Surg Res* 1999;83:13–18.
- Franco N, Picard SF, Mege F, et al. Absence of genetic abnormalities in fibroadenomas of the breast determined at p53 gene mutations and microsatellite alterations. *Cancer Res* 2001;61:7955–8.
- Wang ZC, Buraimoh A, Iglehart JD, et al. Genome-wide analysis for loss of heterozygosity in primary and recurrent phyllodes tumor and fibroadenoma of breast using single nucleotide polymorphism arrays. *Breast Cancer Res Treat* 2006;97:301–9.
- Vousden KH, Prives C. Blinded by the light: the growing complexity of p53. *Cell* 2009;137:413–31.
- Brady CA, Attardi LD. p53 at a glance. *J Cell Sci* 2010;123:2527–32.
- Lang GA, Iwakuma T, Suh YA, et al. Gain of function of a p53 hot spot mutation in a mouse model of Li-Fraumeni syndrome. *Cell* 2004;119:861–72.
- Song H, Hollstein M, Xu Y. p53 gain-of-function cancer mutants induce genetic instability by inactivating ATM. *Nat Cell Biol* 2007;9:573–80.
- Fearon ER, Vogelstein B. A genetic model for colorectal tumorigenesis. *Cell* 1990;61:759–67.
- Malkin D, Li FP, Strong LC, et al. Germ line p53 mutations in a familial syndrome of breast cancer, sarcomas, and other neoplasms. *Science* 1990;250:1233–8.
- Koutselini H, Malliri A, Field JK, et al. p53 expression in cytologic specimens from benign and malignant breast lesions. *Anticancer Res* 1991;11:1415–9.
- Millikan R, Hulka B, Thor A, et al. p53 mutations in benign breast tissue. *J Clin Oncol* 1995;13:2293–300.
- Franco N, Arnould L, Mege F, et al. Comparative analysis of molecular alterations in fibroadenomas associated or not with breast cancer. *Arch Surg* 2003;138:291–5.
- Schneider L, Branchini G, Cericatto R, et al. Gene and protein expression of p53 and p21 in fibroadenomas and adjacent normal mammary tissue. *Endocrine* 2009;35:118–22.
- Bos JL. Ras oncogenes in human cancer: a review. *Cancer Res* 1989;49:4682–9.
- Boguski MS, McCormick F. Proteins regulating Ras and its relatives. *Nature* 1993;366:643–54.
- Donovan S, Shannon KM, Bollag G. GTPase activating proteins: critical regulators of intracellular signaling. *Biochim Biophys Acta* 2002;1602:23–45.
- Bourne HR, Sanders DA, McCormick F. The GTPase superfamily: a conserved switch for diverse cell functions. *Nature* 1990;348:125–32.
- Bourne HR, Sanders DA, McCormick F. The GTPase superfamily: conserved structure and molecular mechanism. *Nature* 1991;349:117–27.
- Vetter IR, Wittinghofer A. The guanine nucleotide-binding switch in three dimensions. *Science* 2001;294:1299–304.
- Hand PH, Thor A, Wunderlich D, et al. Monoclonal antibodies of predefined specificity detect activated ras gene expression in human mammary and colon carcinomas. *Proc Natl Acad Sci U S A* 1984;81:5227–31.
- Kumar R, Sukumar S, Barbacid M. Activation of ras oncogenes preceding the onset of neoplasia. *Science* 1990;248:1101–4.
- von Lintig FC, Dreiling AD, Varki NM, et al. Ras activation in human breast cancer. *Breast Cancer Res Treat* 2000;62:51–62.
- Guha A, Lau N, Huvar I, et al. Ras-GTP levels are elevated in human NF1 peripheral nerve tumors. *Oncogene* 1996;12:507–13.
- Guha A, Feldkamp MM, Lau N, et al. Proliferation of human malignant astrocytomas is dependent on Ras activation. *Oncogene* 1997;15:2755–65.

- 66 Eom M, Han A, Yi SY, *et al.* RHEB expression in fibroadenomas of the breast. *Pathol Int* 2008;58:226–32.
- 67 Geschickter CF, Lewis D, Hartman CG. Tumors of the breast related to the oestrin hormone. *Am J Cancer* 1934;21:828–59.
- 68 Welsch CW, Dombroske SE, McManus MJ, *et al.* Effect of human, bovine and ovine prolactin on DNA synthesis by organ cultures of benign human breast tumours. *Br J Cancer* 1979;40:866–71.
- 69 Manni A, Wright C, Davis G, *et al.* Promotion by prolactin of the growth of human breast neoplasms cultured in vitro in the soft agar clonogenic assay. *Cancer Res* 1986;46:1669–72.
- 70 Kumar S, Mansel RE, Jasani B. Presence and possible significance of immunohistochemically demonstrable prolactin in breast apocrine metaplasia. *Br J Cancer* 1987;55:307–9.
- 71 Touraine P, Martini JF, Zafrani B, *et al.* Increased expression of prolactin receptor gene assessed by quantitative polymerase chain reaction in human breast tumors versus normal breast tissues. *J Clin Endocrinol Metab* 1998;83:667–74.
- 72 Mertani HC, Garcia-Caballero T, Lambert A, *et al.* Cellular expression of growth hormone and prolactin receptors in human breast disorders. *Int J Cancer* 1998;79:202–11.
- 73 Gill S, Peston D, Vonderhaar BK, *et al.* Expression of prolactin receptors in normal, benign, and malignant breast tissue: an immunohistological study. *J Clin Pathol* 2001;54:956–60.
- 74 Bogorad RL, Courtillot C, Mestayer C, *et al.* Identification of a gain-of-function mutation of the prolactin receptor in women with benign breast tumors. *Proc Natl Acad Sci U S A* 2008;105:14533–8.
- 75 Courtillot C, Chakhtoura Z, Bogorad R, *et al.* Characterization of two constitutively active prolactin receptor variants in a cohort of 95 women with multiple breast fibroadenomas. *J Clin Endocrinol Metab* 2010;95:271–9.
- 76 Chakhtoura Z, Laki F, Bernadet M, *et al.* Gain-of-function prolactin receptor variants are not associated with breast cancer and multiple fibroadenoma risk. *J Clin Endocrinol Metab* 2016;101:4449–60.
- 77 Lim WK, Ong CK, Tan J, *et al.* Exome sequencing identifies highly recurrent MED12 somatic mutations in breast fibroadenoma. *Nat Genet* 2014;46:877–80.
- 78 Lien HC, Huang CS, Yang YW, *et al.* Mutational analysis of MED12 exon 2 in a spectrum of fibroepithelial tumours of the breast: implications for pathogenesis and histogenesis. *Histopathology* 2016;68:433–41.
- 79 Pfarr N, Kriegsmann M, Sinn P, *et al.* Distribution of MED12 mutations in fibroadenomas and phyllodes tumors of the breast—implications for tumor biology and pathological diagnosis. *Genes Chromosomes Cancer* 2015;54:444–52.
- 80 Piscuoglio S, Murray M, Fusco N, *et al.* MED12 somatic mutations in fibroadenomas and phyllodes tumours of the breast. *Histopathology* 2015;67:719–29.
- 81 Mishima C, Kagara N, Tanei T, *et al.* Mutational analysis of MED12 in fibroadenomas and phyllodes tumors of the breast by means of targeted next-generation sequencing. *Breast Cancer Res Treat* 2015;152:305–12.
- 82 Tan WJ, Chan JY, Thike AA, *et al.* MED12 protein expression in breast fibroepithelial lesions: correlation with mutation status and oestrogen receptor expression. *J Clin Pathol* 2016;69:858–65.
- 83 Mäkinen N, Mehine M, Tolvanen J, *et al.* MED12, the mediator complex subunit 12 gene, is mutated at high frequency in uterine leiomyomas. *Science* 2011;334:252–5.
- 84 Mäkinen N, Heinonen HR, Sjöberg J, *et al.* Mutation analysis of components of the Mediator kinase module in MED12 mutation-negative uterine leiomyomas. *Br J Cancer* 2014;110:2246–9.
- 85 Kämpjärvi K, Mäkinen N, Kilpivaara O, *et al.* Somatic MED12 mutations in uterine leiomyosarcoma and colorectal cancer. *Br J Cancer* 2012;107:1761–5.
- 86 Zhu BT, Conney AH. Functional role of estrogen metabolism in target cells: review and perspectives. *Carcinogenesis* 1998;19:1–27.
- 87 Kang YK, Guermah M, Yuan CX, *et al.* The TRAP/Mediator coactivator complex interacts directly with estrogen receptors alpha and beta through the TRAP220 subunit and directly enhances estrogen receptor function in vitro. *Proc Natl Acad Sci U S A* 2002;99:2642–7.
- 88 Lozada JR, Burke KA, Maguire A, *et al.* Myxoid fibroadenomas differ from conventional fibroadenomas: a hypothesis-generating study. *Histopathology* 2017;71:626–34.
- 89 Turunen M, Spaeth JM, Keskitalo S, *et al.* Uterine leiomyoma-linked MED12 mutations disrupt mediator-associated CDK activity. *Cell Rep* 2014;7:654–60.
- 90 Kämpjärvi K, Park MJ, Mehine M, *et al.* Mutations in exon 1 highlight the role of MED12 in uterine leiomyomas. *Hum Mutat* 2014;35:1136–41.
- 91 Donner AJ, Szostek S, Hoover JM, *et al.* CDK8 is a stimulus-specific positive coregulator of p53 target genes. *Mol Cell* 2007;27:121–33.
- 92 Zhu HH, Qin YZ, Huang XJ. Resistance to arsenic therapy in acute promyelocytic leukemia. *N Engl J Med* 2014;370:1864–6.
- 93 Tan J, Ong CK, Lim WK, *et al.* Genomic landscapes of breast fibroepithelial tumors. *Nat Genet* 2015;47:1341–5.
- 94 Noguchi S, Motomura K, Inaji H, *et al.* Clonal analysis of fibroadenoma and phyllodes tumor of the breast. *Cancer Res* 1993;53:4071–4.
- 95 Kobayashi S, Iwase H, Kuzushima T, *et al.* Consecutively occurring multiple fibroadenomas of the breast distinguished from phyllodes tumors by clonality analysis of stromal tissue. *Breast Cancer* 1999;6:201–6.
- 96 Kasami M, Vnencak-Jones CL, Manning S, *et al.* Monoclonality in fibroadenomas with complex histology and phyllodal features. *Breast Cancer Res Treat* 1998;50:185–91.
- 97 Kuijper A, Buerger H, Simon R, *et al.* Analysis of the progression of fibroepithelial tumours of the breast by PCR-based clonality assay. *J Pathol* 2002;197:575–81.
- 98 Noguchi S, Yokouchi H, Aihara T, *et al.* Progression of fibroadenoma to phyllodes tumor demonstrated by clonal analysis. *Cancer* 1995;76:1779–85.
- 99 Krings G, Bean GR, Chen YY, *et al.* Fibroepithelial lesions; the WHO spectrum. *Semin Diagn Pathol* 2017;34:438–52.
- 100 Troxell ML, Levine J, Beadling C, *et al.* High prevalence of PIK3CA/AKT pathway mutations in papillary neoplasms of the breast. *Mod Pathol* 2010;23:27–37.
- 101 Konstantinova AM, Vanecek T, Martinek P, *et al.* Molecular alterations in lesions of anogenital mammary-like glands and their mammary counterparts including hidradenoma papilliferum, intraductal papilloma, fibroadenoma and phyllodes tumor. *Ann Diagn Pathol* 2017;28:12–18.
- 102 Volckmar AL, Leichenring J, Flechtenmacher C, *et al.* Tubular, lactating, and ductal adenomas are devoid of MED12 Exon2 mutations, and ductal adenomas show recurrent mutations in GNAS and the PI3K–AKT pathway. *Genes Chromosomes Cancer* 2017;56:11–17.
- 103 Lakhani SR, Ellis IO, Schnitt SJ, *et al.* WHO classification of tumours of the breast. 4th edn. Lyon: IARC Press, 2012.
- 104 Lawton TJ, Acs G, Argani P, *et al.* Interobserver variability by pathologists in the distinction between cellular fibroadenomas and phyllodes tumors. *Int J Surg Pathol* 2014;22:695–8.
- 105 Korcheva VB, Levine J, Beadling C, *et al.* Immunohistochemical and molecular markers in breast phyllodes tumors. *Appl Immunohistochem Mol Morphol* 2011;19:119–25.
- 106 Umekita Y, Yoshida H. Immunohistochemical study of MIB1 expression in phyllodes tumor and fibroadenoma. *Pathol Int* 1999;49:807–10.
- 107 Vilela MH, de Almeida FM, de Paula GM, *et al.* Utility of Ki-67, CD10, CD34, p53, CD117, and mast cell content in the differential diagnosis of cellular fibroadenomas and in the classification of phyllodes tumors of the breast. *Int J Surg Pathol* 2014;22:485–91.
- 108 Kaya R, Pestereli HE, Erdogan G, *et al.* Proliferating activity in differential diagnosis of benign phyllodes tumor and cellular fibroadenomas: is it helpful? *Pathol Oncol Res* 2001;7:213–6.
- 109 Piscuoglio S, Geyer FC, Burke KA, *et al.* Massively parallel sequencing analysis of synchronous fibroepithelial lesions supports the concept of progression from fibroadenoma to phyllodes tumor. *NPJ Breast Cancer* 2016;2:16035.