Gene of the month: *TLE 1*Karen Pinto,¹ Runjan Chetty ²¹Pathology, Kuwait Cancer Control Center, Shuwaikh, Al Asimah, Kuwait²Department of Histopathology, Brighton and Sussex University Hospitals NHS Trust, Brighton, UK

Correspondence to

Professor Runjan Chetty, Department of Histopathology, Brighton and Sussex University Hospitals NHS Trust, Brighton BN2 5BE, UK; runjan.chetty@gmail.com

Accepted 5 December 2020

Published Online First

13 January 2021

ABSTRACT

TLE 1 is the human homologue belonging to a family of four genes and is located on chromosome 9q21. It consists of 19 exons. Although it does not bind directly to DNA, it acts as a repressor of several signalling pathways via transcription factors. *TLE1* protein has several physiological roles in embryogenesis, haematopoiesis, general differentiation, and both neuronal and eye development. Much attention was focused on its expression in the tumour cell nuclei of synovial sarcoma (SS). However, several other soft tissue tumours that do and do not share morphological similarity with SS also display nuclear immunoreactivity for *TLE1*; hence, caution in interpretation is advocated.

GENE STRUCTURE AND PROTEIN

The *Transducin-like enhancer of split 1* (*TLE 1*) gene belongs to a family of four genes, is located on chromosome 9q21 and consists of 19 exons. *TLE1* protein is the human homologue of the *Drosophila* groucho protein. It encodes a non-DNA binding, 770 amino acid transcriptional protein that serves as a co-repressor of other transcription factors and signalling pathways. *TLE1* protein forms homo-oligomers and hetero-oligomers with several proteins, and after binding, it acts to inhibit transcriptional activity, especially in the Wnt signalling pathway where it interacts with β -catenin and T-cell factor.

PHYSIOLOGICAL ROLE

TLE1 protein plays a normal, physiological role in several processes including during embryogenesis, haematopoiesis, epithelial differentiation, segmentation and neuronal/eye development (see [figure 1](#)).

TLE 1 IN DISEASE

Non-cancer

See [figure 2](#).

Inflammation

The nuclear factor-kappa B (NF- κ B) signalling pathway controls the immune response; hence, it plays a role in inflammation. *TLE 1* represses NF- κ B activity. Thus, the loss of *TLE1* results in excessive activation of NF- κ B-mediated inflammatory pathways associated with increased expression of inflammatory cytokines and chemokines in the skin, lung and intestine.¹ Interactions of *TLE1* and nucleotide-binding oligomerisation domain-containing protein 2 (NOD2) are involved with inflammatory bowel disease. *NOD2* gene was the first susceptibility gene identified in Crohn's disease (CD). Nimmo *et al*² suggested that *TLE1* can regulate NOD2 functions. They realised single

nucleotide polymorphisms (SNP) in *TLE1* (intronic SNP rs6559629) result in a *TLE* risk allele which unmasks the mutant NOD2 resulting in CD (ileal phenotype).

Diabetes mellitus (type 2)

TLE1 is a regulator of pancreatic islet β -cell identity. Armour *et al*³ indicated that reduced expression of *TLE1* in islet cells is inversely correlated with α/β cell ratio. Decreased *TLE1* causes an increase in glucagon gene mRNA and misexpression of glucagon in islet cells, which leads to a loss of cellular identity potentially leading to β -cell to α -cell conversion.

Postnatal microcephaly

In postnatal microcephaly, although the child has a normal head size at birth, there is a subsequent deceleration of the head circumference caused by mutations in genes responsible for the development of the forebrain. Mutations in genes like *CASK*, *CDKL5*, *CREBBP*, *EP300*, *FOXG1* and *SLC9A6* are already known, and recently, *TLE1* has been added to the list. A homozygous missense mutation in *TLE1* results in autosomal recessive postnatal microcephaly.^{4 5}

Sheehan syndrome

Sheehan syndrome is postpartum hypopituitarism spawned due to necrosis of the pituitary gland as a result of severe hypotension or shock during or after childbirth. Diri *et al*⁶ studied patients with Sheehan syndrome and concluded that they have abnormal expressions of *TLE1* (and *TLE3*). They suggested this may disrupt differentiation of pituitary cells and lead to a genetic predisposition to Sheehan syndrome.

Role in cancer

See [figure 3](#).

Even though the *TLE1* protein does not bind to DNA directly, it attaches to specific regions of DNA via its association with a variety of transcription factors. For example, *TLE1* binds to the histone, H3, with resultant modification of chromatin structure, thereby allowing *TLE1* to regulate gene expression.

TLE1 exerts its influence in three ways: (1) by downregulating transcriptional activators, (2) by enhancing transcriptional repressors and (3) by conversion of transcriptional activators into repressors.

TLE1 binds to basic helix loop helix (bHLH) proteins and represses target genes in three main pathways:

© Author(s) (or their employer(s)) 2021. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Pinto K, Chetty R. *J Clin Pathol* 2021;**74**:137–140.

Figure 1 Schematic infographic illustrating the mode of action of the *TLE1* gene.

- ▶ *TCF/LEF* (T-cell factor/Lymphoid enhancer factor) in the WNT/β-catenin pathway.
 - ▶ *HES* (hairy and enhancer of split) in the NOTCH pathway.
 - ▶ *NKX* (homeobox protein) in the sonic Hedgehog pathway.
- A dysregulation in any of these can act as a driver in tumourigenesis.⁷

Synovial sarcoma

The *SS18-SSX* gene fusion is specific to synovial sarcoma (SS) and is its primary genetic driver mutation. Su *et al*⁹ determined that *SS18-SSX* forms a functional endogenous complex with activating transcription factor 2 (ATF2) and *TLE1* which represses ATF2 target genes and resulted in abnormal transcriptional activities which drive the malignant transformation in SS. *SS18-SSX* acts as a frame to link these proteins together (although independently through different protein domains) resulting in an *SS18-SSX/TLE1/ATF2* complex which directly binds to the *EGR1* (early growth response 1) promoter. In healthy cells, *EGR1* impairs tumour development and induces expression of tumour suppressors like *p53*, *TGF-β*, *PTEN* and so on. However, when *SS18-SSX/TLE1/ATF2* complex binds to *EGR1*, it represses its transcription and boosts oncogenesis.

Figure 2 An infographic elucidating the role of *TLE1* in disease. NF-κB, nuclear factor-kappa B.

Diagnostic application of *TLE1* immunohistochemistry

From a practical point of view, antibodies have been developed to *TLE1* and its immunohistochemical application in soft tissue sarcoma pathology which has been useful in detecting SS (figure 4). *TLE* immunohistochemistry is by no means specific for S as several other soft tissue lesions have been noted to be positive for *TLE1* antibody: neurofibroma, schwannoma, malignant peripheral nerve sheath tumour, solitary fibrous tumour, rhabdomyosarcoma, leiomyosarcoma, Ewing sarcoma family, high-grade chondrosarcoma and clear cell sarcoma of soft parts.

Thus, judicious application and interpretation of the antibody in soft tissue tumours is advocated. In other words, a spindle cell soft tissue tumour that is *TLE1* positive is not automatically an SS.

T-cell acute lymphoblastic leukaemia

Riz *et al*¹⁰ studied the role of *TLE1* in T-cell acute lymphoblastic leukaemia (T-ALL) with regard to the *TLX1* (T-cell leukaemia homeobox 1) gene. *TLX1* is a transcription regulator which depending on the availability of transcription cofactors can

Figure 3 An infographic elucidating the role and mechanism of action of the *TLE1* gene in oncogenesis. NF-κB, nuclear factor-kappa B.

Figure 4 Image featuring *TLE1* immunohistochemistry (nuclear stain) in synovial sarcoma.

activate or repress gene expression. *TLE1* co-represses NOTCH signalling by establishing negative loops of regulation and mediating NOTCH repressor activity. However, *TLX1-TLE1* interaction causes de-repression of NOTCH-induced genes and activation of downstream transcription regulators. They identified *TLX1/TLE1/NOTCH* network as the cause for developmental arrest and survival of T-ALL. Low expression of the *TLE1* immunostain also indicates a poor prognosis in ALL.¹¹

Acute myeloid leukaemia

TLE1 plays a role in the differentiation of immature haematopoietic cells. At the end of haematopoiesis, *TLE1* causes the repression of target genes from the AML1, Wnt and NOTCH signalling pathways permitting committed progenitors to begin to acquire the characteristic differentiation status of mature haematopoietic cells. *TLE1* binds to AML1 of the AML/CBF α runt domain transcription factor family. However, this binding site is lost when AML1 is translocated with *ETO* (Eight-Twenty one oncoprotein) resulting in an imbalance in the signalling pathway, resulting in the development of AML with *AML1-ETO* fusion. Fraga *et al*¹² found that *TLE1* undergoes promoter CpG island hypermethylation-associated inactivation in haematologic malignancies, such as diffuse large B-cell lymphoma and AML by disrupting cell differentiation and activation growth-suppression pathways.

Glioblastoma

Forkhead box protein G1 (FOXG1) is a transcription factor that promotes the proliferation of progenitor cells in the cerebral cortex. FOXG1 forms complexes with *TLE1* which then repress genes (Cation transport regulator-like protein 1-*CHAC1*) that negatively regulate NOTCH signalling in brain-tumour initiating cells). *CHAC1* acts as a pro-apoptotic factor and is the target for *FOXG1:TLE1* transcription repression complexes which then promotes glioma cell survival.¹³

Lung cancer

In healthy cells, anoikis is a form of programmed cell death that occurs in anchorage-dependent cells when they detach from the surrounding extracellular matrix. Tumour cells have the unique ability to evade this process, grow independently and metastasise. Yao *et al*¹⁴ studied the role of *TLE1* and its interaction with E-cadherin which is a key regulator of epithelial-mesenchymal transition in lung cancer cells. When a healthy cell loses its attachment, E-cadherin expression is transcriptionally induced which then stimulates anoikis. However, in malignant bronchial

epithelial cells, there is an increased expression of *TLE1* which causes suppression of E-cadherin via transcription factor Zinc finger E-box binding homeobox 1 (ZEB1) which confers enhanced anoikis insensitivity and anchorage-independent growth to the malignant cells.

Breast cancer

Because *TLE1* is a 'co-repressor', it does not bind to DNA directly but binds to other DNA-binding transcription factors to form multiprotein complexes. Just like in the lung, *TLE1* plays a role in breast cancer cell anoikis resistance but through the Bit1 anoikis pathway. Mitochondrial Bit forms a complex with AES (amino-terminal enhancer of split) in the cytoplasm leading to apoptosis. However, in breast cancer cells where *TLE1* is over-expressed, *TLE1* sequesters AES in the nucleus, thus preventing the formation of Bit1-AES complexes.¹⁵

Gastric cancer

Long coding RNAs (LncRNA) are a subtype of RNAs that are incapable of protein coding. However, they can bind to proteins, chromosomes and other macromolecules involved in the regulatory network of endogenous RNA which play a role in tumorigenesis by competitively binding to microRNA (miRNA). In gastric cancer, *GIHCG* is the LncRNA that downregulates miR-1281, a miRNA. *TLE1* is the target gene for miR-1281, and they have an inverse relationship. So the downregulation of miR-1281 causes an increased *TLE1* expression in malignant gastric epithelial cells which results in cell proliferation and migration.¹⁶ *TLE1* immunostain expression in gastric cancers is associated with male gender, less frequent lymphatic and perineural invasion, intestinal subtype (Lauren classification), good histologic grade, early pathologic T-stage and longer disease-free survival.¹⁷

Hepatocellular cancer

Hepatitis viral proteins alter host cell miRNA expression. In hepatocellular cancer (HCC), miR-657 plays a pivotal role. In infected cells, viral proteins induce miRNA-657 which directly suppresses *TLE1*. As normal hepatic cells progress from dysplasia to cancer, their *TLE1* mRNA gradually decreases. As outlined previously, *TLE1* and NF- κ B pathways are related. In the liver, NF- κ B modulates basic cellular processes like hepatic apoptosis, proliferation and cancer development. Suppression of *TLE1* activates NF- κ B pathways, which contributes to hepatocellular carcinogenesis.¹⁸

Take home messages

- ▶ *TLE1* belongs to a family of transcription repressors and is located on chromosome 9q21.
- ▶ *TLE1* protein binds to proteins in three pathways: Wnt, Notch and sonic Hedgehog pathways.
- ▶ *TLE1* inhibits transcription activators leading to repression of transcription.
- ▶ *TLE1* overexpression is characteristic of synovial sarcoma, especially those with the (X;18) translocation.
- ▶ *TLE1* is also expressed immunohistochemically in a variety of other soft tissue tumours and epithelial malignancies.

Handling editor Dharendra Govender.

Twitter Runjan Chetty @runjanchetty

Acknowledgements The authors would like to thank Dr Vikram Deshpande, Massachusetts General Hospital and Harvard Medical School, Boston, USA, for the immunohistochemical images of *TLE1*.

Contributors KP and RC contributed equally to concept, collation of data, design and writing of the manuscript.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests RC is a consultant for Deciphex Ltd, Ireland.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

ORCID iD

Runjan Chetty <http://orcid.org/0000-0002-2124-515X>

REFERENCES

- Ramasamy S, Saez B, Mukhopadhyay S, *et al.* Tle1 tumor suppressor negatively regulates inflammation in vivo and modulates NF- κ B inflammatory pathway. *Proc Natl Acad Sci U S A* 2016;113:1871–6.
- Nimmo ER, Stevens C, Phillips AM, *et al.* Tle1 modifies the effects of NOD2 in the pathogenesis of Crohn's disease. *Gastroenterology* 2011;141:972–81.
- Armour SL, Anderson SJ, Richardson SJ, *et al.* Reduced expression of the co-regulator TLE1 in type 2 diabetes is associated with increased islet α -Cell number. *Endocrinology* 2020;161. doi:10.1210/endo/bqaa011. [Epub ahead of print: 01 Apr 2020].
- Cavallin M, Maillard C, Hully M, *et al.* Tle1, a key player in neurogenesis, a new candidate gene for autosomal recessive postnatal microcephaly. *Eur J Med Genet* 2018;61:729–32.
- Seltzer LE, Paciorkowski AR. Genetic disorders associated with postnatal microcephaly. *Am J Med Genet C Semin Med Genet* 2014;166C:140–55.
- Diri H, Sener EF, Bayram F, *et al.* Genetic disorders of pituitary development in patients with Sheehan's syndrome. *Acta Endocrinol* 2016;12:413–7.
- El Beaino M, Jupiter DC, Assi T, *et al.* Diagnostic value of TLE1 in synovial sarcoma: a systematic review and meta-analysis. *Sarcoma* 2020;2020:1–6.
- Su L, Sampaio AV, Jones KB, *et al.* Deconstruction of the SS18-SSX fusion oncoprotein complex: insights into disease etiology and therapeutics. *Cancer Cell* 2012;21:333–47.
- Lubieniecka JM, de Bruijn DRH, Su L, *et al.* Histone deacetylase inhibitors reverse SS18-SSX-mediated polycomb silencing of the tumor suppressor early growth response 1 in synovial sarcoma. *Cancer Res* 2008;68:4303–10.
- Riz I, Hawley TS, Luu TV, *et al.* TLX1 and Notch coregulate transcription in T cell acute lymphoblastic leukemia cells. *Mol Cancer* 2010;9:181.
- Brassesso MS, Pezuk JA, Cortez MA, *et al.* Tle1 as an indicator of adverse prognosis in pediatric acute lymphoblastic leukemia. *Leuk Res* 2018;74:42–6.
- Fraga MF, Berdasco M, Ballestar E, *et al.* Epigenetic inactivation of the Groucho homologue gene TLE1 in hematologic malignancies. *Cancer Res* 2008;68:4116–22.
- Dali R, Verginelli F, Pramatarova A, *et al.* Characterization of a FOXG1:TLE1 transcriptional network in glioblastoma-initiating cells. *Mol Oncol* 2018;12:775–87.
- Yao X, Pham T, Temple B, *et al.* Tle1 inhibits anoikis and promotes tumorigenicity in human lung cancer cells through ZEB1-mediated E-cadherin repression. *Oncotarget* 2017;8:72235–49.
- Brunquell C, Biliran H, Jennings S, *et al.* Tle1 is an anoikis regulator and is downregulated by Bit1 in breast cancer cells. *Mol Cancer Res* 2012;10:1482–95.
- Liu G, Jiang Z, Qiao M, *et al.* Lnc-GIHCG promotes cell proliferation and migration in gastric cancer through miR-1281 adsorption. *Mol Genet Genomic Med* 2019;7:e711.
- Lee J-H, Son M-W, Kim K-J, *et al.* Prognostic and clinicopathological significance of Transducer-Like enhancer of split 1 expression in gastric cancer. *J Gastric Cancer* 2016;16:21–7.
- Zhang L, Yang L, Liu X, *et al.* MicroRNA-657 promotes tumorigenesis in hepatocellular carcinoma by targeting transducin-like enhancer protein 1 through nuclear factor kappa B pathways. *Hepatology* 2013;57:1919–30.